

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/287944996>

Report of the Association Internationale Pour l'Etude Des Argiles (AIPEA) Nomenclature Committee for 1996

Article in *Clays and Clay Minerals* · January 1997

DOI: 10.1180/claymin.1997.032.3.11

CITATIONS

19

READS

105

10 authors, including:



Stephen Guggenheim

University of Illinois at Chicago

122 PUBLICATIONS 4,449 CITATIONS

[SEE PROFILE](#)



Milton L. L. Formoso

Universidade Federal do Rio Grande do Sul

49 PUBLICATIONS 1,386 CITATIONS

[SEE PROFILE](#)



Emilio Galán

Universidad de Sevilla

224 PUBLICATIONS 5,037 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Aproximación multidisciplinar para el estudio de la contaminación y sus efectos en los organismos. Aplicación al Estero Domingo Rubio

[View project](#)



CO2 sequestration [View project](#)

Report of the Association Internationale pour l'Etude des Argiles (AIPEA) Nomenclature Committee for 1996

S. GUGGENHEIM (*Chairman, AIPEA Nomenclature Committee, Department of Geological Sciences, University of Illinois at Chicago, 845 W. Taylor St., Chicago, Illinois 60607, USA*), A. ALIETTI (*Institute di Mineralogia e Petrografia, Universita di Modena, Via s Eufemia 19, 4110 Modena, Italy*), D. C. BAIN (*Macaulay Land Use Research Institute, Craigiebuckler, Aberdeen AB15 8QH, UK*), V. A. DRITS (*Geological Institute of the Russian Academy of Science, 7 Pzhevsky Per, Moscow J-17, Russia*), M. L. L. FORMOSO (*9,500, Ave Bento Gonçalves, Campus do Vale, Instit. of Geosciences, University Federal do Rio Grande do Sul, Porto Alegre - RS - Brazil CEP - 91540-000*), E. GALAN (*Dpto. Cristalografia, Mineralogia y Quimica Agricola, Facultad de Quimica, Universidad de Sevilla, Spain*), W. HUDNALL (*Agronomy Department, Louisiana State University, Baton Rouge, Louisiana, 70803, USA*), H. M. KÖSTER (*Karl-Böhm-Strasse 191, D-85598 Baldham, Germany*), H. PAQUET (*Centre de Sedimentologie et de Géochimie de la Surface, Institut du Geologie, 1 Rue Blessig, 67084 Strasbourg Cedex, France*) and T. WATANABE (*Joetsu University of Education, Joetsu Niigata 943, Japan*).

(Received 4 February 1997)

The most recent report of the AIPEA Nomenclature Committee involved the definition and use of the terms 'clay' and 'clay mineral'; see Guggenheim & Martin (1995; 1996). Additional items acted upon by this committee are reported here.

POLYMORPHISM, POLYTYPISM, AND THE KAOLIN MINERALS

Background

'Polymorphism' is the ability of a given element or compound to crystallize in more than one form, with each form having a distinct crystal structure. 'Polytypism' is defined as the existence of an element or compound in two or more layer-like crystal structures that differ in layer-stacking sequences. The layers need not be crystallographically identical, but they should be similar. In part, polytypism differs from polymorphism in permitting small differences in chemical composition between structures, not to exceed 0.25 atoms per formula unit of any constituent element (see Bailey, 1977, 1980 for additional comments).

In general, polytypes should not receive individual names. Instead, a set of related polytypes

should be designated by a single name followed by a structural symbol suffix that defines the layer stacking differences. Polytype mineral names already in existence that have international acceptance and serve a useful function need not be discarded. Decision on the retention of individual names is the responsibility of the IMA Commission on New Minerals and Mineral Names. (Bailey, 1977, 1980).

Kaolinite, nacrite and dickite have 1:1 layers with close similarities in structure. They are polytypes. Most properly, they should be designated by a single name followed by a structural symbol suffix that defines the layer stacking sequence. In accord with the above procedures, the CIPEA (succeeded by AIPEA) Nomenclature Committee agreed unanimously in 1963 and reconfirmed in 1967 (Brindley, 1967) that kaolinite, nacrite and dickite should be renamed as a single mineral, with symbols expressing the stacking sequence of kaolinite layers. In 1971, the CMS Nomenclature Committee (Bailey *et al.*, 1971) acknowledged these changes in its report, but left open the question of the exact nature of the stacking-sequence symbols pending further study.

Recommendation

The present AIPEA Nomenclature Committee recommends that the names kaolinite, nacrite, and dickite be retained as mineral names, in contrast to previous committee actions. This action is simply a recognition that, after nearly 35 years, it is apparent that the earlier recommendations are not being followed and that the present names are well-established in the literature. Final determination of this matter resides in the IMA Commission.

GROUP AND SUBGROUP NAMES FOR 1:1 PHYLLOSILICATES

Background

In 1980, the AIPEA Nomenclature Committee (Bailey, 1980) recommended a classification scheme of phyllosilicates in which 1:1 and 2:1 layer types are divided into groups (e.g. kaolinite-serpentine) on the basis of charge per formula unit. Subgroups (e.g. kaolinite, serpentine) are based on dioctahedral and trioctahedral characteristics. Each subgroup is further divided into mineral species (e.g. kaolinite, nacrite, dickite, etc. for kaolinite; lizardite, berthierine, amesite, etc. for serpentine). The classification scheme is based not only on details of structure, but it also corresponds to a succession of stages of refinement in identification. It was noted that several 'dual' names exist for which the group, subgroup, or species names were identical, as is the case for kaolinite. The Committee recognized the potential for confusion and stated that "It is anticipated that the precise definitions of the groups and subgroups and their names will evolve and change with time."

The AIPEA Nomenclature Committee (Bailey, 1980) rejected terms such as 'kandite' or 'septechlorite' as alternative group and subgroup

names for kaolinite or serpentine, respectively, and these terms should not be used. Interestingly, the AIPEA Nomenclature Committee used the term 'kaolin' interchangeably with 'kaolinite' group in the text of the report (e.g. Bailey, 1980, p. 86), as did the report of the Clay Minerals Society (CMS) Nomenclature Committee (Martin *et al.*, 1991, Table 1). However, the use of 'kaolin' as the group name was apparently never discussed by either committee.

Studies of the kaolinite-serpentine group minerals (e.g. Ehrenberg *et al.*, 1993) suggest the need for a group name that differs from the species name, since authors use the term 'kaolinite' to refer to any of the three polymorphs, kaolinite, nacrite or dickite. 'Kaolin' was defined (see Kuzvart, 1977) at the Third Kaolin Symposium (1974) as the following: "Kaolin is a rock characterized by a useful content of kaolin (*sic*) minerals". The economic-mineralogy aspects were elaborated by Patterson & Murray (1975): "Kaolin is a clay consisting of substantially pure kaolinite, or related clay minerals, that is naturally or can be beneficiated to be white or nearly white, will fire white or nearly white and is amenable to beneficiation by known methods to make it suitable for use in white ware, paper, rubber, paint and similar uses". In recent years, however, the term 'kaolin' has been used interchangeably with 'kaolinite' group.

Recommendation

It is the recommendation of this committee that the term 'kaolin-serpentine' group be used to designate what was previously referred to as the 'kaolinite-serpentine' group, as shown in Table 1. The Committee believes that little confusion will result in the use of 'kaolin', since the distinction of 'kaolin' as a rock term or as a mineral group name is recognizable by context.

TABLE 1. Recommended use of kaolin and kaolinite in the classification[†] of 1:1 layer silicates

Layer type	Group	Subgroup [‡]	Species [§]
1:1	kaolin-serpentine	kaolin serpentine	kaolinite, dickite, nacrite lizardite, amesite, berthierine

[†] Refer to Martin *et al.* (1991) for a complete classification scheme.

[‡] Octahedral character ('dioctahedral' or 'trioctahedral') may be used also.

[§] Only a few examples are given.

COMMERCIAL NAMES AND SYNTHETIC MATERIAL

Commercially available synthetic material must be explicitly referred to as a commercial name in quotations and, if the name is recorded as a trademark, capitalized. Thus, for a hypothetical hectorite-like material, 'Xyzite', the name of which has been recorded as a trademark, the appropriate reference in a publication should read: 'Xyzite' (a synthetic hectorite-like material, Xyz Manufacturing Co., Chicago, Illinois, USA). Synthetic material should be referred to as 'hectorite-like' and not 'hectorite', because the latter implies a naturally occurring mineral. Because the term 'synthetic' is unambiguous, specifically negates the naturally occurring aspect of the mineral characteristic in question and is commonly used, the use of 'synthetic' as a description modifier to a mineral name (e.g. 'synthetic quartz') is acceptable. The issue of the use of 'synthetic' has recently been discussed by the IMA (Nickel, 1996), and the above usage is consistent with these recommendations.

REFERENCES

- Bailey S.W. (1977) Report of the I.M.A.-I.U.Cr. Joint Committee on Nomenclature. *Am. Miner.* **62**, 411–415.
- Bailey S.W. (1980) Summary of recommendations of AIPEA Nomenclature Committee. *Clay Miner.* **15**, 85.
- Bailey S.W., Brindley G.W., Johns W.D., Martin R.T. & Ross M. (1971) Summary of national and international recommendations on clay mineral nomenclature. *Clays Clay Miner.* **19**, 129–132.
- Brindley G.W. (1967) CIPEA Nomenclature Subcommittee, Minutes of meetings held June 20 and 23 (1966), Jerusalem, Israel. *Proc. Int. Clay Conf. Jerusalem II*, xxvii–xxix.
- Ehrenberg S.N., Aagaard P., Wilson M.J., Fraser A.R. & Duthie D.M.L. (1993) Depth-dependent transformation of kaolinite to dickite in sandstones of the Norwegian Continental shelf. *Clay Miner.* **28**, 325–352.
- Guggenheim S. & Martin R.T. (1995) Definition of clay and clay mineral: Joint report of the AIPEA and CMS Nomenclature Committees. *Clay Miner.* **30**, 257–259.
- Guggenheim S. & Martin R.T. (1996) Reply to the Comment by D.M. Moore on "Definition of Clay and Clay Mineral: Joint Report of the AIPEA and CMS Nomenclature Committees", *Clays Clay Miner.* **44**, 713–715.
- Kuzvart M. (1977) Aspects of kaolin genesis. In: *Proc. 8th Kaolin Sym. and Meeting on Alunite, Madrid-Rome, Madrid, Ministerio de Industria y Energía, K-12*.
- Martin R.T., Bailey S.W., Eberl D.D., Fanning D.S., Guggenheim S., Kodama H., Pevear D.R., Šrodoň J. & Wicks F.J. (1991) Report of the Clay Minerals Society Nomenclature Committee: Revised classification of clay materials. *Clays Clay Miner.* **39**, 333–335.
- Nickel E.H. (1996) Mineral names applied to synthetic substances. *Eur. Miner.* **8**, 461–462.
- Patterson S.H. & Murray H.H. (1975) Clays. Pp. 519–585 in: *Industrial Minerals and Rocks* (S.J. Leford, editor), 4th Ed., New York, Amer. Instit. Mining, Metallurg. Petrol. Eng., Inc.