

The Role of Dissimilatory Fe(III)-Reducing Bacteria in Transformation of Iron Minerals

D. G. Zavarzina

Faculty of Geology, Moscow State University, Vorob'evy gory, Moscow, 119899 Russia; Institute of Microbiology, Russian Academy of Sciences, pr. 60-letiya Oktyabrya 7/2, Moscow, 117811 Russia

e-mail: zavarzinatwo@mail.ru

Received April 24, 2003

Abstract—The possible role of bacteria in the formation and transformation of iron minerals is considered on the basis of the Recent biogeochemical cycle of iron. Dissimilatory Fe(III)-reducing bacteria are the main agents in the reductive part of the cycle. These bacteria are capable of reducing iron oxides and hydroxides to magnetite and siderite. A high degree of crystallinity of the iron oxides is the most important factor preventing bacterial transformation of minerals. The main component that links the oxidative and reductive parts of the cycle is ferrihydrite (a weakly crystalline mineral). Because of its large surface area, it is the optimal acceptor for bacterial iron reduction.

Key words: Dissimilatory Fe(III)-reducing bacteria, ferrihydrite, magnetite, siderite.

INTRODUCTION

Interpretations of the events of the Early Proterozoic should be supported by microbiological data, because bacteria were the main (if not the only) active agents in the Proterozoic biosphere. Such a study usually compares the products of metabolism of recent microorganisms with Precambrian minerals and rocks in the framework of a uniform approach (in this case, uniform bacterial paleoecology, which allows reconstruction of processes that took place in the past). Carrying out a purely paleontological study is an independent method that reveals the presence of fossil microorganisms preserved in various ways in ancient rocks. The presence of iron bacteria (participating in the iron cycle) in the Proterozoic (2 Ga ago) was first established by Barghoorn and Tyler (1965) in the iron formations of Lake Superior. The accumulation of thick sedimentary series of iron rocks that occurred in the Proterozoic and ceased in the Phanerozoic was one of the most important events in the geological history of the Earth. The cessation of this accumulation in the Phanerozoic restricts the use of the comparative method. The reconstruction of the possible role of microorganisms in the Proterozoic environment, which was different from the Recent, can be based on microbiological data. The possible participation of microorganisms in the iron cycle, either by direct precipitation of Fe(III) oxides by iron bacteria or indirect production of O₂ by cyanobacteria with subsequent chemogenic precipitation, has been suggested for some time (Claud, 1980). However, the formation of magnetite, which is the major ore mineral of iron-rich quartzites, remains an unresolved problem. At present, microbiologists have progressed consider-

ably in their understanding of the processes of the formation of magnetite by microorganisms. Because hydrothermal processes are at present considered to be the most likely source of iron in Precambrian beds, magnetite-forming thermophilic microorganisms are particularly important in this respect.

THE BACTERIAL CYCLE OF IRON

Of all the metals involved in biological processes, iron stands out because of the diversity of the physiological and biochemical functions performed by organisms using it. Supposedly, the extremely wide usage of iron by organisms is not due only to its chemical characteristics, but also largely to its wide distribution and availability on Earth (Ehrenreich and Widdel, 1994).

The bacterial cycle of iron is connected to the cycles of C, O, P, and S through minerals that are produced by the metabolism of microorganisms (Fig. 1). A general scheme of the bacterial processes involved in the cycle of iron at neutral pH is shown in Fig. 2, where the stable forms of iron minerals formed both in reactions with bacteria and abiogenically are shown in frames. In natural systems in a neutral and weakly alkaline environment, such as marine sediments and redox zones of chemocline in stratified basins, the iron cycle is combined with the sulfur cycle through the formation of stable sulfides. It is also known that siderite (FeCO₃) and vivianite (Fe₃(PO₄)₂ · 8H₂O) are products of bacterial reduction of iron oxides (Fredrikson *et al.*, 1998). The reductive and oxidative parts of the bacterial iron cycle are closely related (Fig. 2). To understand the influence

BACTERIAL OXIDATION OF IRON

Bacterial oxidation of iron has been known since the mid-19th century, and the formation of sedimentary ores of ferric iron was one of the first problems studied by geological microbiology (see Zavarzin, 1972). The sedimentary environment of thin-layered deposits of iron and manganese in lake sediments was described in detail by Perfil'ev and Gabe (1964). The physiology and ecology of iron bacteria are poorly studied compared to other prokaryotic organisms. This is because the majority of neutrophilic iron-oxidizing bacteria cannot be presently cultivated in the laboratory, and their classification is still based on morphological criteria (Kholodnyi, 1953). These are mostly microaerophils, which develop at very low concentrations of O₂, although some species also grow at atmospheric concentrations of oxygen. Neutrophilic iron bacteria accumulate iron hydroxides produced by two reactions: (1) sorption of colloid iron hydroxides on acidic mucopolysaccharides (Dubinina, 1977) and (2) oxidation of iron by a non-specific peroxide mechanism (Balashova, 1990). Iron deposition of this type has been documented in the sediments of Lake Superior with a characteristic representative of the genus *Eoastrion*. Well-studied acidophilic iron bacteria (e.g., *Leptospirillum ferrooxidans* and *Acidithiobacillus ferrooxidans*) and thermophilic bacteria (e.g., *Acidianus* and *Ferroplasma*) develop in an environment of stable Fe²⁺




Fig. 1. Relationships of the bacterial cycle with the C, O, P, and S cycles.

of biogeochemical factors on the transformation of iron compounds in the reducing environment, it is necessary to take into account the processes occurring in the oxidizing zone.




Fig. 2. Scheme of biogeochemical iron cycle at neutral pH.

mainly by the leaching of sulfides (Karavaiko *et al.*, 1972) with subsequent precipitation of limonite in areas of mixing of acidic and neutral waters. The biogenic oxidation of iron is almost entirely included in the field of stability of hematite in the Eh–pH coordinates (Fig. 3). In their geochemical activity, microorganisms do not contradict the laws of thermodynamics. They mainly affect the kinetics of the process (Zavarzin, 1972) and develop in the area of stability of the product of the oxidative-reductive reaction they perform. Of all the groups of iron-oxidizing microorganisms, only acidophilic bacteria play an active role in the leaching of rocks. The metabolism of other groups of iron-oxidizing bacteria leads to the deposition of iron hydroxides, thus supporting the position of these bacteria in the diagram (Fig. 3).

Although iron bacteria were known as early as the mid-19th century, detailed study of the minerals they produce has been made only relatively recently. Chukhrov *et al.* (1973) studied sediments formed by *Leptothrix ochracea* in the flooded plain of the Yakhroma River and described a new weakly crystalline mineral—ferrihydrite. The crystalline structure of ferrihydrite had been discovered earlier by Towe and Bradley (1967). However, Chukhrov was the first to conduct a detailed study of its properties and structure and described it as a new mineral. The ability of ferrihydrite to transform into hematite or goethite is its most important feature. Chukhrov considered this mineral to be the key form of iron in its transformation in the zone of hypergenesis (*Supergene...*, [1975]). In nature, the formation of ferrihydrite can occur abiogenically in the presence of organic matter, phosphates, or silicates. However, ferrihydrite most often has a biogenic origin (Vodyanitskii and Dobrovol'skii, 1998). The activity of iron bacteria excludes the formation of goethite, lepidocrocite, or delta oxides because the process of oxidation occurs too quickly, which enables only the appearance of the water oxide of iron with the least developed structure (ferrihydrite). The ability of ferrihydrite to transform into hematite leads to rapid solidification of the sediments of iron bacteria, with a loss of the characteristic bacterial structures. Therefore, even recently formed bacterial sediments very quickly lose the features that indicate their microbial origin. Their genesis is, accordingly, difficult to identify.

Formation of ferrihydrite has been recently shown to occur in anaerobic conditions due to oxidation of ferrous iron by anoxygenic phototrophs (Widdel *et al.*, 1993; Ehrenreich and Widdel, 1994) and nitrate reducers (Straub *et al.*, 1996). This discovery suggests the existence of a closed anaerobic cycle of iron, which may have been very important in the Precambrian, given that reducing environments were dominant on Earth at that time.




Fig. 3. Areas of stability of iron compounds in the area of development of major groups of iron-oxidizing and iron-reducing bacteria in Eh–pH coordinates.

BACTERIAL REDUCTION OF IRON

The reduction of iron by bacteria began to attract the attention of scientists much later than the process of biotic oxidation. The ability of microorganisms (mainly bacteria and, to a lesser degree, fungi) to reduce certain metals was established quite early.

The possibility of bacterial reduction of ferric iron in organic-rich conditions (sugars and peptides) was shown as early as 1927 by Halvorson and Stark (see Gabe *et al.*, 1964). Reduction of metals is often necessary to remove the toxic products of metabolism, e.g., hydrogen or hydrogen sulfide. The formation of poorly soluble sulfides by bacteria in the sulfur cycle is a typical example of such reactions.

Up to the end of the 1970s, the problem of the anaerobic cycle of iron had been addressed by soil microbiology and was concerned primarily with reduction of iron in soil and mud by heterotrophic organisms (Duda and Kalakutskii, 1961).

Blakemore (1975) discovered magnetotactic bacteria, which could be orientated in the same direction. It was convincingly shown that these bacteria are oriented in accordance with the Earth's magnetic field. This discovery induced further studies of this group of organisms by not only microbiologists, but also physicists, geologists, chemists, and engineers (Spring and Schliefer, 1995). The presence of magnetite crystals (Fe_3O_4) or greigite (Fe_3S_4) arranged in chains and enclosed in

membranes (Frankel and Blakemore, 1980), which were named magnetosomes, produced an unusual spatial orientation for these microorganisms. The morphology of the crystals proved to be very diverse (from cubical and octahedral to variously shaped hexagonal prisms).

Magnetotactic bacteria have been discovered in marine sediments, soils, and stratified basins and have never been found in the oxygen-saturated, acidic, or thermal habitats (Spring and Schleifer, 1995). These are microaerophilic bacteria requiring a very low level of oxygen (and some bacteria are known to survive in completely anaerobic environments). Supposedly, these organisms need magnetosomes for correct spatial orientation in the narrow zone of the appropriate physicochemical environment (Bazylinsky and Moskowitz, 1998). Their narrow ecological specialization and the possibility of identifying magnetite and greigite of magnetotactic origin due to the high specificity of magnetosomes allow reconstructions of physical, chemical, paleoclimatic, and paleomagnetic conditions for various sedimentary environments.

Magnetotactic bacteria do not use iron in catabolic (exergonic) reactions. However, they affect the primary magnetism of sediments by intracellular formation of magnetite and greigite. Blakemore *et al.* (1985) and Frankel (1986) have suggested that magnetotactic bacteria could play a significant part in the accumulation of magnetite in Precambrian banded iron formations (BIFs). This hypothesis is problematic in various ways. First, despite the recognized role of magnetotactic bacteria in the primary magnetism of sediments, the amount of magnetotactic magnetite accumulated in the deposits is tiny. Second, magnetotactic bacteria are gradient organisms occupying a narrow ecological niche with specific physicochemical conditions that hardly agree with the scale of deposition of BIFs. Finally, the magnetotactic bacteria that are known at present have a relatively narrow phylogenetic affinity. Supposedly, they evolved from several ancestors, which may have included phototrophic, sulfate-reducing, or chemolithotrophic bacteria (Spring and Schleifer, 1995). At the same time, the most ancient representatives of the Bacteria and Archea domains are thermophilic prokaryotes, while no magnetotactic bacteria have been found in thermal environments.

The problem of whether microorganisms can receive energy through iron reduction remained unsolved for a long time. Balashova and Zavarzin (1980) showed the possibility of catabolic reduction of iron by lithotrophs using molecular hydrogen as electron donor. Soon after, a physiological group of dissimilatory iron reducers was discovered. Lovley and Phillips (1986) showed the ability of microorganisms to reduce iron accompanied by oxidation of acetate, butyrate, propionate, ethanol, and methanol, which are the typical metabolic products of heterotrophic microorganisms, i.e., they proved the possibility of complete

oxidation of organic matter to carbon dioxide during iron reduction. The credit for the discovery of magnetite-forming pathways in the biogenic cycle of iron goes to Lovley. In the anaerobic community, hydrogen and acetate are major products of primary anaerobes. The complete decomposition of organic matter depends on the removal of these products by secondary anaerobes using external electron acceptors. While hydrogen is used by many secondary anaerobes, acetate is available to fewer of them, and accumulation of acetate is quite possible. Lovley's studies showed that iron reduction is an energetically favorable reaction for acetate oxidation. *Shewanella putrefaciens* is the first bacterium for which the ability of dissimilatory manganese and iron reduction was shown (Myers and Nealson, 1988). Lovley *et al.* (1993) isolated and described strains of a new microorganism, *Geobacter metallireducens*, capable of catabolic reduction of iron oxides. At present, more than 20 species capable of this are known. Most of them are Proteobacteria (Slobodkin *et al.*, 1999).

The majority of currently known iron reducers are neutrophils with optimum growth at pH = 6.5–7.5. They include strict anaerobes and microaerophils (Slobodkin *et al.*, 1999). It is essential that the range of stability of magnetite (the final reduced product in Eh–pH coordinates) occur at high pH and low Eh values (see Fig. 3), i.e., iron reducers develop in the range of stability of the products of oxidizing-reducing reactions. Apart from iron, most iron reducers are able to reduce manganese, and many iron bacteria are capable of oxidizing it. The geochemistries of iron and manganese in the sedimentary process are closely connected. The isolation of these two elements occurs in the environment of a dominating sulfur cycle: iron is bound in sulfides, while manganese does not form insoluble sulfides.

To understand the biogeochemical function of iron reducers, it is important to show which iron oxides and hydroxides are most intensely involved in processes of anaerobic iron reduction. The most widespread minerals in the zone of hypergenesis include hematite (Fe_2O_3), goethite ($\alpha\text{-FeOOH}$), akagenite ($\beta\text{-FeOOH}$), lepidocrocite ($\gamma\text{-FeOOH}$), ferrihydrite ($5\text{Fe}_2\text{O}_3 \cdot 9\text{H}_2\text{O}$), maghemite ($\gamma\text{-Fe}_2\text{O}_3$), and magnetite (Fe_3O_4). Of these, hematite and goethite are the most thermodynamically stable in aerobic conditions and, therefore, the most widespread in soils and sediments as final products of transformations. It is currently known that goethite can precipitate directly from solution, whereas hematite requires the presence of ferrihydrite as a predecessor. In fact, hematite is formed by the dehydration of ferrihydrite. Lepidocrocite is less widespread in nature than goethite, but its formation is possible by the oxidation of ferrous iron. Thus, the presence of this mineral in soil indicates oxygen deficiency. The presence of carbonates in the medium precludes the formation of lepidocrocite. In this case, the oxidation of the ferrous iron leads to the formation of goethite. The formation of ferrihydrite, as mentioned above, occurs by rapid oxida-

tion or in the presence of compounds prohibiting crystallization processes (organic matter, phosphate, or silica). Apart from spontaneous transformation into hematite, slow transformation by dehydration of ferrihydrite into goethite is possible. This transformation can be considerably accelerated in the presence of organic reducing compounds, e.g., cysteine. In fact, ferrihydrite is a protoxide of iron, which is why it is so widespread. The formation of maghemite is possible either by the oxidation of magnetite or, more likely, by transformation of other oxides, e.g., goethite. In this process, the presence of organic matter is a necessary condition. Hence, maghemite is most widespread in tropical soils (Schwertmann and Cornell, 1991; Vodyanitskii and Dobrovolskii, 1998).

Ottow (1969) showed that the rate of bacterial iron reduction by organotrophs decreases with the growth of crystallinity of the mineral phase in the line $\text{Fe}(\text{OH})_3 > \text{lepidocrocite} > \text{goethite} > \text{hematite}$. These results were completely supported for dissimilatory iron reducers (Lovley, 1987). The large rate of reduction of the least ordered iron minerals is explained by their greater solubility, developed surface, and thermodynamic instability. Arnold *et al.* (1988) have shown that complexing agents, such as chelates or citrates of Fe (III), considerably increase the rate of iron reduction. Another factor that affects the ability of iron reducers to reduce various oxides and hydroxides of iron is the availability of electron donors (Lovley and Phillips, 1986).

Despite the fact that iron reducers, in contrast to sulfate reducers and methanogens, have to reduce iron from poorly soluble oxides and hydroxides, they successfully compete for organic substrates. The level of the partial pressure of hydrogen in the sediment sufficient to develop iron reducers is 3×10^{-7} atm, whereas, for sulfate reducers and methanogens, this value should be at least 2×10^{-6} and 8×10^{-6} atm, respectively. The minimum concentration of acetate required for the development of iron-reducers, sulfate-reducers, and methanogens is 0.5, 2, and 5 mM, respectively (Lovley, 1987), i.e., iron reducers are capable of iron reduction under conditions where sulfido- and methanogenesis would be completely inhibited even if the necessary electron acceptors were present (Lovley and Phillips, 1986). Iron reduction is one of the most beneficial exchange processes for anaerobic microorganisms. Supposedly, it is the availability of iron in various forms rather than the competition with other anaerobic groups for substrates that has constituted a major factor regulating iron reduction in Recent sediments. At the same time, oxidation of iron leading to the formation of the most beneficial electron acceptor (ferrihydrite) is performed by iron bacteria in environments of minimal available oxygen and neutral pH. Hence, ferrihydrite is the key compound through which the connection between the aerobic and anaerobic parts of the bacterial cycle of iron is performed (see Fig. 2).

The most typical reduced iron compounds occurring in the zone of hypergenesis are various iron sulfides, siderite, magnetite, and vivianite. The formation of sulfides is possible through indirect biogenic reduction during the metabolic activity of microorganisms of the sulfur cycle. The rest of the above minerals are formed through the metabolism of iron reducers. Formation of magnetite as a final reduction phase was discovered for the first time by Lovley (1987). Magnetite formed by iron reducers through the reduction of amorphous iron hydroxide had oval or rounded crystals from 10 to 50 nm in diameter, with 90% corresponding to the superparamagnetic (<30 nm) state and the remaining 10% to the monodomain (>30 nm) state (Lovley, 1990). Despite the large amount of monodomain magnetite, the microorganisms were able to produce far more of it than the magnetotactic bacteria. It was shown that *G. metallireducens* produces 5000 times more magnetite than the equivalent biomass of magnetotactic bacteria (Lovley, 1990). This casts doubts on the genesis of monodomain magnetite that occurs in freshwater and marine sediments and in soils and has previously been identified as magnetotactic. In most sediments, magnetite of a typical size occurs in the anaerobic zone, i.e., under conditions that are unfavorable for the ecology of magnetotactic bacteria, which require oxygen for their metabolism. The sediments of Lake Geneva were found to contain superparamagnetic magnetite typical of dissimilatory iron reducers (Gibbs-Eggar *et al.*, 1999). Mössbauer studies of biogenic magnetite formed by thermophilic iron reducers showed that it largely represents thinly dispersed, incompletely formed magnetite. Further aging of the sediment was accompanied by processes of biogenic recrystallization of magnetite leading to the regulation of its structure and increased size of its crystals (Zavarzina, 2001; Chistyakova *et al.*, 2001). Thus, it is very difficult to recognize the biogenic origin of magnetite in ancient sedimentary beds and metamorphic rocks morphologically.

Magnetite is unstable in oxygenated conditions and can transform into hematite or maghemite, although this process is relatively slow. The pseudomorphs of hematite after magnetite (so-called martite structures) are typical of the zone of hypergenesis. The possibility of biogenic oxidation of magnetite into hematite was shown by Brown *et al.* (1997). A community of microorganisms transformed 11% of iron from magnetite into hematite in three weeks. Because two-thirds of iron atoms in magnetite are in a three-valence state, another possible process occurring in Recent sediments and leading to the loss of biogenic material is its further reduction by iron reducers. The possibility of microbial reduction of magnetite was first demonstrated by the example of *S. putrefaciens* (Kostka and Nelson, 1995). Magnetite in direct contact with a bacterium was rapidly reduced. The formation of mineral phases in the process of magnetite reduction was determined by physicochemical conditions, i.e., the concentration of bicarbonate and the presence of phosphorus or organic

compounds (Dong *et al.*, 2000). Thus, both the reduction of magnetite by iron reducers resulting in the formation of siderite or vivianite, depending on the conditions, and relatively rapid bacterial oxidation of magnetite to hematite are possible in natural sediments. When the medium for *G. metallireducens* contained the increased concentration of bicarbonate, siderite was formed in addition to magnetite (Lovley, 1990). A number of subsequent studies have shown the decisive influence of physicochemical conditions on the rate and degree of reduction of various oxides and hydroxides of iron and accompanying mineral phases (Roden and Edmonds, 1997; Fredrikson *et al.*, 1998).

CONCLUSIONS

The biogenic cycle of iron performed by microorganisms of various physiological and phylogenetic groups has been described microbiologically. This cycle is most clearly defined in the conditions of nearly neutral pH and normal temperatures that dominate on the Earth's surface. Iron bacteria, which use iron in catabolic reactions and release weakly crystalline ferrihydrite, are the main agents in the oxidative part of the cycle. Dissimilatory iron reducers, which reduce various oxides and hydroxides of iron to magnetite, siderite, vivianite, or oxides of iron, depending on the physicochemical conditions, are the main agents in the reductive part of the cycle. Ferrihydrite is not only the key compound of the biogenic cycle, because it is the most readily available acceptor for bacterial reduction, it also plays the main part in the formation of the most important iron oxides in the zone of hypergenesis—hematite and goethite. In the course of iron reduction, which is one of the most energetically favorable processes, bacteria are able to carry out complete destruction of organic matter, including acetate, the major compound accumulating in the majority of anaerobic microbial processes. Despite the fact that iron reducers have to reduce poorly soluble minerals of iron, they successfully compete for substrate with two major groups of secondary anaerobes (methanogens and sulfate reducers) because of their high affinity to the substrates. Formation of magnetite as a major reduced product of iron reduction depends on many factors, including the concentration of bicarbonates, humic acids, phosphorous, and the pH of solution. The presence of monodomain biogenic magnetite in soil and sediment strongly affects their total and remaining magnetism. At present, many studies are appearing that point out the decisive role of dissimilatory iron reducers rather than magnetotactic bacteria in the formation of this mineral in the zone of hypergenesis. It is noteworthy that, at present, the anaerobic cycle of iron and processes of iron reduction are among the topics that are most intensely studied by microbiologists and geochemists. There are, however, many problems requiring more detailed examination, including the biochemistry of iron reduction, the kinetics of the forma-

tions of major reduced minerals, and the participation of bacteria in the formation of iron ore. Our studies have shown that one of the key problems is the stabilization of primary products of bacterial reduction in the process of postbiogenic crystallization in diagenesis.

REFERENCES

1. R. G. Arnold, T. J. Dichristina, and M. R. Hoffmann, "Reductive Dissolution of Fe (III) Oxides by *Rseudomonas* sp. 200," *Biotechnol. Bioeng.* **28**, 1081–1096 (1988).
2. V. V. Balashova, "Threadlike Growth and Oxidation of Iron in a Collection Culture of Saprophytic Mycoplasma *Acholeplasma laidlawii*," *Mikrobiol.* **59** (3), 418–423 (1990).
3. V. V. Balashova and G. A. Zavarzin, "Anaerobic Reduction of Ferric Iron by a Hydrogen-Reducing Bacterium," *Mikrobiol.* **48** (5), 773–778 (1979).
4. E. S. Barghoorn and S. A. Tyler, "Microorganisms from the Gunflint Chert," *Science* **147**, 563–577 (1965).
5. D. A. Bazylinski and B. M. Moskowitz, "Microbial Mineralization of Magnetic Iron Minerals: Microbiology, Magnetism, and Environmental Significance," *Rev. Mineral.* **35**, 1–448 (1998).
6. R. B. Blakemore, "Magnetotactic Bacteria," *Science* **190**, 377–379 (1975).
7. R. P. Blakemore, K. A. Short, D. A. Bazylinski, *et al.*, "Microaerobic Conditions Are Required for Magnetite Formation within *Aquaspirillum magnetotactium*," *Geomicrobiol. J.* **4**, 53–71 (1985).
8. D. A. Brown, B. L. Sherriff, and J. A. Sawicki, "Microbial Transformation of Magnetite to Hematite," *Geochim. Cosmochim. Acta* **61** (9), 3341–3348 (1997).
9. N. I. Chistyakova, V. S. Rusakov, D. G. Zavarzina, *et al.*, "Formation of the Magnetotactic Phase by Thermophilic Fe(III)-Reducing Bacteria: Mössbauer Study," *Phys. Metals Metallogr.* **92** (Suppl. 1), S138–S142 (2001).
10. F. V. Chukhrov, B. B. Zvyagin, A. I. Gorshkov, *et al.*, "Ferrihydrite," *Izv. Akad. Nauk SSSR, Ser. Geol.*, No. 4, 23–33 (1973).
11. P. Claud, "Early Biogeochemical Systems," in *Biogeochemistry of Ancient and Modern Environments*, Ed. by P. A. Trudinger *et al.* (Springer-Verlag, Berlin, 1980), pp. 7–27.
12. H. Dong, J. K. Fredrikson, D. W. Kennedy, *et al.*, "Mineral Transformation Associated with Microbial Reduction of Magnetite," *Chem. Geol.* **169**, 299–318 (2000).
13. G. A. Dubinina, "Biology of Iron Bacteria and Their Geochemical Activity," *Doctoral Dissertation in Biology* (Moscow, 1977).
14. V. I. Duda and L. V. Kalakutskii, "On the Role of Microorganisms in the Reduction Processes in Soil: Reduction of Iron by a Pure Culture of *Pseudomonas tralucida*," *Nauch. Dokl. Vyssh. Shkoly, Biol. Nauki*, No. 2, 1961.
15. A. Ehrenreich and F. Widdel, "Anaerobic Oxidation of Ferrous Iron by Purple Bacteria, a New Type of Phototrophic Metabolism," *Appl. Environ. Microbiol.* **60**, 4517–4526 (1994).

16. R. B. Frankel and R. P. Blakemore, "Navigation Compass in Magnetic Bacteria," *J. Magn. Magn. Mater.* **15-18**, 1562-1565 (1980).
17. R. B. Frankel, "Magnetic Skeletons in Davy Jones' Locker," *Nature* **320**, 575 (1986).
18. R. B. Frankel, "Anaerobes Pumping Iron," *Nature* **330**, 208 (1987).
19. J. K. Fredrikson, J. M. Zachara, D. W. Kennedy, *et al.*, "Biogenic Iron Mineralization Accompanying the Dissimilatory Reduction of Hydrous Ferric Oxide by a Groundwater Bacterium," *Geochim. Cosmochim. Acta* **62** (19-20), 3239-3257 (1998).
20. D. R. Gabe, E. P. Troshanov, and E. E. Sherman, "Formation of Manganese-Iron Intercalations in Silt as a Biogenic Process," *The Role of Microorganisms in the Formation of Manganese-Iron Lake Ores* (Nauka, Moscow, 1964), pp. 95-117 [in Russian].
21. Z. Gibbs-Eggar, B. Jude, J. Dominik, *et al.*, "Possible Evidence for Dissimilatory Bacterial Magnetite Dominating the Magnetic Properties of Recent Lake Sediments," *Earth Planet. Sci. Lett.* **168**, 1-6 (1999).
22. G. I. Karavaiko, S. I. Kuznetsov, and A. I. Golomzik, *The Role of Microorganisms in Leaching of Metals from Ores* (Nauka, Moscow, 1972) [in Russian].
23. N. G. Kholodnyi, *Iron Bacteria* (Akad. Nauk SSSR, Moscow, 1953) [in Russian].
24. J. E. Kostka and K. H. Nealson, "Dissolution and Reduction of Magnetite by Bacteria," *Environ. Sci. Technol.* **29**, 2535-2540 (1995).
25. D. R. Lovley, "Organic Matter Mineralization with the Reduction of Ferric Iron: A Review," *Geomicrobiol. J.* **5** (3-4), 375-399 (1987).
26. D. R. Lovley, "Magnetite Formation during Microbial Dissimilatory Iron Reduction," in *Iron Biominerals*, Ed. by R. B. Frankel and R. P. Blakemore (Plenum, New York, 1990), pp. 150-160.
27. D. R. Lovley and E. J. P. Phillips, "Organic Matter Mineralization with the Reduction of Ferric Iron in Anaerobic Sediments," *Appl. Environ. Microbiol.* **51** (4), 683-689 (1986).
28. D. R. Lovley, S. J. Giovannoni, D. C. White, *et al.*, "*Geobacter metallireducens* gen. nov. sp. nov., a Microorganism Capable of Coupling the Complete Oxidation of Organic Compounds to the Reduction of Iron and Other Metals," *Arch. Microbiol.* **159**, 336-344 (1993).
29. C. R. Myers and K. H. Nealson, "Bacterial Manganese Reduction and Growth with Manganese Oxide as Sole Electron Acceptor," *Science* **240**, 1319-1321 (1988).
30. J. C. G. Ottow, "Der Einfluß von Nitrat, Chlorat, Sulfat, Eisenoxidform und Wachstumsbedingungen auf das Ausmaß der bakterielle Eisenreduktion," *Z. Pflanzenern. Bodenk.* **124**, 238-253 (1969).
31. B. V. Perfil'ev and D. R. Gabe, "The Study by the Method of Microbial Landscape of Bacteria Accumulating Manganese and Iron in Bottom Sediments," in *The Role of Microorganisms in the Formation of Iron-Manganese Lake Ores* (Nauka, Moscow, 1964), pp. 16-53 [in Russian].
32. E. E. Roden and J. W. Edmonds, "Phosphate Mobilization in Iron-Rich Anaerobic Sediments: Microbial Fe(III) Oxide Reduction versus Iron-Sulfide Formation," *Arch. Hydrobiol.* **139** (3), 347-378 (1997).
33. U. Schwertmann and R. M. Cornell, *Iron Oxides in the Laboratory: Preparation and Characterization* (VCH, Weinheim, 1991).
34. A. I. Slobodkin, D. G. Zavarzina, T. G. Sokolova, *et al.*, "Dissimilatory Reduction of Inorganic Electron Acceptors by Thermophilic Anaerobic Prokaryotes," *Mikrobiol.* **68** (5), 600-622 (1999).
35. S. Spring and K.-H. Schleifer, "Diversity of Magnetotactic Bacteria," *System. Appl. Microbiol.* **18**, 147-153 (1995).
36. K. L. Straub, M. Benz, B. Schink, *et al.*, "Anaerobic, Nitrate-Dependent Microbial Oxidation of Ferrous Iron," *Appl. Environ. Microbiol.* **62** (4), 1458-1460 (1996).
37. *Supergene Iron Oxides in Geological Processes*, Ed. by N. V. Petrovskaya (Nauka, Moscow, 1975).
38. K. M. Towe and W. F. Bradley, "Mineralogical Constitution of Colloidal 'Hydrous Ferric Oxide'," *J. Colloid Interface Sci.* **24**, 384-392 (1967).
39. Yu. N. Vodyanitskii and V. V. Dobrovol'skii, *Iron Minerals and Heavy Metals in Soils* (Nauka, Moscow, 1998).
40. F. Widdel, S. Schnell, S. Heising, *et al.*, "Ferrous Iron Oxidation by Anoxygenic Phototrophic Bacteria," *Nature* **362**, 834-836 (1993).
41. G. A. Zavarzin, *Lithotrophic Microorganisms* (Nauka, Moscow, 1972).
42. D. G. Zavarzina, "Biogeochemical Factors in the Transformation of Iron in Reducing Environment," Candidate's Dissertation in Geology and Mineralogy (Moscow State Univ., Moscow, 2001).