

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/225879676>

An internally consistent model for the Thermodynamic properties of Fe-Mg-titanomagnetitealuminat spinels

Article in Contributions to Mineralogy and Petrology · February 1991

DOI: 10.1007/BF00321989

CITATIONS

214

READS

116

2 authors:



Richard Sack
OFM Research

83 PUBLICATIONS 6,534 CITATIONS

[SEE PROFILE](#)



Mark S. Ghiorso
OFM Research

187 PUBLICATIONS 11,330 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Trace Element Modeling [View project](#)



Ti3+— and Ti4+—rich Fassaïtes at the birth of the solar system: Thermodynamics and Applications [View project](#)

Erratum

An internally consistent model for the thermodynamic properties of Fe – Mg-titanomagnetite-aluminate spinels

Richard O. Sack¹ and Mark S. Ghiorso²

¹ Department Earth and Atmospheric Sciences, Purdue University, West Lafayette, IN 47907, USA

² Department Geological Sciences, AJ-20, University of Washington, Seattle, WA 98195, USA

Contrib Mineral Petrol (1991) 106:474–505

An unfortunate error occurred in Eqs. (3a) and (3b) on p. 476 of the above article. A factor of one-half was omitted in both equations, which should have read as follows:

$$s_2 = \frac{1}{2} [2 X_{\text{Al}^{3+}}^{\text{oct}} - X_{\text{Al}^{3+}}^{\text{tet}}] \quad (3b)$$

$$s_4 = \frac{1}{2} [2 X_{\text{Fe}^{3+}}^{\text{oct}} - X_{\text{Fe}^{3+}}^{\text{tet}}]. \quad (3c)$$