

The Solubility of Sulphur in Hydrous Rhyolitic Melts

BEATRICE CLEMENTE, BRUNO SCAILLET* AND MICHEL PICHAVANT

INSTITUT DES SCIENCES DE LA TERRE D'ORLÉANS, CNRS, 1A RUE DE LA FÉROLLERIE, F-45071 ORLÉANS CEDEX 2, FRANCE

RECEIVED JULY 5, 2000; ACCEPTED JUNE 23, 2004
ADVANCE ACCESS PUBLICATION SEPTEMBER 9, 2004

Experiments performed at 2 kbar, in the temperature range 800–1000°C, with fO_2 between $NNO-2.3$ and $NNO+2.9$ (where NNO is the nickel–nickel oxide buffer), and varying amounts of sulphur added to hydrous metaluminous rhyolite bulk compositions, were used to constrain the solubility of sulphur in rhyolite melts. The results show that fS_2 exerts a dominant control on the sulphur solubility in hydrous silicate melts and that, depending on fO_2 , a rhyolitic melt can reach sulphur contents close to 1000 ppm at high fS_2 . At fO_2 below $NNO+1$, the addition of iron to a sulphur-bearing rhyolite magma produces massive crystallization of pyrrhotite and does not enhance the sulphur solubility of the melt. For a given fO_2 , the melt-sulphur-content increases with fS_2 . For fixed fO_2 and fS_2 , temperature exerts a positive control on sulphur solubilities, at least for fO_2 below $NNO+1$. The mole fraction of dissolved sulphur exhibits essentially linear dependence on fH_2S at low fO_2 and, although the experimental evidence is less clear, on fSO_2 at high fO_2 . The minimum in sulphur solubility corresponds to the redox range where both fH_2S and fSO_2 are approximately equal. A thermodynamic model of sulphur solubility in hydrous rhyolite melts is derived assuming that total dissolved sulphur results from the additive effects of H_2S and SO_2 dissolution reactions. The model reproduces well the minimum of sulphur solubility at around $NNO+1$, in addition to the variation of the sulphide to sulphate ratio with fO_2 . A simple empirical model of sulphur solubility in rhyolitic melts is derived, and shows good correspondence between model and observations for high-silica rhyolites.

KEY WORDS: sulphur; solubility; rhyolite; thermodynamics; fO_2 ; fS_2

INTRODUCTION

Sulphur species are the most abundant magmatic volatiles, after H_2O and CO_2 . Sulphur is also a major

component of magmatic solid and melt phases (e.g. pyrrhotite, anhydrite, Fe–S–O liquid). Experiments performed at 1 atm on a wide variety of silicate liquid compositions have shown that sulphur solubility is a function of fO_2 , fS_2 , temperature and melt composition (Fincham & Richardson, 1954; Richardson & Fincham, 1956; Abraham & Richardson, 1960; Abraham *et al.*, 1960; Haughton *et al.*, 1974; Katsura & Nagashima, 1974; Shima & Naldrett, 1975; Buchanan & Nolan, 1979; O'Neill & Mavrogenes, 2002). Experimental studies in which fO_2 varied significantly have shown that a minimum in sulphur solubility occurs at around NNO to $NNO+1$ (NNO being the Ni–NiO solid buffer assemblage) (see Carroll & Webster, 1994). Experimental studies carried out at high pressure (Carroll & Rutherford, 1985, 1987; Luhr, 1990) on hydrous andesitic to dacitic compositions have also shown a minimum in sulphur solubility in this redox range. The effect of temperature on sulphur solubility has been studied, mostly at 1 atm (Richardson & Fincham, 1954; St Pierre & Chipman, 1956; Nagashima & Katsura, 1973; Katsura & Nagashima, 1974; Buchanan *et al.*, 1983). These studies showed that for a constant SO_2 input in 1 atm gas furnaces [or constant absolute value of fS_2 for the study by Buchanan *et al.* (1983)], and a constant absolute value of fO_2 , the sulphur solubility increases with temperature under reducing conditions (below $NNO-1$) and decreases when temperature increases under oxidizing conditions. Luhr (1990) observed a positive dependence of sulphur solubility on temperature under oxidizing conditions at high pressure, but fS_2 was not kept constant in these experiments. In fact, with the exception of Luhr (1990), most high-pressure sulphur solubility experiments have been

*Corresponding author. Fax: 00 33 (0) 2 38 63 64 88.
E-mail: bscaille@cnrs-orleans.fr

performed without the precise and systematic control of fS_2 . This hinders the understanding of the control that fS_2 may have on sulphur solubility in hydrous silicate liquids, which, in turn, renders difficult any thermodynamic modeling of the sulphur solubility in silicate melts.

To address this issue, we have determined the relationship between fO_2 , fS_2 , T (temperature) and X_S (dissolved melt sulphur content) for rhyolitic compositions. We have used rhyolitic compositions primarily because it makes it possible to work at relatively low temperatures ($\leq 1000^\circ\text{C}$), where accurate control of critical parameters (namely T , fO_2 , fH_2O , and fS_2) is comparatively easy. In addition, sulphur solubility data on rhyolitic melts are still scarce (Carroll & Rutherford, 1987; Luhr, 1990; Baker & Rutherford, 1996a; Scaillet *et al.*, 1998), and our study aims at filling this gap. Rhyolitic melts represent important compositional end-members in the evolution of S-rich intermediate to silicic magmas, such as those found in arc settings (e.g. Martel *et al.*, 1998; Scaillet & Evans, 1999). Here, we present a detailed set of sulphur solubility data, acquired at between 800 and 1000°C at 2 kbar and under various fO_2 and fS_2 . We emphasize how fS_2 , fO_2 and T control the solubility of sulphur in rhyolitic melts and we derive both a thermodynamic and an empirical model for the solubility of sulphur in hydrous rhyolite melts.

EXPERIMENTAL TECHNIQUES

Apparatus


All experiments were performed in an internally heated pressure vessel (IHPV), working vertically using Ar–H₂ mixtures as the pressurizing medium (Scaillet *et al.*, 1992). For reducing conditions and at 800 and 930°C , we used a Shaw membrane to measure the fH_2 (Shaw, 1963; Scaillet *et al.*, 1992), connected to a Protaiis gauge (accurate to 0.1 bar). For experiments performed at 1000°C , the experimental fH_2 was measured using Ni–Pd-based solid sensors [Taylor *et al.*, 1992; Pownceby & O'Neill, 1994; see Scaillet & Evans (1999) for additional details]. Sensor capsules were prepared using Pt tubes, lined with ZrO₂ to prevent reaction between Pt and the sensor material. The sensors were sealed. For each experiment, we used two starting sensor compositions with different bulk Ni/Pd ratios. In all cases, both sensors gave nearly identical final compositions, tightly bracketing the experimental fO_2 (or fH_2). Temperature was measured with three sheathed chromel–alumel thermocouples, calibrated at 1 atm against the melting point of NaCl and is accurate to within $\pm 5^\circ\text{C}$. Pressure was monitored by a factory-calibrated Heise gauge, accurate to within 20 bar. All experiments were performed at 2 kbar, with run durations of between 6 and 9 days. Quenching was accomplished by turning off the furnace power, with quench rates of around $100^\circ\text{C}/\text{min}$ between 800 and 250°C .

Starting materials and capsule preparation

The experiments were performed with crushed anhydrous glasses of rhyolitic composition (Table 1), obtained by fusion of a synthetic gel at 1400°C and 1 atm. The compositions correspond to that of the interstitial glasses of the recent Mont Pelée andesites (P1 Plinian eruption, dated at 650 BP; Martel *et al.*, 1998) and that of Mt Pinatubo 1991 dacite (Table 1). Experimental charges were prepared by weighing finely ground glass and elemental sulphur (S) into Au capsules. Between 5 and 6 wt % deionized and distilled H₂O was added, so as to reach conditions slightly below H₂O saturation for a system without sulphur under our experimental conditions. A preliminary study showed that the starting form of sulphur in the experimental charge (as elemental, pyrrhotite or anhydrite) has no detectable effect on the stability of sulphur-bearing phases. The addition of sulphur as anhydrite or pyrrhotite can, however, lead to an increase of the FeO_{tot} and CaO contents of the melt, eventually resulting in the crystallization of a silicate or oxide phase. Therefore, to avoid significant modifications of the starting glass composition as a result of the addition of elements other than sulphur, we used mainly elemental sulphur as starting material. Different concentrations of sulphur were used to generate different fS_2 . In some charges, synthetic pyrrhotites of differing starting compositions (i.e. Fe/S atomic ratio) were used as a sulphur source, with the purpose of monitoring the kinetics of attainment of equilibrium fS_2 . In addition, to investigate the role of iron on sulphur solubility in hydrous silicic melt, some charges were doped with magnetite. The bulk compositions of these magnetite-added charges are listed in Table 1. After quenching, the capsules were re-weighed, to check for leaks. They were then pierced under a binocular microscope, in order to detect a fluid phase, and any H₂S smell was noted. After complete opening, glass chips were mounted in epoxy resin and polished using diamond solutions for SEM and electron microprobe analyses.

Control of fO_2 and fS_2

Oxygen fugacity was varied between NNO–2.3 and NNO+2.9, by varying the fH_2 and fH_2O of the experiment. The sample capsules were always positioned near the Shaw membrane or near the sensor capsules. The fO_2 of any given charge was calculated through the water dissociation reaction



$$K_w = \frac{fH_2 \times fO_2^{1/2}}{fH_2O} \quad (2)$$

We determined the fH_2O in each charge from the water content of the glass, using the model of Burnham

Table 1: Starting compositions (wt %)

	MP natural ¹	MP synthetic ²	MP15VII ³	MP16VII ³	MP17VII ³	MP18XII ³	MP19XII ³	Pinatubo ⁴
SiO ₂	75.65	75.92	70.43	71.60	71.47	72.06	72.06	78.29
Al ₂ O ₃	12.97	13.10	12.15	12.35	12.33	12.43	12.43	12.85
FeO	2.15	2.00	9.09	7.59	7.75	6.99	6.99	1.06
MgO	0.34	0.32	0.29	0.29	0.29	0.29	0.29	0.20
CaO	2.26	2.58	2.39	2.43	2.43	2.45	2.45	1.31
Na ₂ O	4.34	4.14	3.84	3.90	3.93	3.93	3.93	3.29
K ₂ O	1.94	1.94	1.80	1.83	1.83	1.84	1.84	2.85
FeO/(FeO+MgO)	0.86	0.87	0.97	0.96	0.96	0.96	0.96	0.84

The compositions are recalculated to 100% anhydrous. Total iron is given as FeO.

¹Natural interstitial glass composition of Mt Pelée andesite erupted in 1929 (Martel *et al.*, 1998). It also contains 0.26 wt % TiO₂ and 0.08 wt % MnO.

²Dry synthetic glass used in all non magnetite-doped charges.

³Magnetite-doped charges, obtained by mechanical mixture between the dry MP synthetic glass and stoichiometric magnetite.

⁴Dry synthetic glass similar to the Pinatubo 1991 dacite interstitial glass (Scaillet & Evans, 1999). It also contains 0.15 wt % TiO₂.

(1979). For experiments performed with a Shaw membrane, the oxygen fugacity of a given charge was computed from equation (2), knowing the $f\text{H}_2$ of the experiment, the $f\text{H}_2\text{O}$ of the charge and the equilibrium constant of the reaction, K_w (Robie *et al.*, 1979). For experiments with solid sensors, the prevailing $f\text{H}_2$ was first computed from equation (2), using the sensor composition and the model of Pownceby & O'Neill (1994) and assuming that $f\text{H}_2\text{O}$ in the sensor was that of pure H₂O. Then, the $f\text{O}_2$ of the charge was calculated with $f\text{H}_2$ and $f\text{H}_2\text{O}$ from the model of Burnham (1979). It should be noted that the above approach in controlling $f\text{O}_2$ does not require the presence of a fluid phase in the charge. Equilibrium (1) can be either homogeneous, with all components dissolved in the melt, or heterogeneous, with components either in the melt or fluid phases.

The uncertainty in $f\text{O}_2$ calculation depends on the uncertainties in temperature, $f\text{H}_2$ and $f\text{H}_2\text{O}$. Errors arising from uncertainties attached to either $f\text{H}_2$ (± 0.1 bar) or temperature ($\pm 5^\circ\text{C}$) measurements are minor, i.e. less than $0.01 \log f\text{O}_2$. In contrast, the uncertainty attached to $f\text{H}_2\text{O}$ is the main source of error, as it depends both on the uncertainty of the determination of the melt H₂O content, which is, here, estimated using the by-difference method (see below), and on the accuracy of the thermodynamic model used to calculate H₂O fugacities from melt H₂O contents. An error of ± 0.5 wt % in the melt H₂O content, which corresponds to the analytical uncertainty of most charges, translates into an error of $c. 10\%$ in $f\text{H}_2\text{O}$ using the model of Burnham (1979), which, in turn, translates into an

uncertainty of $c. 0.2$ log units in $f\text{O}_2$, if all other parameters are held constant. The error associated with the thermodynamic model is more difficult to evaluate. In this study, all melt compositions are rhyolitic (see below), with SiO₂ contents in the range 75–79 wt % (anhydrous basis) and are, therefore, in a compositional domain that is well covered by existing thermodynamic models of H₂O solubility in silicate melts (see Zhang, 1999). Although it is well known that the Burnham model does not properly account for the actual water species in the melt, a recent compilation by Zhang (1999) has shown that, in terms of bulk water solubility, the Burnham model yields water contents almost identical to those obtained using speciation models (e.g. Stolper, 1982), at least up to 3 kbar. Thus, besides the speciation of water in silicate melt, the use of either model (Burnham- or Stolper-type) makes no difference in terms of $f\text{H}_2\text{O}$ calculations that we are interested in. The Burnham model is based on the thermodynamic properties of pure water, as determined by Burnham *et al.* (1969), which were also used to derive the a and b parameters used in the MRK equation of state (EOS) of H₂O by Holloway (1977). In the updated version of the Stolper model, provided by Zhang (1999), the EOS of Pitzer & Sterner (1994) is used to calculate $f\text{H}_2\text{O}$. As explained below, the MRK-EOS of Holloway (1977) is used for the calculation of $f\text{S}_2$. Therefore, for the sake of internal consistency, the Burnham model has been used in this work. Standard error propagation formalism of analytical or experimental sources of error through equation (2) yields an uncertainty in $f\text{O}_2$ that is always less than 0.6 log units and with an average ($\pm 1\sigma$) of 0.18 log

units for all charges. All experimental fO_2 are referenced to the NNO buffer (Holloway *et al.*, 1992).

The fS_2 was determined from the composition of pyrrhotite, crystallized during the experiment (Froese & Gunter, 1976). The fS_2 of the experiments was thus not fixed by a standard buffer assemblage such as pyrite–pyrrhotite, but, instead, it was a function of the initial amount of sulphur in the charge and of the imposed experimental conditions. By changing the bulk sulphur content of the starting material, we changed the fS_2 of the system. Sulphur fugacities calculated with pyrrhotite composition determined by XRD (see below) are known to within ± 0.2 log units. However, under oxidizing conditions, pyrrhotite is not a stable phase and is replaced either by an Fe–S–O liquid (see below) or by anhydrite (Carroll & Rutherford, 1985, 1987; Luhr, 1990). In our experiments, progressive addition of sulphur leads to saturation of the hydrous silicate liquid with respect to a fluid phase. In such cases (i.e. fluid phase present), we used an MRK equation of state (e.g. Holloway, 1977, 1981, 1987) to determine fS_2 and the fluid phase composition.

The fluid phase can be described in the H–O–S system. By the phase rule, we have

$$v = 2 + C - \phi \quad (3)$$

where C is the number of components, ϕ is the number of phases and v is the degree of freedom (4 in this case). The pressure, the temperature and the fH_2 are fixed in our experiments and we know the fH_2O of each charge. Therefore, the system is invariant and we can calculate the fluid phase composition and, thus, the fS_2 . Note that this allows calculation of the fugacities of all S-bearing species in the fluid, even if none is known.

In this study, we used the MRK equation of state (MRK-EOS) of Holloway (1977):

$$P = \frac{RT}{V - b} - \frac{a}{\sqrt{T}(V^2 + Vb)} \quad (4)$$

where P is the pressure, T the temperature, V the volume, and a and b are two parameters as a function of temperature and/or pressure. For pure phases, a and b can be estimated from the critical pressure (P_c) and temperature (T_c) (Breedveld & Prausnitz, 1973; Ferry & Baumgartner, 1987). Calculation of the fluid phase compositions indicates that the major species under our experimental conditions are H_2O , H_2 , H_2S , SO_2 and S_2 . H_2O is a polar molecule, the parameters a and b for water, taken from Holloway (1981). H_2S and SO_2 are also polar molecules but no P – V – T data are available under our experimental conditions. Therefore, we considered these molecules as non-polar, a and b being determined from P_c and T_c , as given by Shi & Saxena (1992). For S_2 , P_c and T_c are not known and we used the estimated values of Shi & Saxena (1992). The full set of

Table 2: a and b parameters used in the MRK equation of state

	a (atm \times cm ⁶ K ^{0.5} /mole ²)	b (cm ³ /mole)
H ₂ ¹	3.56 \times 10 ⁶	15.15
H ₂ S ²	89.38 \times 10 ⁶	29.89
SO ₂ ²	142.92 \times 10 ⁶	39.46
S ₂ ²	25 \times 10 ⁶	20.57
	a_o	b
H ₂ O ¹	35 \times 10 ⁶	14.6

¹From Holloway (1981).

²From Shi & Saxena (1992).




Fig. 1. Comparison between fS_2 measured by the pyrrhotite composition [$\Delta FFS(Po)$] and fS_2 calculated from an MRK equation of state [$\Delta FFS(V)$].

a and b parameters used in the MRK-EOS is listed in Table 2. The MRK program was checked against the program of Connolly & Cesare (1993) and, for given P , T , fH_2 and fH_2O conditions, both methods were found to agree within 0.2 log fS_2 .

As discussed previously, for reducing conditions ($fO_2 < NNO$), fS_2 was calculated from the pyrrhotite composition. Some pyrrhotite-bearing charges that were saturated with a fluid phase provided a test of the validity of the MRK-EOS approach to compute fS_2 and fluid phase compositions. As shown in Fig. 1, the fS_2 given by the pyrrhotite composition and the fS_2 given by the fluid phase composition are usually found to agree with each other to within 1 log unit for most charges, which is considered encouraging, given our current poor knowledge of the thermodynamic behavior of the H–O–S fluids at magmatic conditions. Charges at an fO_2 above NNO+1 or at 1000°C are those for which the two methods yielded significantly different results (Fig. 1). This misfit could be due, in part, to the modification of

sulphide composition during cooling (see below). Despite this limitation, we conclude that it is possible to use the MRK equation of state to estimate fS_2 of charges lacking pyrrhotite, at least for H_2O -rich fluid compositions. An additional check on the accuracy of the calculated fS_2 was also provided by comparing the H_2O/S ratio of the fluid phase, as calculated from MRK, against that obtained through mass balance. In the latter, the amount of S and H_2O left over for the fluid phase can be determined. Thus, the H_2O/S ratio of the fluid can be obtained and compared with H_2O/S values calculated from the MRK-EOS approach. Pyrrhotite-free charges for which the calculated H_2O/S ratios from both methods failed to converge were not considered for deriving either the thermodynamic or the empirical models of sulphur solubility presented below. Sulphur fugacities calculated using an MRK approach have larger associated uncertainties than those calculated from the pyrrhotite composition (around ± 0.5 log units), largely because of the propagation of the uncertainty associated with the determination of melt water contents.


All values of fS_2 are referenced to the FFS buffer, which corresponds to the equilibrium between iron and troilite:




$$\ln fS_2 = \frac{2}{RT}(-0.2655(P-1) - 35910 + 12.56T) \quad (\text{Froese \& Gunter, 1976}) \quad (6)$$

with T in K, P in bar and R in cal/molK. We extrapolated the data from Froese & Gunter (1976) to high temperature (above 900°C) and, at 2 kbar, the $-\log fS_2$ (FFS) is 9.36 at 800°C , 7.75 at 930°C and 7.02 at 1000°C .

Knowing fH_2 , fS_2 , and fO_2 , the equilibrium values of both fH_2S and fSO_2 were calculated using the following equilibria:



and



using the equilibrium constants of Ohmoto & Kerrick (1977).

Attainment of equilibrium

Kinetic studies performed by Turkdogan & Pearce (1963) on simultaneous oxidation and sulfidation reactions in anhydrous iron-bearing silicate liquids showed that sulfidation is faster than iron oxidation. The corresponding

Table 3: Composition of synthetic pyrrhotites

	Po5	Po7
$d_{(102)}^1$	2.0815	2.0500
$MFeS^2$	0.975	0.920
% at Fe^3	48.77	45.97

¹Distance between inter-reticular 102 plane of hexagonal pyrrhotite determined by XRD.


²Mole fraction of FeS in pyrrhotite from Toulmin & Barton (1964).

³Atomic proportion of Fe in pyrrhotite calculated after Yund & Hall (1969).

two reactions are



Both the sulfidation and oxidation reactions are related through the following equilibrium:



Baker & Rutherford (1996b) and Gaillard *et al.* (2001) have shown that the Fe^{3+}/Fe^{2+} ratio is constant after 24 h run durations for hydrous rhyolitic compositions, equilibrated between 725 and 1150°C , 100 and 2000 bar, and fO_2 between NNO and $MnO-Mn_2O_3$. Given that our run durations are between 6 and 9 days, we can expect that the iron redox ratio is at equilibrium and, consequently, that sulphur solubility is also at equilibrium. The Fe content of the capsules was always found to be below detection (270 ppm), even under reduced conditions. At fO_2 higher than NNO, sulphur-free supra-liquidus glasses have the same iron content as the starting composition. At lower fO_2 , Fe loss occurred but remained negligible. The maximum Fe loss occurred at 1000°C , where the glass has an FeO content of 1.5 wt % (instead of 2 wt %). However, in all cases, Fe concentration profiles toward the glass-capsules contact were flat in the glass, which suggests that equilibrium between melt and capsule was attained with respect to Fe. To check the attainment of the fS_2 equilibrium in some experiments, we added two additional capsules, where 1 wt % sulphur was added as synthetic pyrrhotite (Table 3). Two different pyrrhotites (charges with Po5 and Po7 in Table 4) were synthesized in evacuated SiO_2 -glass tubes (Kullerud, 1971) using iron and sulphur powder (both from Aldrich, 99.99% purity). For the synthesis, the sample was heated rapidly to 400°C , kept at this temperature over 12 h and then held at 700°C for

Table 4: Experimental conditions and run products

Charges	wt % S ¹	aH ₂ O ²	logfO ₂ ³	ΔNNO ⁴	d(10a) ⁵	at % Fe ⁶	log(S ₂ /pO ₂) ⁷	log(S ₂ /V) ⁸	ΔFFS (ppm) ⁹	ΔFFS (V) ¹⁰	GI	PI	Px	Qz	Po	An	V	
P = 2.152 kbar, T = 785°C, $HH_2^{11} = 27.4$ bar, 143 h																		
MP15 I	1.18 (S)	0.81 (0.09)	-15.30 (0.13)	-1.12 (0.10)	2.0688	47.57	-2.14 (0.42)	-0.78 (0.50)	7.45 (0.21)	8.81 (0.25)	94.41 (3.43)	3.52 (1.77)	0.87 (0.51)	—	1.17	—	1.39 (0.50)	
MP15 II	2.29 (S)	0.80 (0.29)	-15.34 (0.32)	-1.16 (0.29)	2.0532	46.23	-0.16 (0.38)	-0.53 (0.57)	9.43 (0.19)	9.06 (0.26)	74.99 (2.89)	14.38 (1.59)	0.95 (0.43)	8.30 (1.59)	1.37	—	3.25 (0.84)	
MP15 III	3.72 (S)	0.37 (0.09)	-16.01 (0.30)	-1.83 (0.22)	2.0481	45.52	0.36 (0.37)	0.03 (0.09)	9.95 (0.19)	9.62 (0.06)	57.64 (2.33)	25.04 (1.27)	0.55 (0.40)	15.41 (8.51)	1.36	—	6.18 (0.36)	
MP15 IV	5.69 (S)	0.35 (0.15)	-16.11 (0.54)	-1.93 (0.39)	2.0449	45.58	0.66 (0.37)	0.05 (0.14)	10.25 (0.18)	9.63 (0.10)	47.82 (2.12)	32.59 (1.31)	0.73 (0.42)	17.90 (1.17)	1.37	—	9.15 (0.44)	
MP15 V	10.62(S)	0.01	-17.65	-3.47	2.0441	45.52	0.73 (0.37)	0.29	10.32 (0.19)	9.88	53.59 (2.84)	26.77 (1.77)	1.89 (0.58)	16.82 (1.35)	0.92	—	15.23 (0.20)	
MP15 VI	1.36 (po5-por7)	1.00	-14.84	-0.93	2.0780	48.23	-3.21 (0.13)	-1.36 (0.30)	6.19 (0.13)	8.03 (0.30)	95.60 (1.21)	4.30 (0.91)	—	—	0.10 (0.16)	—	1.06 (0.78)	
MP15 VII ¹³	3.70 (S)	1.00	-14.84	-0.93	2.0754	48.18	-3.23 (0.11)	—	6.36 (0.22)	—	90.74 (0.56)	—	—	—	9.26	—	—	
MP15 X	1.02 (S)	0.92 (0.09)	-15.19 (0.11)	-1.03 (0.08)	2.0690	47.58	-2.17 (0.42)	-1.12 (0.22)	7.42 (0.21)	8.46 (0.14)	98.90 (1.21)	traces	—	—	1.10 (0.16)	—	3.21 (1.11)	
MP15 XI	1.11 (S)	0.97 (0.03)	-15.14 (0.04)	-0.96 (0.09)	2.0708	47.75	-2.46 (0.43)	-1.60 (0.30)	7.13 (0.21)	7.99 (0.18)	98.89 (1.20)	—	—	—	1.11 (0.11)	—	5.05 (1.15)	
MP15 SN	1.10 (S)	0.79 (0.18)	-15.34 (0.28)	-1.18 (0.20)	2.0702	47.69	-2.35 (0.42)	-0.59 (0.15)	7.24 (0.21)	9.00 (0.31)	96.42 (1.20)	—	0.56 (0.40)	—	1.27 (0.40)	—	2.67 (0.87)	
PM 15 VIII	3.62 (S)	0.78 (0.06)	-15.33 (0.09)	-1.16 (0.07)	2.0490	45.89	0.27 (0.38)	-0.71 (0.23)	9.86 (0.19)	8.88 (0.15)	62.99 (1.86)	20.30 (1.17)	—	15.77 (1.03)	0.94	—	5.23 (0.27)	
P = 1.973 kbar, T = 798°C, $HH_2^{11} = 1.8$ bar, 139 h																		
MP16 I	1.16 (S)	0.88 (0.13)	-12.59 (0.18)	1.30 (0.12)	2.0524	45.16	0.00 (0.15)	1.37 (0.31)	9.39 (0.07)	10.76 (0.19)	53.98 (1.81)	25.40 (0.97)	—	17.88 (1.14)	1.74	—	2.28 (0.48)	
MP16 II	2.98 (S)	0.77 (0.12)	-12.71 (0.19)	1.18 (0.13)	2.0563	46.48	-0.42 (0.15)	1.61 (0.56)	8.97 (0.08)	10.99 (0.26)	41.67 (1.66)	34.02 (0.94)	—	22.40 (1.04)	1.9	traces	5.33 (0.29)	
MP16 III	4.40 (S)	0.75 (0.12)	-12.73 (0.18)	1.16 (0.19)	2.0567	46.51	-0.46 (0.19)	1.66 (0.46)	8.92 (0.10)	11.04 (0.24)	36.30 (1.51)	36.71 (0.88)	—	25.11 (0.97)	1.88	—	7.14 (0.29)	
MP16 IV	6.10 (S)	0.61 (0.09)	-12.91 (0.18)	0.98 (0.13)	2.0562	46.47	-0.41 (0.16)	1.97 (0.17)	8.98 (0.08)	11.35 (0.11)	29.96 (1.30)	41.97 (0.87)	—	28.46 (0.87)	1.9	—	9.30 (0.24)	
MP16 V	10.74 (S)	0.67 (0.12)	-12.90 (0.12)	1.07 (0.16)	2.0570	46.54	-0.50 (0.20)	0.99 (0.12)	8.89 (0.10)	11.35 (0.08)	33.72 (1.50)	40.70 (0.96)	—	23.66 (0.94)	1.91	—	13.64 (0.30)	
MP16 VII ¹³	4.00 (S)	0.86	-12.60	1.29	2.0574	46.58	-0.55 (0.15)	0.99 (0.12)	8.84 (0.80)	10.38 (0.46)	79.20 (1.17)	11.20 (0.84)	—	—	6.80 (0.20)	—	4.48 (0.46)	
MP16 VIII	1.11 (S)	0.98 (0.03)	-12.49 (0.04)	1.41 (0.09)	2.0608	46.87	-1.00 (0.30)	0.67 (0.94)	8.43 (0.10)	10.06 (0.30)	36.30 (1.51)	36.71 (0.88)	—	25.11 (0.97)	1.88	—	7.14 (0.29)	
MP16 IX	1.37 (S)	0.98 (0.02)	-12.48 (0.08)	1.41 (0.02)	2.0583	46.65	-0.70 (0.30)	0.69 (0.63)	8.73 (0.10)	10.08 (0.27)	57.46 (2.02)	21.04 (1.10)	—	20.21 (1.16)	1.29 (0.16)	—	6.64 (0.29)	
PM16 VI	4.39 (S)	0.68 (0.10)	-12.80 (0.18)	1.06 (0.13)	2.0516	46.10	0.08 (0.30)	1.81 (0.28)	9.46 (0.30)	11.20 (0.17)	42.30 (1.15)	30.31 (0.77)	—	26.44 (0.83)	0.95	—	7.22 (0.27)	
P = 1.976 kbar, T = 811°C, $HH_2^{12} = 1$ bar, 150 h																		
MP17 I	1.21 (S)	1.00	-11.78	1.83 (0.20)	2.0546	46.24	-0.13 (0.15)	-0.96	9.08 (0.07)	8.25	85.33 (3.31)	12.64 (1.65)	—	traces	1.86	—	0.47 (0.45)	
MP17 II	2.24 (S)	1.00	-11.78	1.83 (0.20)	2.0525	46.17	0.09 (0.20)	-0.96	9.30 (0.10)	8.25	78.96 (3.06)	15.95 (1.54)	—	3.19 (1.68)	1.9	—	1.86 (0.42)	
MP17 III	4.43 (S)	1.00 (0.01)	-11.79 (0.01)	1.83 (0.20)	2.0549	46.37	-0.17 (0.16)	-1.10 (0.17)	9.04 (0.08)	8.11 (0.11)	78.82 (3.61)	15.64 (1.85)	—	3.82 (1.92)	1.87	traces	4.40 (0.36)	
MP17 IV	6.03 (S)	0.97 (0.04)	-11.82 (0.05)	1.80 (0.09)	2.0572	46.56	-0.42 (0.15)	0.71 (0.88)	8.79 (0.08)	9.91 (0.30)	78.50 (3.62)	15.70 (1.88)	—	3.82 (1.91)	1.87	traces	6.47 (0.48)	
MP17 V	10.17 (S)	0.98 (0.01)	-11.79 (0.01)	1.82 (0.01)	2.0583	46.65	-0.54 (0.16)	-0.39 (0.40)	8.67 (0.08)	8.82 (0.22)	75.25 (3.43)	17.30 (1.78)	—	5.47 (1.83)	1.91	traces	9.78 (0.58)	
MP17 VII ¹³	3.80 (S)	1.00	-11.78	1.83 (0.20)	2.0547	46.35	-0.14 (0.15)	—	9.06 (0.70)	—	89.96 (0.57)	—	—	—	10.14	traces	—	
MP17 VIII	1.10 (po5-por7)	1.00	-11.78	1.83 (0.20)	2.0652	47.25	-1.41 (0.13)	—	7.80 (0.13)	—	88.70 (1.37)	10.60 (1.11)	—	26.46 (0.83)	0.96	—	5.87 (0.32)	
PM17 VI	3.60 (S)	0.95 (0.05)	-11.83	1.79 (0.05)	—	—	—	1.06 (1.16)	—	10.27	43.62 (1.14)	28.96 (0.75)	—	—	—	—	—	
P = 2.054 kbar, T = 930°C, $HH_2^{13} = 2.2$ bar, 169 h																		
MP2 I	1.00 (S)	0.86	-9.99	1.38 (0.08)	2.0488	45.65	1.46	—	9.22	—	98.34 (0.56)	—	—	—	1.66	—	—	
MP2 II	1.00 (po)	0.77	-10.09	1.29 (0.15)	2.0388	45.13	1.97	—	9.72	—	98.23 (0.56)	—	—	—	1.77	—	1.39 (0.65)	
MP2 III	1.00 (am)	0.81	-10.04	1.34 (0.15)	—	—	—	—	—	—	96.07 (0.56)	—	—	—	1.50	—	2.43	
P = 2.257 kbar, T = 933°C, $HH_2^{13} = 0.41$ bar, 175 h																		
MP3 I	1.42 (S)	0.56	-8.90	2.42 (0.29)	—	—	—	—	—	—	98.10 (0.56)	—	—	—	traces	1.7	5.61 (0.06)	
MP3 II	1.00 (po)	0.49	-9.02	2.30 (0.29)	—	—	—	—	—	—	99.13 (0.56)	—	—	—	—	0.87	4.56 (0.07)	
MP3 III	1.00 (am)	0.99	-8.41	2.91 (0.29)	—	—	—	—	—	—	97.69 (0.56)	—	—	—	—	2.31	3.39 (0.08)	
P = 1.986 kbar, T = 932°C, $HH_2^{12} = 46.7$ bar, 175 h																		
MP4 I	1.03 (S)	0.62	-12.93	-1.58 (0.14)	2.0636	47.10	-0.21	—	7.51	—	98.33 (0.56)	—	—	—	1.67	—	1.95 (0.77)	
MP4 II	1.00 (po)	0.43	-13.24	-1.89 (0.14)	2.0780	48.43	-2.26	—	5.46	—	97.12 (0.56)	—	—	—	2.88	—	2.33 (0.69)	
MP4 III	1.00 (am)	0.78	-12.73	-1.38 (0.14)	2.0684	47.53	0.81	—	6.92	—	98.64 (0.56)	—	—	—	1.36	—	1.43 (0.58)	

Charges	wt % S ¹	$a_{H_2O}^2$	$\log O_2^3$	ΔNNO^4	$d(102)^5$	at % Fe ⁶	$\log S_2$ (po) ⁷	$\log S_2$ (V) ⁸	ΔFFS (po) ⁹	ΔFFS (V) ¹⁰	GI	PI	Px	Oz	Po	An	V
<i>P = 2.116 kbar, T = 936°C, $HH_2^{12} = 0.9$ bar, 164 h</i>																	
MP9 I	1.07 (S)	0.96	-9.12	2.16 (0.13)	2.0606	46.85	0.16	-	7.86	-	99.22 (0.56)	-	-	-	0.76	traces	1.16 (0.65)
MP9 III	1.00 (po)	1.00	-9.01	2.27 (0.16)	2.0562	46.47	0.62	-	8.31	-	98.27 (0.56)	-	-	-	1.72	traces	-
MP9 V	1.00 (an)	1.00	-8.99	2.29 (0.16)	2.0567	47.10	-0.17	-	7.53	-	97.97 (0.56)	-	-	-	-	2.03	-
<i>P = 2.013 kbar, T = 933°C, $HH_2^{11} = 47.3$ bar, 168 h</i>																	
MP10 I	2 (S)	0.81	-12.71 (0.12)	-1.38 (0.09)	2.0685	47.19	-0.31 (0.14)	-0.34	7.42 (0.07)	7.38 (0.22)	97.84	-	-	-	2.16	-	1.93 (0.45)
MP10 II	3 (S)	0.75	-12.78 (0.18)	-1.45 (0.13)	2.0585	46.67	0.35 (0.15)	-1.27 (1.96)	8.08 (0.07)	7.60 (0.22)	98.02	0.60 (0.74)	-	-	1.38	-	3.15 (0.61)
MP10 III	4 (S)	0.72	-12.82 (0.16)	-1.48 (0.11)	2.0572	46.55	0.49 (0.14)	-0.23 (0.61)	8.21 (0.07)	7.68 (0.16)	96.8	1.89 (0.80)	-	-	1.3	-	4.66 (0.53)
MP10 IV	10 (S)	0.5	-13.18 (0.37)	-1.85 (0.27)	2.0540	46.30	0.79 (0.14)	0.27 (0.48)	8.51 (0.07)	8.11 (0.14)	92.83	5.36 (1.04)	-	-	1.71	-	10.73 (0.96)
MP10 V	1 (po05)	1	-12.53 (0.14)	-1.18 (0.14)	2.0761	48.25	-1.93 (0.18)	-	5.79 (0.08)	-	96.16	-	-	-	3.85	-	-
MP10 VI	1 (po07)	1	-12.53 (0.14)	-1.18 (0.14)	2.0760	48.23	-1.91 (0.15)	-	5.82 (0.09)	-	96.55	-	-	-	3.45	-	-
<i>P = 2.202 kbar, T = 935°C, $HH_2^{11} = 20.4$ bar, 164 h</i>																	
MP11 I	2.12 (S)	0.74	-11.96 (0.20)	-0.65 (0.14)	2.0553	46.40	0.70 (0.13)	-0.45 (2.18)	8.43 (0.07)	8.52 (0.22)	98.18	-	-	-	1.82	-	2.20 (0.79)
MP11 II	4.83 (S)	0.78	-11.92 (0.18)	-0.61 (0.13)	2.0512	46.13	1.01 (0.14)	1.02	8.73 (0.07)	8.42 (0.24)	98.96	-	-	-	1.04	-	2.95 (0.64)
MP11 III	4.83 (S)	0.63	-12.11 (0.19)	-0.80 (0.14)	2.0570	46.54	1.50 (0.14)	0.94 (0.42)	9.29 (0.07)	8.76 (0.13)	93.15	2.50 (1.36)	-	-	1.65	-	5.68 (0.65)
MP11 IV	10.12 (S)	0.51	-12.31 (0.37)	-1.00 (0.26)	n.d.	-	-	1.09 (0.50)	-	8.95 (0.15)	89.7	4.40 (1.33)	-	-	1.65	-	11.39 (0.95)
<i>P = 2.202 kbar, T = 935°C, $HH_2^{11} = 20.4$ bar, 164 h</i>																	
MP11 V	1.20 (Po6)	0.97	-11.71 (0.05)	-0.41 (0.05)	2.0748	48.12	-1.68 (0.15)	-	6.05 (0.07)	-	96.9	-	-	-	3.1	-	-
MP11 VI	1.10 (Po7)	0.97	-11.71 (0.05)	-0.41 (0.05)	2.0750	48.15	-1.72 (0.16)	-	6.00 (0.08)	-	97.09	-	-	-	2.91	-	-
<i>P = 2.003 kbar, T = 934°C, $HH_2^{11} = 15$ bar, 213 h</i>																	
MP6 II	0.62 (S)	0.83	-11.66 (0.20)	-0.34 (0.14)	2.0739	48.04	-1.57 (0.21)	-	6.15 (0.10)	-	99.13	-	-	-	0.87	-	-
MP6 III	1.02 (S)	0.74 (0.11)	-11.79 (0.18)	-0.46 (0.13)	2.0709	47.76	-1.14 (0.15)	0.57 (0.92)	6.88 (0.08)	6.66 (0.20)	98.34	-	-	-	1.66	-	1.43 (0.51)
MP6 IV	1.33 (S)	0.76 (0.15)	-11.76 (0.24)	-0.43 (0.17)	2.0631	47.06	-0.14 (0.14)	1.25	7.57 (0.07)	6.62 (0.27)	98.31	-	-	-	1.69	-	1.49 (0.68)
MP6 V	1.03 (Po7)	0.93	-11.56 (0.08)	-0.24 (0.08)	2.0740	48.05	-1.59 (0.15)	-	6.13 (0.07)	-	97.31	-	-	-	2.69	-	-
MP6 VI	1.04 (Po6)	1	-11.50 (0.08)	-0.17 (0.08)	2.0740	48.05	-1.59 (0.15)	-	6.12 (0.08)	-	97.52	-	-	-	2.48	-	-
<i>P = 2.052 kbar, T = 934°C, $HH_2^{11} = 6$ bar, 147 h</i>																	
MP12 I	2.31 (S)	0.77 (0.13)	-10.92 (0.21)	0.40 (0.15)	2.0541	46.30	0.80 (0.14)	1.97	6.52 (0.07)	9.35 (0.26)	97.85	-	-	-	2.15	-	1.67 (0.65)
MP12 II	3.90 (S)	0.79 (0.14)	-10.90 (0.21)	0.41 (0.15)	2.0552	46.39	0.70 (0.14)	1.96	8.42 (0.07)	9.33 (0.27)	97.89	-	-	-	2.11	-	4.04 (0.85)
MP12 III	5.86 (S)	0.65 (0.17)	-11.09 (0.33)	0.22 (0.23)	2.0569	46.53	0.53 (0.17)	2.23	8.25 (0.07)	9.65 (0.22)	91.85 (2.66)	3.60 (1.30)	-	-	1.65	-	6.23 (0.78)
MP12 IV	10.31 (S)	0.57 (0.18)	-11.22 (0.40)	0.10 (0.29)	2.0588	46.70	0.33 (0.20)	1.71	6.06 (0.10)	9.78 (0.19)	86.20 (2.75)	9.10 (1.34)	-	-	1.65	-	11.66 (0.88)
<i>P = 1.980 kbar, T = 938°C, $HH_2^{11} = 4$ bar, 162 h</i>																	
MP7 III	1.00 (S)	0.78 (0.13)	-10.50 (0.21)	0.75 (0.15)	2.0443	45.53	1.62 (0.12)	2.16	9.29 (0.06)	9.46 (0.27)	98.61	-	-	-	1.39	-	1.40 (0.63)
MP7 IV	1.25 (S)	0.72 (0.08)	-10.56 (0.14)	0.69 (0.10)	2.0509	46.04	1.10 (0.13)	1.74 (0.60)	8.76 (0.06)	9.58 (0.16)	98.95	-	-	-	1.05	-	2.25 (0.43)
<i>P = 2.001 kbar, T = 930°C, $HH_2^{11} = 0.9$ bar, 168 h</i>																	
MP13 I	2.00 (S)	0.36 (0.03)	-9.92 (0.06)	1.47 (0.05)	2.0530	46.21	0.86 (0.14)	2.75 (0.18)	8.62 (0.07)	10.52 (0.05)	96.21 (0.89)	-	-	-	1.63	0.37	3.80 (0.18)
MP13 II	3.87 (S)	0.38 (0.05)	-9.97 (0.15)	1.42 (0.11)	2.0577	46.60	0.41 (0.14)	2.79 (0.27)	8.17 (0.07)	10.59 (0.06)	95.74 (0.88)	-	-	-	1.69	0.43	5.80 (0.30)
MP13 III	6.04 (S)	0.29 (0.03)	-10.15 (0.13)	1.23 (0.09)	2.0538	46.28	0.79 (0.16)	3.00 (0.15)	8.55 (0.08)	10.77 (0.05)	97.41 (0.90)	-	-	-	1.79 (0.85)	0.42	8.82 (0.63)
MP13 IV	10.11 (S)	0.22 (0.02)	-10.38 (0.13)	1.00 (0.09)	2.0554	46.41	0.64 (0.22)	3.16 (0.09)	8.39 (0.11)	10.92 (0.03)	98.03	traces	-	-	1.73	0.24	13.37 (0.41)
<i>P = 2.188 kbar, T = 984°C, $HH_2^{12} = 30$ bar, 152 h</i>																	
MP18 I	1.25 (S)	0.55 (0.09)	-11.64 (0.43)	-1.28 (0.43)	2.0737	48.02	-1.04 (0.16)	0.69 (1.12)	6.07 (0.08)	6.30 (0.20)	99.27	-	-	-	0.73	-	2.46 (0.13)
MP18 II	0.60 (S)	0.75 (0.05)	-11.29 (0.44)	-0.83 (0.44)	2.0734	47.99	-1.00 (0.22)	0.41 (0.98)	6.11 (0.11)	7.94 (0.20)	98.68	-	-	-	1.32	-	1.16 (0.20)
MP18 IV	2.04 (S)	0.65 (0.04)	-11.69 (0.52)	-1.34 (0.37)	2.0655	47.27	0.02 (0.14)	-0.10 (1.67)	7.13 (0.07)	7.89 (0.21)	98.42	-	-	-	1.88	-	2.72 (0.18)
MP18 V	4.32 (S)	0.37 (0.02)	-12.11 (0.42)	-1.75 (0.42)	2.0580	46.63	0.81 (0.13)	0.94 (0.78)	7.92 (0.06)	8.34 (0.18)	98.41	-	-	-	1.59	-	6.01 (0.20)
MP18 VI	6.33 (S)	0.23 (0.02)	-12.52 (0.42)	-2.15 (0.30)	2.0533	46.23	1.24 (0.12)	1.09 (0.77)	8.35 (0.06)	7.48 (0.16)	98.27	-	-	-	1.73	-	8.63 (0.15)

Table 4: continued

Charges	wt % S ¹	$\theta_{H_2O}^2$	$\log f_{O_2}^3$	ΔNNO^4	$d(102)^5$	at % Fe ⁶	$\log f_{S_2}(\text{po})^7$	$\log f_{S_2}(\text{IV})^8$	$\Delta FFS(\text{po})^9$	$\Delta FFS(\text{V})^{10}$	GI	Pl	Px	Qz	Po	An	V
P = 2.188 kbar, T = 994°C, $HH_2^{12} = 30 \text{ bar}$, 162 h																	
MP18 VII	10.84 (S)	0.33 (0.03)	-12.21 (0.42)	-1.85 (0.30)	2.0509	46.04	1.44 (0.13)	0.99 (0.77)	8.55 (0.06)	8.38 (0.18)	98.17	—	—	—	1.83	—	12.73 (0.31)
MP18 X	3.62 (S)	0.56 (0.11)	-11.77 (0.40)	-1.41 (0.48)	2.0661	47.33	-0.05 (0.14)	0.63 (0.88)	7.06 (0.07)	8.09 (0.19)	98.45	—	—	—	1.55	—	6.52 (0.56)
MP18 XI	3.77 (S)	0.72 (0.15)	-11.55 (0.40)	-1.19 (0.48)	2.0727	47.93	-0.90 (0.15)	0.52 (0.52)	6.21 (0.07)	7.79 (0.28)	98.66	—	—	—	1.34	—	9.75 (0.82)
MP18 XIII ¹³	3.30 (S)	0.83 (0.07)	-11.11 (0.07)	-0.75 (0.07)	2.1718	47.84	-0.78 (0.14)	-1.36 (1.39)	6.33 (0.07)	8.23 (0.70)	97.24 (0.56)	—	—	—	2.76	—	2.54 (0.52)
MP18 SN	1.32 (po5-po7)	0.83	-11.1	-0.74	2.0741	48.06	-1.11 (0.11)	—	6.00 (0.12)	—	98.47	—	—	—	3.53*	—	—
PN18 IX	3.26 (S)	0.24 (0.12)	-12.62 (0.64)	-2.26 (0.77)	2.0577	46.60	1.00 (0.13)	1.18 (0.62)	8.11 (0.06)	8.48 (0.27)	98.13 (0.56)	—	—	—	1.87	—	5.74 (0.63)
P = 2.068 kbar, T = 996°C, $HH_2^{12} = 2 \text{ bar}$, 162 h																	
MP19 VII	0.54 (S)	0.49 (0.11)	-9.26 (0.22)	1.06 (0.19)	—	—	—	2.22 (0.80)	—	9.59 (0.19)	100	—	—	—	—	—	2.08 (0.47)
MP19 II	1.42 (S)	0.75 (0.10)	-8.86 (0.16)	1.46 (0.12)	—	—	—	0.97 (0.97)	—	8.67 (0.25)	100	—	—	—	—	—	2.33 (0.46)
MP19 III	2.34 (S)	0.61 (0.16)	-9.08 (0.33)	1.25 (0.24)	—	—	—	1.31 (1.31)	—	9.34 (0.27)	98.34	—	—	0.66	—	—	3.57 (0.74)
MP19 IV	3.09 (S)	0.39 (0.17)	-9.52 (0.57)	0.81 (0.41)	—	—	—	1.55 (1.55)	—	9.86 (0.23)	99.39	—	—	0.61	—	—	5.33 (0.82)
MP19 VI	5.03 (S)	0.01	-12.61	-2.29	—	—	—	3.24	—	10.31	36.70	39	—	23	1.40	—	4.47 (0.10)
MP19 IX	3.39 (S)	0.88 (0.13)	-8.74 (0.18)	1.59 (0.12)	—	—	—	0.75 (0.57)	—	8.20 (0.31)	98.64	—	—	0.36	—	—	5.97 (0.60)
MP19 X	3.30 (S)	0.87 (0.13)	-8.74 (0.18)	1.59 (0.19)	—	—	—	0.78 (0.59)	—	8.25 (0.31)	100	—	—	traces	—	—	6.12 (1.12)
MP19 XI	3.29 (S)	0.88 (0.09)	-8.74 (0.12)	1.59 (0.09)	—	—	—	0.67 (0.50)	—	8.03 (0.29)	98.80	—	—	0.20	—	—	8.90 (0.47)
MP19 XIII ¹³	3.50 (S)	0.83 (0.17)	-8.80 (0.19)	1.53 (0.18)	—	—	—	-0.81 (2.85)	—	6.25 (1.32)	98.10 (0.56)	—	—	0.90	—	—	3.79 (0.80)
MP19 XIII	1.27 (po5-po7)	0.47	-9.27	1.05	—	—	—	2.48 (0.30)	—	9.55 (0.30)	98.29	—	—	1.71*	—	—	2.31 (0.78)
PN19 VIII	3.15 (S)	0.41 (0.14)	-9.45 (0.46)	0.88 (0.32)	—	—	—	2.18 (1.21)	—	9.81 (0.21)	100	—	—	—	—	—	5.60 (0.66)
P = 1.992 kbar, T = 987°C, $HH_2^{12} = 0.5 \text{ bar}$, 157 h																	
MP14 I	1.99 (S)	0.47 (0.04)	-8.24 (0.12)	2.23 (0.09)	—	—	—	0.61 (0.49)	—	7.89 (0.14)	98.15 (0.92)	—	—	—	trace	0.85	3.99 (0.22)
MP14 II	3.86 (S)	0.43 (0.02)	-8.29 (0.07)	2.18 (0.06)	—	—	—	0.80 (0.29)	—	8.00 (0.09)	99.08	—	—	—	trace	0.92	6.00 (0.21)
MP14 III	6.04 (S)	0.31 (0.06)	-8.62 (0.19)	1.85 (0.14)	2.0529	46.20	1.21 (0.17)	1.42 (0.78)	8.37 (0.08)	8.85 (0.18)	97.90 (0.88)	—	—	1.08	—	1.02	8.31 (0.53)
MP14 IV	10.11 (S)	0.23 (0.03)	-8.86 (0.19)	2.00 (0.31)	2.0552	46.39	1.01 (0.14)	2.09 (0.37)	8.17 (0.07)	9.32 (0.12)	97.62	traces	—	1.38	1	12.09 (0.46)	
MP14 VI	1.50 (po5-po7)	0.40 (0.05)	-8.38 (0.11)	2.03 (0.06)	2.0556	46.44	0.95 (0.12)	1.08 (0.30)	8.11 (0.12)	8.49 (0.30)	98.00 (1.25)	—	—	1.20 (0.22)	traces	3.45 (0.29)	
MP14 VII ¹³	3.50 (S)	0.57 (0.04)	-8.12 (0.02)	2.36 (0.03)	—	—	—	0.22 (0.19)	—	7.37 (0.09)	95.70 (0.59)	—	—	3.10	0.90	—	5.08 (0.22)
PN14 VIII	3.78 (S)	0.43 (0.03)	-8.32 (0.08)	2.14 (0.06)	—	—	—	0.87 (0.32)	—	8.09 (0.10)	98.08 (0.91)	—	—	traces	traces	—	6.00 (0.21)

¹Amount of sulphur added to the charge either as elemental (S) or as pyrrhotite (Po5—Po7; see Table 3).

²Water activity calculated from Burnham (1979).

³ f_{O_2} computed from equation (2).

⁴ f_{O_2} referenced to the NNO buffer (Holloway *et al.*, 1992).

⁵Distance between inter-reticular planes 102 of hexagonal pyrrhotite determined by XRD.

⁶Iron-content of pyrrhotite (atomic) from Yund & Hall (1976) using $d(102)$.

⁷ $\log f_{S_2}$ given by the pyrrhotite composition, using the calibration of Froese & Gunter (1976). Numbers in italics are for $f_{O_2} > \text{NNO}$ (quenched Fe—S—O liquid).

⁸ $\log f_{S_2}$ calculated using and MRK-EOS (see text). Numbers in italics are for charges whose total sulphur content computed from MRK-EOS calculation is significantly different from the sulphur content calculated from mass balance.

⁹ $\log f_{S_2}$ calculated from pyrrhotite composition referenced to the iron—troilite (FFS) buffer (see text).

¹⁰ $\log f_{S_2}$ given by the MRK-EOS (see text). Numbers in italics are for charges whose total sulphur content computed from MRK-EOS calculation is significantly different from the sulphur content calculated from mass balance.

¹¹ f_{H_2} from membrane reading.

¹² f_{H_2} from NIPd sensor measurement.

¹³Magnetite-added charges.

Modal composition determined by mass balance. Phase abbreviations: Gl, glass; Pl, plagioclase; Px, pyroxene; Qz, quartz; Po, pyrrhotite; An, anhydrite; V, fluid phase.

1 σ for Gl, Po, An, which are not given in parentheses, are 0.56, 0.16 and 0.13, respectively.




Fig. 2. Attainment of equilibrium in fS_2 for the experiment MP10. $d(102)$ is the inter-reticular distance of the 102 plane of hexagonal pyrrhotite determined by XRD. Error bars are smaller than symbols. The starting pyrrhotite compositions (Po5 and Po7) are shown as black dots. Open dots are the final pyrrhotite compositions. The arrows indicate the changes in pyrrhotite compositions. Each pyrrhotite was loaded in separate capsules run at the same P - T - fH_2 conditions and the compositional similarity [similar $d(102)$] observed after the experiment indicates that both pyrrhotites record the same fS_2 .

48 h. The compositions of the two synthetic pyrrhotites are given in Table 3. In the first capsule, the synthetic starting pyrrhotite was iron-rich [$NFeS = 0.975$; $NFeS$ is the mole fraction of FeS in the system $FeS-S_2$ (Toulmin & Barton, 1964)] and, in the second one, the pyrrhotite was sulphur-rich ($NFeS = 0.92$). During the experiment, pyrrhotites in these two capsules re-equilibrated to the prevailing conditions, yielding final pyrrhotite compositions identical within error and suggesting attainment of equilibrium fS_2 , as illustrated in Fig. 2 for the experiments MP10. In all pyrrhotite-bearing charges, the two sensor capsules yielded identical fS_2 (except when the prevailing fO_2 is outside the stability field of pyrrhotite, such as MP19 charges) and we conclude that fS_2 is at equilibrium for all experiments presented here. Attainment of equilibrium conditions is corroborated by the compositional homogeneity of glasses, both for major (e.g. SiO_2) and minor element (S) concentrations.

ANALYTICAL TECHNIQUES

Run products were examined by optical and scanning electron microscopy (SEM). We used X-ray diffractometry to confirm the phase assemblage determined by optical microscopy and SEM. Besides the characterization of the run products, X-ray diffractometry of powdered run products was also used to measure the composition of pyrrhotites (Arnold & Reicher, 1962; Toulmin & Barton, 1964; Boorman, 1967; Yund & Hall, 1969). Our measurements were made on an Inel CPS 120 diffractometer at ambient temperature using Co radiation ($\lambda = 1.78897 \text{ \AA}$) with quartz as an external

standard. The measured peak positions are accurate to within 0.01° (Roux & Volfinger, 1996).

Analyses of the major elements of the experimental charges were performed on both Camebax and SX50 electron microprobes. The operating analytical conditions were: accelerating voltage 15 kV, beam current 6 and 12 nA for glass and crystals, respectively, beam diameter $10 \mu\text{m}$ for glass and focused beam ($1-2 \mu\text{m}$) for crystals. A ZAF correction procedure was applied. Correction for electron beam induced alkali-migration and determination of H_2O concentration in glasses by the by-difference method (Devine *et al.*, 1995) were performed by using calibration curves constructed from three hydrous standard glasses, analysed by wet chemistry for their Na and K content (Pichavant, 1987). These standards have the same composition as our starting material (without sulphur) and their water contents (up to 6.5 wt %) are known from Karl Fisher titration. Major element calibration used wollastonite (Ca), hematite (Fe), albite (Na, Si), corundum (Al), olivine (Mg) and orthoclase (K) standards. The detection limit for major elements is about 1000 ppm. For the fH_2 sensor, the analytical conditions for Pt, Pd, Ni were: accelerating voltage 20 kV and a focused beam current of 20 nA. Counting time was 10 s on the peak and 5 s on the background for all major element analyses. Anhydrite was analyzed (using a JEOL JXA-8600 electron microprobe) under analytical conditions of 15 kV, 6 nA and $5 \mu\text{m}$ beam diameter, using anhydrite as a standard.

The concentration of total sulphur in glasses was determined by electron microprobe analysis (Camebax). In silicate melts, sulphur is assumed to be present as sulphide (S^{2-}) or sulfate (S^{6+}) (Richardson & Withers, 1950; Ricke, 1960; Schneider, 1970). The position of the SK_a X-ray has been shown to be a function of the sulphur valence state and of the complexation of sulphur (e.g. Carroll & Rutherford, 1988; Kucha *et al.*, 1989). In our glasses, sulphur is present either as sulphide or sulfate, or as a mixture of the two. In this last case, the SK_a X-ray peak is between the sulphide and sulfate positions. Therefore, at the beginning of each analytical session, we established two calibration curves—one for S as sulphide and one for S as sulfate. These curves were constructed with three synthetic hydrous dacitic standard glasses, containing 750, 1400, and 1900 ppm sulphur, determined by wet chemistry. The difference between the two calibration curves increases with the sulphur content, but remains below detection for S contents below 400 ppm. Consequently, the S-peak position was determined for all glasses with sulphur contents higher than 400 ppm, and the counting was performed on each specific peak. For S analyses, we used three PET crystal spectrometers, using the following analytical procedure: accelerating voltage 15 kV, sample current 50 nA, beam diameter $10 \mu\text{m}$. Counting time was 60 s on each spectrometer, resulting

in a total counting time of 180 s. The background was determined by analysing a glass of the same composition without sulphur, using the above analytical procedure. The detection limit under these analytical conditions is about 80 ppm, as calculated from Ancy *et al.* (1978). Single spot analyses of 1 h duration with these analytical conditions showed no migration of sulphur.

Qualitative analyses of oxygen in sulphides by electron microprobe (SX50) have been performed to check whether the pyrrhotites are the products of back reactions of immiscible Fe–S–O liquids upon quenching. Specifically, under conditions of high fO_2 , pyrrhotite may partially break down to an Fe–O–S liquid. It appeared that, indeed, for $fO_2 > NNO$, most quench sulphides contained detectable amounts of oxygen. Therefore, the pyrrhotite in these charges may be because of the crystallization of Fe–S–O immiscible liquids during the quench, and we did not consider as reliable the fS_2 retrieved from pyrrhotite composition in those charges.

RESULTS

The experimental conditions, phase assemblages and proportions, and glass compositions are summarised in Tables 4 and 5.

Phase assemblages

The phase assemblages are portrayed in Fig. 3. Besides direct evidence of fluid in excess upon opening of the capsules, such as drops of water and hissing, fluid-saturated conditions were also assumed when bubble sizes in quenched glasses exceeded 10 μm . However, virtually all charges contain micrometric to sub-micrometric bubbles, which most probably represent remnants of air trapped in the capsules. No magnetite crystallized, except at high fO_2 ($> NNO+1.3$), where a trace amount of this oxide was detected (charges MP16 VII, MP17 VII and MP14 VII, Table 4). Previous experimental work, performed on various rhyolitic compositions, has shown that the quench rates used in the present study are high enough to prevent quench crystallization of oxides (Gaillard *et al.*, 2001). The comparison between magnetite-doped and undoped charges shows that below $NNO+1$, addition of magnetite to a sulphur-bearing charge merely results in extensive sulphide crystallization (Table 4). At all temperatures with $fO_2 < NNO$, the sulphide is pyrrhotite, whereas above NNO , the sulphide is an Fe–S–O immiscible liquid \pm pyrrhotite, as indicated by detection of O by EMPA (Fig. 3). For charges in which the amount of added sulphur exceeds the sulphur solubility of the melt, an S–O–H fluid phase forms, and water becomes partitioned between melt and fluid. Further addition of sulphur to the system decreases the activity of water through a simple dilution effect (i.e. the H_2O/S ratio in the fluid

decreases) and because of interaction with sulphur species (i.e. production of H_2S). A significant decrease in aH_2O may lead to the partial crystallization of other, S-free phases, such as plagioclase, pyroxene or quartz. For instance, at 800°C, plagioclase is stable for aH_2O slightly below 1, pyroxene is stable under reducing conditions for aH_2O slightly below 1, and quartz crystallizes for $aH_2O \leq 0.85$. Similarly, at 930°C, plagioclase is stable for $aH_2O \leq 0.75$ and quartz is stable for $aH_2O \leq 0.65$ above $NNO-1$. At 1000°C, apart from one charge which crystallized plagioclase, only S-bearing phases were present in charges loaded with elemental sulphur. Charges doped with magnetite resulted in extensive sulphide crystallization (up to 10 wt %), depending on the amount of sulphur added (Table 4), except at high fO_2 at 800 and 930°C, where iron oxides crystallized (MP16 VII, MP17 VII and MP14 VI charges, Table 4). Anhydrite is stable under oxidizing conditions only, confirming other experimental studies (Carroll & Rutherford, 1987; Luhr, 1990; Scaillet & Evans, 1999). At 800 and 930°C, anhydrite is stable at $fO_2 > NNO+1$. At 1000°C, anhydrite appears at $fO_2 > NNO+1.8$, which indicates that the fO_2 needed to crystallize anhydrite in rhyolitic melts increases with T (Fig. 3). Anhydrite stability in rhyolitic magmas appears to be in the same fO_2 range as that for trachyandesite magmas at $T < 950^\circ\text{C}$ (Carroll & Rutherford, 1987; Luhr, 1990).

Fluid phase composition

The composition of the coexisting fluid-phase was calculated using the MRK equation of state, as explained above. The mole fraction of H_2O in the fluid (X_{H_2Ofl}) ranges between near 1 and 0.1, yet, for the vast majority of charges, it is within the range 0.75–0.30. The few charges having X_{H_2Ofl} significantly lower than 0.2 are those for which the by-difference method yields H_2O melt content close to zero (e.g. charge MP15V, Table 4). At all temperatures, as long as the fO_2 is below $NNO+1$, H_2S is the dominant sulphur-bearing species, and its mole fraction in the fluid (X_{H_2Sfl}) ranges from 0.1 up to 0.85. Thus, at 2 kbar and at an fO_2 below $NNO+1$, the fluid phase can be considered as a binary mixture of H_2O and H_2S , other fluid species (H_2 , SO_2 , S_2) amounting to no more than 0.01 mole fraction. SO_2 becomes abundant at higher fO_2 and its mole fraction (X_{SO_2fl}) may even exceed that of H_2S at 930 and 1000°C. At 930°C, an fO_2 of $NNO+1.4$ is required to have $X_{SO_2fl} > X_{H_2Sfl}$. At 1000°C, the same happens at slightly lower fO_2 , that is around $NNO+1$. At 800°C, however, H_2S dominates over SO_2 , even with an fO_2 as high as $NNO+1.8$. At $fO_2 > NNO+1$, and in charges loaded with massive amounts of sulphur (5–10 wt %, Table 4), S_2 makes up a significant proportion of sulphur-bearing species in the fluid—up to 0.3 of X_{S_2fl} .

Table 5: Compositions of experimental glasses (normalized to 100 wt % anhydrous)

Charges	ΔNNO^1	ΔFFS^2	SiO_2	Al_2O_3	FeO^3	MgO	CaO	Na_2O	K_2O	H_2O^4	S (ppm) ⁵	log S
P = 2.152 kbar, T = 765°C												
MP15 I	-1.12 (0.10)	7.45 (0.21)	78.35 (0.91)	13.25 (0.37)	0.28 (0.10)	0.29 (0.10)	1.62 (0.13)	4.17 (0.21)	2.02 (0.14)	4.69 (0.43)	≤D	—
MP15 II	-1.16 (0.29)	9.43 (0.19)	78.52 (0.99)	13.08 (0.41)	0.14 (0.10)	0.27 (0.10)	1.01 (0.11)	4.44 (0.24)	2.52 (0.18)	4.70 (1.00)	440 (11)	2.64
MP15 III	-1.83 (0.22)	9.95 (0.19)	78.63 (0.90)	12.54 (0.36)	0.14 (0.10)	0.26 (0.10)	0.94 (0.10)	4.18 (0.21)	3.30 (0.18)	2.65 (0.44)	612 (19)	2.79
MP15 IV	-1.93 (0.39)	10.25 (0.18)	78.59 (0.65)	12.57 (0.26)	0.18 (0.10)	0.18 (0.10)	0.67 (0.10)	3.92 (0.15)	3.88 (0.14)	2.56 (0.70)	753 (40)	2.88
MP15 VI ⁶	-0.93	6.19 (0.13)	78.03 (0.65)	13.02 (0.26)	1.51 (0.10)	0.17 (0.10)	1.65 (0.10)	3.71 (0.14)	1.89 (0.10)	5.54 (0.75)	≤D	—
MP15 VII ⁷	-0.93	6.36 (0.22)	76.57 (1.27)	12.98 (0.52)	1.59 (0.18)	0.28 (0.10)	2.08 (0.21)	4.44 (0.32)	2.07 (0.23)	6.78 (0.65)	≤D	—
MP15 X	-1.03 (0.08)	7.42 (0.21)	77.53 (0.65)	13.71 (0.27)	0.17 (0.10)	0.32 (0.10)	1.98 (0.10)	4.34 (0.15)	1.94 (0.10)	5.93 (1.06)	126 (34)	2.10
MP15 XI	-0.98 (0.09)	7.13 (0.21)	77.46 (1.01)	13.75 (0.42)	0.63 (0.11)	0.35 (0.10)	2.00 (0.16)	3.87 (0.23)	1.93 (0.16)	6.37 (1.10)	≤D	—
MP15 SN	-1.18 (0.20)	7.24 (0.21)	77.96 (0.58)	13.51 (0.24)	0.23 (0.10)	0.34 (0.10)	1.77 (0.10)	4.27 (0.14)	1.92 (0.10)	4.60 (0.60)	96 (61)	1.98
PIN15VIII	-1.16 (0.07)	9.86 (0.19)	77.84 (1.17)	12.79 (0.47)	0.09 (0.10)	0.29 (0.10)	0.57 (0.10)	4.03 (0.28)	4.38 (0.29)	4.70 (0.30)	787 (21)	2.90
P = 1.973 kbar, T = 798°C												
MP16 I	1.30 (0.12)	10.76 (0.19)	78.78 (0.83)	12.31 (0.33)	0.16 (0.10)	0.33 (0.10)	1.45 (0.11)	2.87 (0.17)	4.10 (0.19)	5.01 (0.75)	108 (64)	2.03
MP16 II	1.18 (0.13)	10.99 (0.26)	78.83 (1.06)	12.20 (0.42)	0.08 (0.10)	0.33 (0.10)	1.04 (0.12)	2.79 (0.21)	4.71 (0.26)	4.32 (0.52)	294 (9)	2.47
MP16 III	1.16 (0.19)	11.04 (0.24)	78.10 (0.49)	12.16 (0.19)	0.10 (0.10)	0.36 (0.10)	0.99 (0.10)	3.02 (0.10)	5.27 (0.13)	4.31 (0.52)	240 (72)	2.38
MP16 IV	0.98 (0.13)	11.35 (0.11)	77.28 (0.74)	12.50 (0.30)	0.10 (0.10)	0.34 (0.10)	0.65 (0.10)	3.02 (0.15)	6.16 (0.21)	3.71 (0.41)	356 (22)	2.55
MP16 V	1.07 (0.16)	11.35 (0.08)	78.09 (0.57)	12.28 (0.22)	0.10 (0.10)	0.37 (0.10)	0.85 (0.10)	2.98 (0.11)	5.39 (0.15)	3.97 (0.54)	274 (16)	2.44
MP16 VII ⁷	1.29	10.38 (0.46)	74.36 (0.49)	12.67 (0.20)	0.26 (0.10)	0.28 (0.10)	1.88 (0.10)	6.93 (0.15)	3.60 (0.11)	4.84 (0.41)	≤D	—
MP16VIII	1.41 (0.09)	10.06 (0.30)	78.04 (0.50)	12.47 (0.20)	0.58 (0.10)	0.29 (0.10)	1.67 (0.10)	2.96 (0.10)	3.97 (0.11)	5.71 (0.52)	≤D	—
MP16 IX	1.41 (0.02)	10.08 (0.27)	77.27 (0.39)	13.62 (0.16)	0.33 (0.10)	0.33 (0.10)	2.01 (0.10)	3.12 (0.10)	3.76 (0.10)	5.58 (0.34)	≤D	—
PIN16 VI	1.08 (0.13)	11.20 (0.17)	77.01 (0.52)	12.53 (0.21)	0.10 (0.10)	0.23 (0.10)	0.73 (0.10)	2.51 (0.10)	6.89 (0.15)	4.04 (0.47)	373 (14)	2.57
P = 1.976 kbar, T = 811°C												
MP17 I	1.83 (0.20)	—	80.01 (0.73)	12.06 (0.30)	0.10 (0.10)	0.37 (0.10)	1.41 (0.12)	3.89 (0.10)	2.15 (0.12)	5.88 (0.39)	125 (16)	2.1
MP17 II	1.83 (0.20)	—	79.93 (0.83)	12.13 (0.14)	0.08 (0.10)	0.40 (0.10)	1.21 (0.11)	3.93 (0.19)	2.32 (0.10)	5.86 (0.39)	263 (11)	2.42
MP17 III	1.83 (0.20)	—	79.90 (0.39)	12.17 (0.12)	0.11 (0.12)	0.40 (0.10)	1.12 (0.10)	3.98 (0.10)	2.32 (0.10)	5.49 (0.20)	304 (19)	2.48
MP17 IV	1.80 (0.09)	—	79.91 (0.70)	12.23 (0.26)	0.12 (0.10)	0.40 (0.10)	1.06 (0.10)	3.94 (0.15)	2.33 (0.18)	5.46 (0.43)	279 (18)	2.45
MP17 V	1.82 (0.01)	—	79.96 (0.70)	12.21 (0.22)	0.06 (0.10)	0.42 (0.10)	1.04 (0.16)	3.87 (0.21)	2.44 (0.17)	5.62 (0.38)	307 (11)	2.49
MP17 VII ⁷	1.83 (0.20)	—	77.84 (0.81)	13.42 (0.41)	0.40 (0.43)	0.31 (0.10)	1.99 (0.15)	4.19 (0.18)	1.95 (0.10)	6.21 (0.47)	83 (24)	1.92
MP17 VIII ⁶	1.83 (0.20)	—	78.62 (1.33)	12.48 (0.24)	1.39 (0.27)	0.21 (0.10)	1.51 (0.10)	3.67 (0.41)	2.12 (0.23)	6.74 (1.25)	≤D	—
PIN 17 VI	1.79 (0.05)	—	79.05 (0.70)	13.04 (0.20)	0.11 (0.14)	0.22 (0.10)	1.05 (0.10)	3.30 (0.13)	3.22 (0.11)	5.43 (0.56)	281 (11)	2.45
P = 2.054 kbar, T = 930°C												
MP2 I	1.39 (0.08)	9.22	76.92 (0.10)	13.62 (0.10)	0.48 (0.10)	0.35 (0.10)	2.07 (0.10)	4.57 (0.10)	1.99 (0.10)	4.97 (0.37)	158 (12)	2.20
MP2 II	1.29 (0.15)	9.72	75.83 (0.23)	13.34 (0.10)	2.41 (0.10)	0.33 (0.10)	1.95 (0.10)	4.22 (0.10)	1.92 (0.10)	4.44 (0.59)	112 (13)	2.05
MP2 III	1.34 (0.15)	—	77.25 (0.33)	13.55 (0.14)	0.45 (0.10)	0.35 (0.10)	2.01 (0.10)	4.43 (0.10)	1.94 (0.10)	4.71 (0.79)	285 (15)	2.46

Table 5: continued

Charges	ΔNNO^1	ΔFFS^2	SiO_2	Al_2O_3	FeO^3	MgO	CaO	Na_2O	K_2O	H_2O^4	S (ppm)^5	$\log S$
P = 2.257 kbar, T = 933°C												
MP3 I	2.42 (0.29)	—	77.30 (0.23)	13.73 (0.10)	1.85 (0.10)	0.32 (0.10)	0.39 (0.10)	4.40 (0.10)	2.01 (0.10)	3.04 (0.47)	989 (19)	2.99
MP3 II	2.30 (0.29)	—	76.90 (0.32)	13.50 (0.13)	1.81 (0.10)	0.33 (0.10)	1.18 (0.10)	4.37 (0.10)	1.90 (0.10)	2.83 (0.70)	307 (14)	2.49
MP3 III	2.91 (0.29)	—	76.22 (0.30)	13.40 (0.13)	1.93 (0.10)	0.33 (0.10)	2.10 (0.10)	4.05 (0.10)	1.96 (0.10)	4.89 (0.60)	186 (13)	2.27
P = 1.986 kbar, T = 932°C												
MP4 I	-1.58 (0.14)	7.51	77.19 (0.10)	13.66 (0.23)	0.19 (0.10)	0.32 (0.10)	2.04 (0.10)	4.60 (0.13)	1.99 (0.10)	3.72 (0.72)	313 (13)	2.50
MP4 II	-1.89 (0.14)	5.46	76.31 (0.26)	13.59 (0.11)	1.43 (0.10)	0.34 (0.10)	2.02 (0.10)	4.38 (0.10)	1.93 (0.10)	2.81 (0.65)	$\leq D$	—
MP4 III	-1.38 (0.14)	6.92	75.55 (0.22)	13.38 (0.10)	0.43 (0.10)	0.20 (0.10)	4.23 (0.10)	4.28 (0.10)	1.93 (0.10)	4.37 (0.53)	148 (11)	2.17
P = 2.116 kbar, T = 936°C												
MP9 I	2.16 (0.13)	7.86	76.85 (0.66)	13.54 (0.31)	1.16 (0.16)	0.32 (0.10)	2.02 (0.10)	3.99 (0.12)	2.12 (0.10)	4.74 (0.58)	77 (61)	1.89
MP9 III	2.27 (0.16)	8.31	76.08 (0.77)	13.53 (0.28)	2.40 (0.27)	0.34 (0.10)	1.91 (0.10)	3.70 (0.59)	2.03 (0.10)	5.13 (0.56)	$\leq D$	—
MP9 V	2.29 (0.16)	7.53	75.81 (0.63)	13.24 (0.34)	1.87 (0.17)	0.35 (0.10)	2.64 (0.10)	3.98 (0.17)	2.11 (0.10)	5.76 (0.49)	154 (6)	2.19
P = 2.013 kbar, T = 933°C												
MP10 I	-1.38 (0.09)	7.42 (0.07)	77.34 (0.67)	13.52 (0.22)	0.31 (0.17)	0.30 (0.10)	2.04 (0.10)	4.38 (0.12)	2.10 (0.10)	4.51 (0.36)	189 (36)	2.28
MP10 II	-1.45 (0.13)	8.08 (0.07)	77.42 (0.61)	13.60 (0.25)	0.13 (0.25)	0.32 (0.10)	1.99 (0.10)	1.39 (0.14)	2.15 (0.10)	4.25 (0.48)	544 (20)	2.74
MP10 III	-1.48 (0.11)	8.21 (0.07)	77.71 (0.38)	13.35 (0.16)	0.20 (0.16)	0.32 (0.10)	1.95 (0.10)	4.38 (0.10)	2.08 (0.10)	4.09 (0.40)	691 (76)	2.84
MP10 IV	-1.85 (0.27)	8.51 (0.07)	77.08 (1.72)	12.94 (0.73)	0.12 (0.16)	0.33 (0.10)	1.70 (0.18)	5.55 (1.39)	2.26 (0.10)	3.14 (0.67)	883 (33)	2.95
MP10 V ⁶	-1.18 (0.14)	5.79 (0.09)	76.76 (0.76)	13.37 (0.23)	1.35 (0.22)	0.31 (0.10)	1.99 (0.10)	4.15 (0.15)	2.07 (0.10)	5.90 (0.55)	$\leq D$	—
MP10 VI ⁶	-1.18 (0.14)	5.82 (0.09)	76.81 (0.77)	13.35 (0.25)	1.22 (0.25)	0.32 (0.10)	1.99 (0.10)	4.26 (0.15)	2.04 (0.10)	5.54 (0.52)	$\leq D$	—
P = 2.202 kbar, T = 935°C												
MP11 I	-0.65 (0.14)	8.43 (0.07)	77.31 (0.67)	13.40 (0.29)	0.34 (0.25)	0.32 (0.10)	2.04 (0.10)	4.48 (0.14)	2.11 (0.10)	4.39 (0.55)	211 (52)	2.32
MP11 II	-0.61 (0.13)	8.73 (0.07)	77.44 (0.70)	13.55 (0.20)	0.05 (0.10)	0.33 (0.10)	2.08 (0.10)	4.48 (0.12)	2.07 (0.11)	4.56 (0.54)	644 (31)	2.81
MP11 III	-0.80 (0.14)	9.29 (0.07)	77.27 (0.66)	13.60 (0.22)	0.11 (0.14)	0.32 (0.10)	2.05 (0.10)	4.52 (0.10)	2.12 (0.10)	3.86 (0.46)	718 (44)	2.86
MP11 IV	-1.00 (0.26)	8.95 (0.15)	77.39 (0.80)	13.54 (0.28)	0.09 (0.10)	0.35 (0.10)	1.92 (0.10)	4.50 (0.22)	2.21 (0.10)	3.31 (0.71)	901 (22)	2.95
MP11 V ⁶	-0.41 (0.05)	6.05 (0.07)	76.38 (0.68)	13.16 (0.24)	1.86 (0.21)	0.33 (0.10)	2.01 (0.14)	4.17 (0.26)	2.08 (0.10)	5.36 (0.37)	$\leq D$	—
MP11 VI ⁶	-0.41 (0.05)	6.00 (0.08)	76.40 (0.50)	13.38 (0.27)	1.54 (0.17)	0.31 (0.10)	2.01 (0.10)	4.24 (0.19)	2.11 (0.11)	5.38 (0.30)	$\leq D$	—
P = 2.003 kbar, T = 934°C												
MP6 II	-0.34 (0.14)	6.15 (0.10)	76.79 (1.10)	13.67 (0.27)	1.06 (0.56)	0.35 (0.10)	1.97 (0.10)	3.99 (0.15)	2.16 (0.10)	4.50 (0.57)	72 (62)	1.86
MP6 III	-0.46 (0.13)	6.58 (0.08)	77.34 (0.51)	13.77 (0.25)	0.44 (0.22)	0.35 (0.10)	2.00 (0.10)	4.02 (0.18)	2.08 (0.10)	4.07 (0.46)	74 (62)	1.87
MP6 IV	-0.43 (0.13)	7.57 (0.07)	77.33 (0.68)	13.67 (0.22)	0.43 (0.27)	0.34 (0.10)	1.98 (0.11)	4.13 (0.15)	2.11 (0.13)	4.21 (0.29)	$\leq D$	—
MP6 VI ⁶	-0.24 (0.08)	6.13 (0.07)	76.49 (0.83)	13.55 (0.16)	1.62 (0.21)	0.31 (0.10)	2.06 (0.11)	3.88 (0.18)	2.08 (0.10)	4.86 (0.79)	$\leq D$	—
MP6 VII ⁶	-0.17 (0.08)	6.12 (0.08)	76.03 (0.59)	13.51 (0.24)	2.11 (0.23)	0.33 (0.10)	1.97 (0.10)	3.94 (0.10)	2.10 (0.10)	5.38 (0.45)	$\leq D$	—

Charges	ΔNNO^1	ΔFFS^2	SiO_2	Al_2O_3	FeO^3	MgO	CaO	Na_2O	K_2O	H_2O^4	S (ppm) ⁵	log S
<i>P = 2.092 kbar, T = 934°C</i>												
MP12 I	0.40 (0.15)	9.35 (0.26)	77.43 (0.65)	13.53 (0.24)	0.09 (0.10)	0.34 (0.10)	2.01 (0.10)	4.45 (0.11)	2.13 (0.10)	4.43 (0.59)	273 (38)	1.58
MP12 II	0.41 (0.15)	9.33 (0.27)	77.33 (0.83)	13.62 (0.20)	0.12 (0.12)	0.33 (0.10)	2.05 (0.12)	4.39 (0.16)	2.15 (0.10)	4.48 (0.62)	334 (77)	2.52
MP12 III	0.22 (0.23)	9.65 (0.22)	77.60 (1.17)	13.48 (0.31)	0.11 (0.10)	0.33 (0.10)	1.94 (0.10)	4.35 (0.23)	2.18 (0.13)	3.83 (0.77)	535 (47)	2.73
MP12 IV	0.10 (0.29)	9.78 (0.19)	78.77 (1.03)	12.75 (0.43)	0.07 (0.12)	0.32 (0.10)	1.47 (0.14)	4.36 (0.23)	2.25 (0.10)	3.47 (0.82)	554 (43)	2.74
<i>P = 1.980 kbar, T = 938°C</i>												
MP7 III	0.75 (0.15)	9.46 (0.27)	76.98 (0.54)	13.79 (0.19)	0.69 (0.14)	0.32 (0.10)	2.04 (0.10)	4.08 (0.11)	2.08 (0.10)	4.27 (0.57)	$\leq D$	—
MP7 IV	0.69 (0.10)	9.58 (0.16)	76.90 (0.74)	13.60 (0.20)	0.94 (0.59)	0.36 (0.10)	2.03 (0.10)	4.06 (0.15)	2.11 (0.10)	4.02 (0.36)	$\leq D$	—
<i>P = 2.001 kbar, T = 930°C</i>												
MP13 I	1.47 (0.05)	10.52 (0.05)	77.31 (0.44)	13.94 (0.19)	0.38 (0.10)	0.31 (0.10)	1.79 (0.10)	4.39 (0.11)	1.88 (0.10)	2.51 (0.11)	371 (51)	2.57
MP13 II	1.42 (0.11)	10.59 (0.09)	77.21 (0.76)	13.97 (0.32)	0.31 (0.10)	0.30 (0.10)	1.72 (0.11)	4.63 (0.19)	1.86 (0.12)	2.45 (0.19)	422 (25)	2.63
MP13 III	1.23 (0.09)	10.77 (0.05)	77.75 (0.54)	13.75 (0.23)	0.21 (0.10)	0.33 (0.10)	1.70 (0.10)	4.38 (0.13)	1.87 (0.10)	2.10 (0.13)	494 (86)	2.69
MP13 IV	1.00 (0.09)	10.92 (0.03)	77.19 (0.44)	13.87 (0.19)	0.29 (0.10)	0.38 (0.10)	1.89 (0.10)	4.53 (0.11)	1.85 (0.10)	1.83 (0.10)	429 (31)	2.63
<i>P = 2.188 kbar, T = 994°C</i>												
MP18 I	-1.28 (0.43)	6.07 (0.08)	77.32 (0.40)	13.73 (0.17)	0.46 (0.10)	0.34 (0.10)	2.02 (0.10)	4.23 (0.10)	1.89 (0.10)	3.37 (0.12)	284 (47)	2.45
MP18 II	-0.93 (0.44)	6.11 (0.11)	77.08 (0.59)	13.68 (0.25)	0.92 (0.10)	0.32 (0.10)	2.07 (0.10)	4.12 (0.14)	1.80 (0.10)	4.24 (0.19)	147 (29)	2.17
MP18 IV	-1.34 (0.37)	7.13 (0.07)	77.51 (0.49)	13.69 (0.21)	0.26 (0.10)	0.33 (0.10)	2.07 (0.10)	4.26 (0.12)	1.87 (0.10)	3.82 (0.15)	453 (98)	2.66
MP18 V	-1.75 (0.42)	7.92 (0.06)	77.22 (0.59)	13.80 (0.25)	0.16 (0.10)	0.31 (0.10)	2.10 (0.10)	4.53 (0.14)	1.88 (0.0)	2.62 (0.15)	660 (38)	2.82
MP18 VI	-2.15 (0.30)	8.35 (0.06)	77.31 (0.46)	13.78 (0.20)	0.09 (0.10)	0.34 (0.10)	2.12 (0.10)	4.52 (0.11)	1.83 (0.10)	1.92 (0.10)	907 (41)	2.96
MP18 VII	-1.85 (0.30)	8.55 (0.06)	77.52 (0.58)	13.64 (0.24)	0.12 (0.10)	0.31 (0.10)	2.09 (0.10)	4.46 (0.14)	1.85 (0.10)	2.34 (0.14)	1154 (21)	3.06
MP18 XII	-0.75 (0.07)	6.33 (0.07)	76.70 (0.51)	13.67 (0.22)	1.09 (0.10)	0.33 (0.10)	2.07 (0.10)	4.24 (0.12)	1.90 (0.10)	4.82 (0.49)	178 (57)	2.25
MP18 SN ⁶	-0.74	6.00 (0.12)	76.33 (0.44)	13.72 (0.19)	1.50 (0.10)	0.32 (0.10)	2.03 (0.10)	4.20 (0.10)	1.90 (0.10)	4.47 (0.61)	116 (23)	2.06
PIN18 IX	-2.26 (0.77)	8.11 (0.06)	77.00 (0.77)	13.51 (0.32)	0.27 (0.10)	0.33 (0.10)	2.09 (0.13)	4.89 (0.19)	1.90 (0.12)	1.80 (0.54)	698 (64)	2.84
<i>P = 2.068 kbar, T = 996°C</i>												
MP19 VII	1.06 (0.19)	9.59 (0.19)	75.98 (0.48)	13.43 (0.20)	1.94 (0.10)	0.33 (0.10)	2.07 (0.10)	4.36 (0.12)	1.89 (0.10)	3.01 (0.46)	$\leq D$	—
MP19 II	1.46 (0.12)	8.67 (0.25)	76.06 (0.49)	13.41 (0.20)	2.11 (0.10)	0.33 (0.10)	2.11 (0.10)	4.13 (0.11)	1.85 (0.10)	4.17 (0.43)	$\leq D$	—
MP19 III	1.25 (0.24)	9.34 (0.27)	76.36 (0.51)	13.54 (0.22)	1.46 (0.10)	0.35 (0.10)	2.08 (0.10)	4.28 (0.12)	1.91 (0.10)	3.57 (0.71)	132 (70)	2.12
MP19 IV	0.81 (0.41)	9.86 (0.23)	76.23 (0.68)	13.54 (0.29)	1.50 (0.10)	0.32 (0.10)	2.15 (0.11)	4.33 (0.16)	1.93 (0.11)	2.61 (0.77)	$\leq D$	—
MP19 VI	-2.29	10.31	77.44 (0.88)	11.79 (0.34)	1.13 (0.11)	0.39 (0.10)	0.91 (0.10)	3.17 (0.18)	5.17 (0.23)	0.25 (0.25)	$\leq D$	—
MP19 IX	1.59 (0.12)	8.20 (0.31)	76.91 (0.53)	13.83 (0.22)	1.71 (0.10)	0.33 (0.10)	2.08 (0.10)	3.28 (0.11)	1.85 (0.10)	4.53 (0.52)	$\leq D$	—
MP19 X	1.59 (0.19)	8.25 (0.31)	76.28 (0.80)	13.47 (0.33)	2.09 (0.13)	0.35 (0.10)	2.14 (0.13)	3.74 (0.18)	1.92 (0.13)	5.15 (1.08)	122 (70)	2.09
MP19 XI	1.59 (0.09)	8.03 (0.29)	76.40 (0.50)	13.57 (0.21)	1.84 (0.10)	0.33 (0.10)	2.06 (0.10)	3.90 (0.11)	1.89 (0.10)	4.65 (0.37)	145 (21)	2.16
MP19 XII	1.53 (0.18)	6.25 (1.23)	75.36 (0.48)	13.44 (0.20)	3.06 (0.10)	0.32 (0.10)	2.03 (0.10)	3.85 (0.11)	1.93 (0.10)	4.44 (0.73)	167 (23)	2.22
MP19 XIII ⁶	1.05	9.55 (0.30)	74.93 (0.48)	13.49 (0.21)	3.40 (0.11)	0.32 (0.10)	1.97 (0.10)	4.04 (0.11)	1.85 (0.10)	2.84 (0.79)	$\leq D$	—
PIN19 VIII	0.88 (0.32)	9.81 (0.21)	78.32 (0.62)	12.99 (0.25)	0.80 (0.10)	0.25 (0.10)	1.29 (0.10)	3.49 (0.13)	2.85 (0.12)	2.60 (0.61)	$\leq D$	—

Table 5: continued

Charges	ΔNNO^1	ΔFFS^2	SiO_2	Al_2O_3	FeO^3	MgO	CaO	Na_2O	K_2O	H_2O^4	S (ppm) ⁵	log S
P = 1.992 kbar, T = 987°C												
MP14 I	2.23 (0.09)	7.89 (0.14)	76.71 (0.66)	13.67 (0.28)	2.20 (0.12)	0.34 (0.10)	1.30 (0.10)	3.93 (0.15)	1.84 (0.10)	2.80 (0.18)	674 (104)	2.83
MP14 II	2.18 (0.06)	8.00 (0.09)	76.88 (0.44)	13.73 (0.18)	2.13 (0.10)	0.32 (0.10)	1.24 (0.10)	3.90 (0.10)	1.79 (0.10)	2.67 (0.11)	563 (77)	2.75
MP14 III	1.85 (0.14)	8.85 (0.18)	77.60 (0.83)	13.90 (0.35)	0.92 (0.11)	0.35 (0.10)	1.15 (0.10)	4.16 (0.19)	1.92 (0.13)	2.11 (0.20)	734 (49)	2.87
MP14 IV	2.00 (0.31)	9.32 (0.12)	77.60 (0.45)	14.02 (0.19)	0.63 (0.10)	0.33 (0.10)	1.18 (0.10)	4.34 (0.11)	1.89 (0.10)	1.78 (0.10)	771 (31)	2.89
MP14 VI ⁶	2.03 (0.06)	8.49 (0.30)	75.88 (0.77)	13.35 (0.32)	3.17 (0.16)	0.30 (0.10)	1.62 (0.11)	3.88 (0.18)	1.79 (0.12)	2.53 (0.20)	399 (88)	2.60
MP14 VII	2.36 (0.03)	10.88 (0.09)	76.08 (0.58)	13.48 (0.24)	3.26 (0.12)	0.31 (0.10)	1.15 (0.10)	3.90 (0.13)	1.81 (0.10)	3.19 (0.16)	645 (75)	2.64
PI14 VIII	2.14 (0.06)	—	78.66 (0.48)	12.99 (0.19)	0.89 (0.10)	0.23 (0.10)	1.19 (0.10)	3.28 (0.10)	2.76 (0.10)	2.64 (0.12)	186 (14)	2.27

Numbers in parentheses indicate 1 standard deviation of replicate analyses.

¹ f_{O_2} normalized to the NNO buffer (see Table 4).

² f_{S_2} normalized to the FFS buffer (see Table 4).

³Total Fe, given as FeO.

⁴Water content determined using the by-difference method.

⁵Total sulphur content. D is the detection limit (80 ppm).

⁶Pyrrhotite-added charges.

⁷Magnetite-added charges.




Fig. 3. Stability domains of sulphur-bearing phases as a function of fO_2 and temperature. Phase abbreviations: anh, anhydrite; po, pyrrhotite; S, immiscible Fe–S–O liquid; mt, magnetite.




Fig. 4. Fugacities of H_2S and SO_2 fluid species versus the corresponding fO_2 for all charges for which fS_2 could be determined, calculated using equations (7) and (8) (see Table 4 and text). The increase in fO_2 favours SO_2 relative to H_2S species in the fluid phase. At any given fO_2 , the spread in either fH_2S or fSO_2 reflects the variations in fS_2 that are a result of the varying amounts of sulphur loaded to the charge.

Figure 4 shows the calculated fugacities of both H_2S and SO_2 species in the fluid for all charges for which fS_2 is available, plotted against the corresponding fO_2 . The fugacity of H_2S (fH_2S) ranges from 10 to 4.977 kbar, whereas that of SO_2 (fSO_2) covers a wider range, from 10^{-4} to 7.804 kbar. The crossover between fH_2S and fSO_2 occurs at around $NNO+1$. In fact, it is only at $fO_2 > NNO+2$ and at 930 and 1000°C that fSO_2 is several orders of magnitude larger than fH_2S .

Major element glass composition

Glasses obtained using the Pinatubo composition, which is richer in alkalis than that from Mt Pelée (Table 1), display similar sulphur concentrations when held under similar fO_2 and fS_2 , except at high fO_2 , where Pinatubo




Fig. 5. FeO_{tot} vs melt-sulphur content of experimental glasses at all investigated temperatures for reduced and oxidized conditions. The latter correspond in general to anhydrite-bearing charges. Note the general leftward displacement relative to the starting composition (dashed line at 2 wt %) of most charges. The arrow indicates that, below $NNO+1$, the general increase in melt-sulphur content is a result of the increase in fH_2S (and temperature).

charges are slightly richer in sulphur than the Mt Pelée ones. Overall, however, because Pinatubo charges do not depart significantly from the trends defined by the Mt Pelée ones, in the following, both groups are considered together. The glasses are rhyolitic, with SiO_2 contents predominantly in the range 76–78 wt % and $Al_2O_3 = 13 \pm 1$ wt %. Variations in the concentrations of other elements affect primarily Fe and Ca, as both elements are removed by either sulphide or sulfate crystallization. In addition, at 930 and 800°C, the crystallization of plagioclase, quartz and pyroxene also affects the melt chemistry, most notably its CaO and K_2O concentrations (Tables 4 and 5). In particular, at 800°C, a continuous increase in K_2O arises as a result of the dominantly incompatible behaviour of this element (see MP16 series, Table 5). At all values of fS_2 , fO_2 and T , a broad negative correlation can be established between FeO and melt sulphur content in sulphide-bearing charges (Fig. 5). All glasses to which elemental sulphur was added and which were held at $fO_2 < NNO+1$ are depleted to various extents in iron compared with the starting composition (2 wt % FeO), because of crystallization of sulphide (Fig. 5). In contrast, most charges with sulphur added as pyrrhotite have glasses richer in iron than other charges run under the same T , fO_2 but loaded with elemental sulphur (see Table 5). Charges doped with magnetite also display higher iron contents than undoped ones held at the same T and fO_2 . It is important to note that most glasses with low to very low iron contents display the highest sulphur contents, whereas those with the highest iron content have the lowest sulphur content, often below the detection limit (Fig. 5). Glasses of charges held at an $fO_2 > NNO+1$ display consistently higher iron contents than all other charges having similar amounts of dissolved




Fig. 6. Sulphur concentration in glass as a function of fO_2 and fS_2 at 930°C. Dotted lines are contours of iso- fS_2 , labelled with ΔFFS in bold and corresponding fS_2 in parentheses.

sulphur (Fig. 5)—an effect of the crystallization of anhydrite in lieu of pyrrhotite at high fO_2 .

Solubility of sulphur as a function of intensive parameters


The interrelationships between melt sulphur content, fS_2 and fO_2 at 930°C, where most of the data were obtained, are shown in Fig. 6. The solubility of sulphur is clearly a function of both fO_2 and fS_2 . Below $NNO+1$, for a fixed fO_2 , the solubility of sulphur increases with fS_2 . Above $NNO+1$, the data are scarce but suggest the same trend of increasing sulphur content with increasing fS_2 . For any given fS_2 , the sulphur solubility decreases when fO_2 increases, so as to reach a minimum between NNO and $NNO+1$. The data suggest that this minimum in sulphur solubility moves towards more oxidizing conditions when fS_2 increases.

The relationships between melt sulphur concentration (S_{melt}), fS_2 , fH_2S , fSO_2 for various fO_2 and at the three investigated temperatures are shown in Fig. 7. There is a general systematic increase in S_{melt} with fS_2 , most clearly seen at low fO_2 ($< NNO+1$). As a rule, an increase in fO_2 leads to a decrease in S_{melt} at fixed fS_2 . This effect is most apparent at 800 and 930°C. Exceptions to this behaviour concern experiments performed at 930 and 1000°C in the fO_2 range $NNO+0.7$ to $NNO+1$, where the S_{melt} content displays few variations, being close to the detection limit (Fig. 7). No clear trend appears when using fSO_2 in place of fS_2 (Fig. 7), even under high fO_2 , where SO_2 is the dominant sulphur-bearing species in the fluid. In contrast, S_{melt} correlates strongly with fH_2S . At 800°C, the two data groups at $NNO-1.4$ and $NNO+1.2$ merge into a single, well-defined trend, whereas using

fS_2 leads to two distinct series. Similarly, at 930°C, a general single positive correlation emerges, with much less dispersion than when fS_2 is used as a variable. At 1000°C, a strong positive correlation is clearly defined only for an fO_2 of $NNO-1.4$. At all temperatures, S_{melt} exceeds 200 ppm once fH_2S is above a few hundred bars. At $NNO-1.4$, when the mole fraction of S (X_S , calculated using common rock-forming oxides such as SiO_2 , Al_2O_3 , etc.) is used in place of concentration (S_{melt}), the correlations defined with fH_2S appear to be linear at all temperatures (Fig. 8). Under high fO_2 , a broad positive correlation exists between X_S and fSO_2 , but with significant scatter and no apparent temperature effect can be resolved, unlike at low fO_2 . The scatter is because of the difficulties in precisely determining fS_2 under high fO_2 conditions.

Sulphur speciation in hydrous rhyolite melt

Since the pioneering work of Fincham & Richardson (1954), it is common to envision sulphur dissolution in anhydrous silicate melts either as sulphide (low- fO_2) or as sulphate (high- fO_2) species, depending on prevailing redox state, through the following two reactions (see Carroll & Webster, 1994):



and



where S^{2-} and SO_4^{2-} refer respectively to sulphide and sulphate species dissolved in the melt and O^{2-} represents oxygen anions in the silicate melt. A quantitative assessment of reactions (12) and (13) has been so far limited by our inability to rigorously evaluate the proportion, as well as the activity–composition relationships, of O^{2-} in complex hydrous aluminosilicate melts such as magmatic liquids. The strong correlation existing between Fe and S_{melt} in both natural and experimental anhydrous glasses suggests that S^{2-} exchanges preferentially with O^{2-} bound to ferrous iron in melts under low fO_2 , whereas at high fO_2 , anhydrite precipitation suggests that the exchange primarily takes place with oxygens associated to Ca. In hydrous melts, the same reactions are supposed to operate (Carroll & Webster, 1994). In particular, under low fO_2 , sulphur dissolution might happen through a reaction such as (Carroll & Webster, 1994)



This reaction predicts that sulphur solubility in iron-bearing hydrous melt should increase with the activity of FeO, which is contrary to what our experimental data





Fig. 7. Relationships between melt-sulphur concentration (S_{melt}), and fS_2 , fH_2S , fSO_2 for various fO_2 , at the three investigated temperatures. For clarity at 930°C, the data have been grouped into three fO_2 intervals, each broadly corresponding to a different type of S-bearing phase (as fO_2 increases: pyrrhotite, occurrence of an immiscible Fe–S–O liquid, and anhydrite).

show (Fig. 5). However, based on experimental as well as on energetic and volumetric considerations, in particular the similarity of H_2O and H_2S molecules, Burnham (1979) proposed that sulphur dissolution into iron-poor hydrous rhyolitic melts is analogous to the H_2O solution reaction, and can be described by the following reaction:



Burnham (1979) reported a sulphur solubility of 1700 ppm at 2 kbar, 850°C in a hydrous, pyrrhotite saturated, rhyolitic melt (initial iron content of 0.3 wt %). This sulphur concentration is higher than the maxima attained in our study, and, upon completion of the run, it was found that, essentially, all the iron initially present in the system is fixed in pyrrhotite (Burnham, 1979)—an observation that corroborates our findings at low fO_2 . The very fact that similar, or even higher, sulphur concentrations compared with those achieved in the present work have been reported in either H_2O -bearing (Keppler, 1999) or H_2O -free (Mysen & Popp,

1980) haplogranite or albite melts (i.e. melts without iron) shows that iron is not instrumental in achieving high sulphur concentration in silicate melts. That, first, a negative correlation exists between dissolved iron and sulphur (Fig. 5), secondly, the glasses having the highest sulphur concentration are also those having the lowest iron content, close to the detection limit and, thirdly, a strong positive correlation exists between fH_2S and S_{melt} all suggest that reactions such as (15) may better account for our experimental observations, at least those below $NNO+1$, rather than reactions in which iron-bearing species are involved. Under oxidizing conditions, the situation is less clear but sulphur dissolution in hydrous melts may occur via the following reaction:



Unfortunately, a rigorous evaluation of reactions (14)–(16) is hampered primarily by the lack of knowledge about the nature of sulphur species dissolved in silicate melts, as well as of the activity–composition relationships




Fig. 8. X_i - f_i relationships. (a) Under reduced conditions ($\Delta\text{NNO} \leq 0$), where the dominant S-bearing species in the fluid is H_2S . (b) Under oxidizing conditions ($\Delta\text{NNO} \geq 1.0$), where the dominant S-bearing species in the fluid is SO_2 . Dashed curves are sulphur solubilities calculated using the thermodynamic model developed in the text.

of melt components, and these are clearly two main avenues for future research on the behaviour of sulphur in silicate melts.

Thermodynamic model of sulphur solubility in hydrous and iron-poor rhyolite melts

In this section, in the absence of specific speciation data and on the basis that H_2S and SO_2 are the major sulphur-bearing species in the coexisting fluid phase, we assume that the total dissolved sulphur in the melt results from the addition of sulphur dissolved from H_2S and SO_2 dissolution reactions. The extent of each dissolution reaction is controlled by the magnitude of both H_2S and SO_2 fugacities in the system. We thus have

$$X_S = X_{\text{reduced}} + X_{\text{oxidized}} \quad (17)$$

where X_{reduced} and X_{oxidized} represent the mole fractions of sulphur species dissolved under the reduced and oxidized forms, hereafter referred to as $X_{\text{H}_2\text{S}_{\text{melt}}}$ and $X_{\text{SO}_2_{\text{melt}}}$, respectively. Regardless of the actual sulphur species in the melt, the data allow us to evaluate the

standard thermodynamic parameters of the following heterogeneous reactions between fluid and melt (e.g. Spera & Bergman, 1980):



and



whose equilibrium constants are

$$K_{18} = f_{\text{H}_2\text{S}_{\text{fluid}}} / a_{\text{H}_2\text{S}_{\text{melt}}}$$

and

$$K_{19} = f_{\text{SO}_{2\text{fluid}}} / a_{\text{SO}_{2\text{melt}}}$$

Assuming that $a_i = X_i$, then

$$K_{18} = f_{\text{H}_2\text{S}_{\text{fluid}}} / X_{\text{H}_2\text{S}_{\text{melt}}}$$

and

$$K_{19} = f_{\text{SO}_{2\text{fluid}}} / X_{\text{SO}_{2\text{melt}}}$$

To evaluate the above reactions, we proceed by considering separately charges at $f\text{O}_2 < \text{NNO}$, in which H_2S is the dominant sulphur-bearing species in the fluid, and then charges at $f\text{O}_2 > \text{NNO}+1.5$, in which SO_2 is the dominant sulphur-bearing species. Under those conditions, the right-hand side of equation (17) simplifies to either $X_{\text{H}_2\text{S}_{\text{melt}}}$ ($f\text{O}_2 < \text{NNO}$) or $X_{\text{SO}_{2\text{melt}}}$ ($f\text{O}_2 > \text{NNO}+1.5$).

H_2S dissolution

At all temperatures and $f\text{O}_2 \leq \text{NNO}$, the relationships between X_S and $f\text{H}_2\text{S}$ is within error linear (Fig. 8) and, thus, the solubility of H_2S in rhyolite melt obeys Henry's Law, i.e. at fixed P and T ,

$$f\text{H}_2\text{S} = X_{\text{H}_2\text{S}_{\text{melt}}} k_H \quad (20)$$

where k_H is the Henry's Law constant and is obviously equivalent to K_{18} . As shown in Fig. 8, the $f\text{H}_2\text{S}$ - $X_{\text{H}_2\text{S}_{\text{melt}}}$ relationships display well organized trends with temperature. Departure from the simple Henrian behaviour, because of both temperature and pressure variations, can be accounted for by using the following classical equation (e.g. Fogel & Rutherford, 1990):

$$\ln(f\text{H}_2\text{S}/X_{\text{H}_2\text{S}_{\text{melt}}}) = \ln k_{H(\text{Pr},\text{Tr})} + (\Delta H/R)(1/T - 1/\text{Tr}) + (V_{\text{H}_2\text{S}}(P - P_r))/RT \quad (21)$$

which, upon rearrangement, gives

$$X_{\text{H}_2\text{S}_{\text{melt}}} = f\text{H}_2\text{S}/k_{H(\text{Pr},\text{Tr})} \exp[-(\Delta H/R)(1/T - 1/\text{Tr}) - (V_{\text{H}_2\text{S}}(P - P_r))/RT] \quad (22)$$

in which T_r and P_r are the reference temperature and pressure, respectively, ΔH is the heat of H_2S dissolution into the melt, R is the gas constant and $V_{\text{H}_2\text{S}}$ the partial molar volume of H_2S melt species. T_r and P_r were, respectively, chosen to be 930°C , as most of the data were gathered along or close to this isotherm, and 0 bar. Equation (22) has been extensively used in the thermodynamic modelling of H_2O and CO_2 solubilities in various silicate–melt compositions (e.g. Stolper *et al.*, 1987; Fogel & Rutherford, 1990; Blank *et al.*, 1993). Non-linear least-squares regression using charges with an $f\text{O}_2 \leq \text{NNO}-1.4$ gives the following values: $k_{\text{H},930^\circ\text{C}} = 0.801 \cdot 10^6$ bar, $\Delta H = 54.85$ kJ/mol and $V_{\text{H}_2\text{S}} = 26.03$ cm³/mol. The Henry constant is similar to that found for CO_2 dissolution in rhyolite melt ($1.106 \cdot 10^6$ bar, Fogel & Rutherford, 1990). However, contrary to both H_2O and CO_2 species, the enthalpy of H_2S solution is positive in sign, which simply means that, at a given $f\text{H}_2\text{S}$, temperature increases the solubility of sulphur. Although the partial molar volume of H_2S in rhyolite melt appears to be similar to that of H_2O , we stress that our data cover a range in $f\text{H}_2\text{S}$ too small to allow extraction of volume properties with great accuracy. In fact, the investigated pressure domain falls in the range where sulphur exhibits essentially linear dependence on $f\text{H}_2\text{S}$, i.e. where $V_{\text{H}_2\text{S}}$ plays a negligible role in equation (22).

SO₂ dissolution

At $f\text{O}_2 > \text{NNO}+1.5$, because of the difficulties in retrieving fluid species fugacities in the oxidized range, the dispersion in the $X_{\text{S}}-f\text{SO}_2$ (Fig. 8) plot is much more severe than for H_2S , and it can only be assumed that sulphur dissolution at high $f\text{O}_2$ also follows Henry's Law. Linear regression of the data obtained at an $f\text{O}_2 > \text{NNO}+1.5$ gives the following value for the Henry constant: $K_{\text{SO}_2} = K_{19} = 4.13 \cdot 10^6$ bar.

Sources of error

The calculated versus measured sulphur mole fractions for various $f\text{H}_2\text{S}$ are shown in Fig. 9 for charges held at an $f\text{O}_2$ of $\text{NNO}-1.4$ or lower. Although the model correctly reproduces the measured values, there is a substantial scatter around the 1:1 correlation line. Besides the obvious fact that not all dissolved sulphur needs to be in the H_2S form, one major source of error that affects the derivation of thermodynamic parameters comes from the calculated values of $f\text{H}_2\text{S}$. In experimental studies designed to calibrate H_2O or CO_2 solubility models, the fluid species fugacities are calculated from equations of state in which uncertainties in fugacities essentially come from uncertainties in P and T measurements. Here, the H_2S fugacities are also affected by this source of




Fig. 9. Experimental versus calculated mole fractions of sulphur for charges at an $f\text{O}_2$ of below $\text{NNO}-1.4$, using the thermodynamic model developed in the text.

uncertainty, but also by (1) the analytical uncertainty attached to the determination of pyrrhotite composition and (2) the lack of well-calibrated EOS for sulphur-bearing volatile species. The analytical uncertainty in measured $f\text{S}_2$ is typically 0.2 log units (Table 3). Taking charge MP15V as an example, its measured $f\text{S}_2$ is 5.46 bar ($\log f\text{S}_2 = 0.74$), to which corresponds an $f\text{H}_2\text{S}$ of 4.977 kbar, given the $f\text{H}_2$ of 27 bar. If, instead, $f\text{S}_2$ is 8.66 bar ($\log f\text{S}_2 = 0.94$), then the calculated $f\text{H}_2\text{S}$ would be 6.265 kbar or *c.* 25% higher absolute. There is thus no doubt that the thermodynamic quantities derived above need to be confirmed, in particular by designing specific experimental procedures to achieve equilibration of rhyolite melts with pure H_2S fluid (or SO_2) and by exploring the behaviour at higher $f\text{H}_2\text{S}$ and $f\text{SO}_2$ than those attained in the present study to constrain more rigorously the partial molar volumes of H_2S and SO_2 -related species (see below also). The data suggest, however, that, under low $f\text{O}_2$ and in iron-poor silicate melts, H_2S dissolution may occur through formation of HS-like complexes, and that at high $f\text{O}_2$, SO_2 reacts with OH^- groups to yield SO_4^{2-} . Clearly, however, there is a need for more direct information (spectroscopic) about potential S complexes present in silicate melts (e.g. Winther *et al.*, 1998), as the common peak-shift method is unable to distinguish between S^{2-} -associated with Fe^{2+} from those linked to H^+ cations.

Solubility minimum and speciation

Despite these caveats, the thermodynamic properties derived above for H_2S and SO_2 dissolution in hydrous rhyolite melt can be combined to compute the behaviour of total dissolved sulphur, $X_{\text{S}} (= X_{\text{H}_2\text{S}\text{melt}} + X_{\text{SO}_2\text{melt}}$, as defined above in equations (18)–(22) under various T and $f\text{O}_2$ conditions. An Excel spreadsheet, which calculates the fugacities of S_2 , H_2S , SO_2 and H_2O , species at fixed P , T , S_{melt} and $f\text{O}_2$, using the thermodynamic approach




Fig. 10. Application of the thermodynamic model. (a) Calculated total dissolved sulphur (S_{ppm}) against fO_2 (ΔNNO) at 930°C and for an fS_2 of $\Delta\text{FFS}+7$ ($\log fS_2 = -0.75$), $\Delta\text{FFS}+8$ ($\log fS_2 = 0.24$), and $\Delta\text{FFS}+9$ ($\log fS_2 = 1.24$). All calculations have been performed with $fH_2O = 1$ kbar, which corresponds to a melt-water content of ≈ 4.5 wt %. (b) Comparison between observed and calculated melt sulphur contents for all charges in which fS_2 could be estimated. (c) Calculated proportions of S^{6+} at 1100 and 800°C, as a function of ΔNNO . The gray squares represent the experimental data of Carroll & Rutherford (1988).

detailed above, is available upon request from the second author. The results of such calculations are shown in Fig. 10a for 930°C, 2 kbar, several fS_2 within the fO_2 range $\text{NNO}-2$ to $\text{NNO}+3$. As shown in Fig. 10a, the

model reproduces the classical inverted bell-shaped pattern of sulphur solubility with varying fO_2 . At all fS_2 , the minimum calculated centers at $\text{NNO}+1$, in good agreement with experimental observations in the present (Fig. 6) as well as in other studies (see Carroll & Webster, 1994). It shows also that as fS_2 decreases, this bell-shaped pattern opens wide to the extent of becoming almost flat at very low values of fS_2 . Specifically, at 930°C, any fS_2 lower than 0.17 bar ($\text{FFS}+7$) results in no apparent variations in sulphur concentrations with varying fO_2 (note that the curve still has a bell-shaped pattern, even at very low sulphur concentrations, but the concentration scale over which it occurs is well below the detection limit of EPMA). Conversely, an fS_2 of 18 bar ($\text{FFS}+9$) leads to a strongly asymmetrical and steep-sided shape of the sulphur concentration curve with varying fO_2 . The calculated melt-sulphur concentrations for all charges for which fS_2 and fH_2 are available are plotted against those measured in Fig. 10b. The model reproduces well the experiments in the high concentration range, above 200 ppm, which reflects, in part, that the analytical accuracy is better at high concentration.

Given the above model assumptions, we can also calculate the proportions of S^{2-} and S^{6+} melt species, assuming that S^{2-} and S^{6+} abundances are fixed by H_2S and SO_2 dissolution reactions (18) and (19), respectively. Figure 10c shows the proportion of S^{6+} calculated for 1100 and 800°C isotherms. At a given P , T and fO_2 , the fS_2 has no effect on species proportions. In contrast, temperature does affect the calculated proportions so that lower temperature increases the proportion of sulphide species at any fixed fO_2 . As shown in Fig. 10c, the calculated pattern of species proportion broadly conforms with that obtained by Carroll & Rutherford (1988) from peak-shift measurements carried out on a series of synthetic sulphur-bearing glasses, annealed under various redox conditions in the temperature range 920–1150°C. The difference between model and observations may be a result of the fact that melt compositions synthesized by Carroll & Rutherford (1988) are less silicic than those of the present work.

Empirical model of sulphur solubility in rhyolite melts

The application of the thermodynamic model presented above requires the determination of both fH_2S and fSO_2 . These two quantities are not usually measured on magmatic rocks, in contrast to both fO_2 and fS_2 , which are more easily estimated from Fe–Ti oxides or sulfide barometry (Whitney, 1984). Although derivation of fSO_2 from fO_2 and fS_2 is straightforward, calculation of fH_2S requires fH_2 to be known in addition to fS_2 or, alternatively, the determination of fH_2O if fO_2 is known. This supplementary step introduces an additional source of




Fig. 11. Comparison between measured and calculated sulphur fugacities using the empirical model [equation (24)]. See text for explanation.

uncertainty, that is, the determination of melt water content from which $f_{\text{H}_2\text{O}}$ is calculated. To overcome this potential problem and given the restricted compositional range displayed by the experimental glasses, we have derived an empirical model in which T , f_{O_2} and f_{S_2} are the only required input parameters: such a model is simpler in its application and, therefore, of more direct petrological use. Given that many arc magmas have f_{O_2} below $\text{NNO}+1.5$, we have restricted the fitting procedure to charges lying below this f_{O_2} threshold: this has the advantage that only the charges for which f_{S_2} is best known are taken into account. Considering the different effects of f_{O_2} , f_{S_2} and T on the melt–sulphur content illustrated above, the experimental data were fitted by least-squares regression to the following equation, taking into account only glasses with sulphur-content higher than the detection limit:

$$\log S \text{ (ppm)} = 0.001 T (^{\circ}\text{C}) - 0.2567 \Delta\text{NNO} + 0.1713 \Delta\text{FFS} + 0.0034 \Delta\text{NNO} \times \Delta\text{FFS} \quad (23)$$

or

$$\frac{\log S - 0.001 T + 0.2567 \Delta\text{NNO}}{0.1713 + 0.0034 \Delta\text{NNO}} = \Delta\text{FFS}. \quad (24)$$

As with f_{O_2} , referencing f_{S_2} to a standard solid buffer such as FFS removes the temperature dependence and improves the quality of the fit. Knowing the sulphur content of a glass, the temperature and f_{O_2} of the system, equation (24) allows calculation of f_{S_2} . Figure 11 shows model performances for all charges held at an $f_{\text{O}_2} < \text{NNO}+1.5$. Equation (24) reproduces observed values to within 1 log units at $\Delta\text{FFS} > 6$, apart from a few charges in the f_{O_2} range NNO to $\text{NNO}+1.5$. At lower ΔFFS , the model tends to underestimate f_{S_2} —a

reflection of the difficulty in measuring sulphur content in glasses in the low concentration range (<100 ppm). We stress that equation (24) is merely a convenient mathematical way to describe the interdependence between the various parameters that control the sulphur solubility in rhyolite melts at 2 kbar and it has no thermodynamic meaning.

DISCUSSION

Fluid and melt-sulphur relationships

The general coherency of the experimental trends obtained between intensive (f_{O_2} , f_{S_2}) and extensive (S in melt) parameters shows that the adopted experimental procedure has been successful in either accurately controlling f_{O_2} or measuring f_{S_2} . The observed solubility trends can be interpreted in terms of different solubility behaviour of the different S-species present in the coexisting fluid. Because the fluid phase composition is sensitive to redox conditions, any change in f_{O_2} profoundly affects the sulphur solubility. The more soluble sulphur species appear to be H_2S and SO_2 . The sulphur behaviour in hydrous silicate melts can thus be modelled using a simple thermodynamic approach, similar to that employed for characterizing the solubilities of both H_2O and CO_2 volatiles in silicate melts, considering the additive effects of two basic dissolution reactions involving either H_2S or SO_2 fluid species. At fixed P and T , Henry's Law is obeyed for H_2S and possibly SO_2 species, up to fugacities of 5 and 8 kbar, respectively. At equal fugacities, H_2S appears to be more soluble than SO_2 , and, hence, the asymmetrical behaviour of sulphur on both sides of the solubility minimum. This asymmetry is also a result of the differing PVT properties of H_2S and SO_2 fluid species. The MRK model predicts that SO_2 exhibits a more pronounced non-ideal behaviour than H_2S (that is higher fugacity coefficients), and thus the fugacities of either end-member (i.e. pure SO_2 or H_2S fluids at any fixed P and T) and their mixing behaviour will also control the shape of the solubility curve. In other words, even if both H_2S and SO_2 were to have identical Henry's Law constants, because the fugacities of pure H_2S or SO_2 differ at fixed P and T , this will ultimately impart an asymmetrical shape on the solubility curve with changing redox conditions. In this regard, it would be of interest to constrain the solubility of silicate melts with respect to H_2S and SO_2 by performing experiments with pure H_2S and SO_2 fluids. The minimum in solubility appears to lie in the redox range where H_2S and SO_2 fugacities in the fluid are approximately equal (Fig. 4), and MRK calculations show that S_2 is comparatively more abundant in this domain. This latter species is thus apparently weakly soluble in silicate melts and plays a minor role in the melt-sulphur budget.

The effect of iron

The experimental procedure adopted here does not allow independent variation of fS_2 and the melt-FeO content (Fig. 5). In particular, at any given temperature, it has not been possible to produce charges displaying the same fS_2 and fO_2 but with widely different iron-melt contents. Only charges belonging to the MP12 and MP7 series (Table 5) are suitable to test the effect of varying iron content on sulphur solubility under quasi-constant fS_2 and fO_2 , albeit for iron content below 1 wt % only. These charges also display a negative trend between iron and sulphur in melt. Such a negative correlation is corroborated by the iron-rich but sulphur-poor melt in all magnetite-doped charges (Table 5). Altogether, the above lines of evidence demonstrate unequivocally that an iron-rich rhyolite will not dissolve more sulphur than a iron-poor one under the experimental conditions explored here—a result that is at variance with what has been found in mafic melts (e.g. Haughton *et al.*, 1974). Clearly, addition of sulphur to a silicic magma in the stability field of sulphide has the main effect of crystallizing sulphide, which removes both iron and sulphur from the melt (and fluid). This scavenging of sulphur results in low fS_2 , as long as there is enough iron to produce pyrrhotite. One reason for the fundamental difference between mafic and felsic melts is that the liquidus temperature of S-free hydrous silicic melts is much lower than that of dry mafic melts. Experiments performed in mafic systems were performed at about 1200°C, where much of the iron remains in solution in the melt and where it may complex with sulphur, preventing extensive crystallization of either sulphide or oxides. In contrast, the experiments of the present study are performed at temperatures well below the liquidus of sulphides, as indicated by their occurrence in nearly all run products. It is mostly when $fO_2 > NNO+1.5$, or in the anhydrite stability field, that the negative trend observed between FeO and sulphur solubility becomes weaker, indicating conditions less appropriate for sulphide crystallization (Fig. 5). We note, in addition, that most of the 1 bar experiments investigated the sulphur behaviour without the involvement of H-bearing species (e.g. O'Neill & Mavrogenes, 2002), and, thus, the solubility mechanisms of sulphur in those studies are probably different from those proposed here.

The foregoing considerations show that the application of the present thermodynamic model to melt compositions other than metaluminous rhyolite, in particular to more mafic compositions, is not warranted. Studies carried out on the H_2O and CO_2 solubilities in various silicate-melt compositions have shown that the thermodynamic parameters, i.e. enthalpy of species dissolution and its partial molar volume, are strongly dependent on melt composition (see Holloway & Blank, 1994). This stems from the fact that dissolution mechanisms, and, thus, the energetics of solution, may widely vary with

melt composition. An obvious difference is that sulphur may complex with iron in melts with higher FeO contents than those of the present study, which may lead to significantly different solution parameters. Another example concerns the alkalis over alumina balance (i.e. peralkaline, peraluminous) which evidently affects the proportion of O^{2-} and, thus, the sulphur behaviour. Therefore, although the thermodynamic model appears to reproduce successfully first-order observations bearing on the sulphur behaviour in silicate melts, its quantitative use on melt compositions other than metaluminous rhyolite requires prior experimental investigation of ΔH and V_i values associated with any given melt composition.

The effect of temperature on the solubility of sulphur and its speciation

Temperature appears to exert a positive effect on the solubility of H_2S —a behaviour at variance with that of both CO_2 and H_2O , at least in the low-pressure range, whereas SO_2 appears to be less sensitive to this parameter. As Stolper *et al.* (1987) have emphasized, the bulk effect of temperature on volatile solubility in silicate melts is controlled by the energetic balance between the reactions controlling molecular (e.g. H_2S , SO_2 , H_2O , CO_2) and anion groups' solubility (e.g. HS^- , SO_4^{2-} , OH^- , CO_3^{2-}). In general, the solubility of molecular species decreases with T (Stolper *et al.*, 1987; Fogel & Rutherford, 1990), whereas the abundance of species formed by reaction with the melt is enhanced by temperature (see Zhang, 1999). Thus, the fact that under reduced conditions, the bulk sulphur solubility has a strong and positive temperature dependence might be interpreted as the dominant presence of S^{2-} or HS^- species over molecular H_2S in rhyolite melts, i.e. that, essentially, all dissolved H_2S reacts with the aluminosilicate melt framework to yield HS^- units. Our data favor a reaction such as (15), in which exchange between gaseous H_2S and hydroxyl groups is the dominant mechanism of sulphur incorporation into hydrous and iron-poor silicate melts. At high fO_2 , it is not yet possible to decipher the details of the solution reaction but if all SO_2 reacts with hydroxyl groups to form SO_4^{2-} units, then it might be expected that there is a positive effect of temperature on solubility under oxidizing conditions as well. Thus, compared with H_2O and CO_2 , which dissolve both physically and chemically, the case for sulphur is different, as its dissolution seems to proceed dominantly chemically, by reaction of gaseous species with the melt framework, keeping the abundance of molecular species at a marginal level. Another major difference is that, as far as we know, for fixed melt chemistry, both H_2O and CO_2 speciations are insensitive to redox conditions, whereas, clearly, this parameter is the major controlling factor of sulphur speciation in silicate melts.

Table 6: Application of the empirical model to natural silicic magmas

Rock	SiO ₂ (wt %)	T (°C)	ΔNNO	log <i>f</i> S ₂	S _{melt} (ppm)	ΔFFS _{mes} ¹	ΔFFS _{calc} ²
Bishop	77	770	0.1	-2.70	100	7.09	7.18
Krakatau	69	890	1.20	-0.39	130	7.72	8.68
Katmai	78	850	0.25	-1.50	68	7.20	6.29
St Helens	73	930	1.18	0.20	65	7.98	6.83
El Chichon	67	800	1.00	0.00	200	8.70	10.01

Data sources for *T*, ΔNNO, log *f*S₂, S_{melt} are from Hildreth (1979, 1983), Luhr *et al.* (1984), Whitney (1984), Rutherford *et al.* (1985), Anderson *et al.* (1989), Luhr (1990), Westrich *et al.* (1991), Lowenstern (1993) and Mandeville *et al.* (1996, 1998).

¹Measured *f*S₂ in the rock from pyrrhotite composition (Whitney, 1984) referenced to the FFS buffer (see text).

²Calculated *f*S₂ (referenced to the FFS buffer) from equation (24).

Although significant, the dependence on temperature appears to be low compared with the effects exerted by *f*O₂ and *f*S₂. Below NNO, the temperature dependence established in other studies performed on hydrous silicic melts (Carroll & Rutherford, 1987; Luhr, 1990) is lower than that reported here. For instance, the experiments of Luhr (1990) on the El Chichón trachyandesite show almost no effect of temperature on the melt S content between 800 and 1000°C at an *f*O₂ buffered by the quartz–fayalite–magnetite solid buffer (NNO–0.7). In contrast, at high *f*O₂, a strong positive temperature dependence has been evidenced by both Carroll & Rutherford (1987) and Luhr (1990). In these experiments, however, low-temperature runs are largely below the liquidus of the bulk composition used as starting material, which means that the residual melt composition changes significantly with rising temperature, unlike in the present study. In such experiments, the temperature effect on sulphur solubility may thus be hindered by the interplay of additional variables such as *a*SiO₂ or *f*S₂, even when performed at fixed redox conditions. Thus, it remains to be demonstrated whether the strong positive temperature dependence of sulphur solubility observed under oxidizing conditions in hydrous silicic melts by Carroll & Rutherford (1987) and Luhr (1990) is, indeed, solely because of temperature, or the combined effects of varying melt composition with temperature in addition to varying *f*S₂.

The effect of pressure

The last parameter of interest is pressure. Recent experimental studies have confirmed the earlier findings of Wendlandt (1982) that pressure decreases the sulphur solubility in dry mafic systems (Mavrogenes & O'Neill, 1999; Holzheid & Grove, 2002). By contrast, in hydrous systems, Luhr (1990) concluded that pressure has an opposite effect. Nevertheless, as for temperature, the experiments of Luhr (1990) in which pressure was varied involved also varying other parameters, in particular the

melt composition, and it cannot be ruled out that the observed increase in S_{melt} also reflects a change in melt chemistry and not only an intrinsic and positive effect of pressure. Our experiments suggest that one of the controlling parameters of sulphur solubility in H-bearing systems is *f*H₂S. By analogy with what happens with water, as total pressure decreases, *f*H₂S will decrease too, so that the melt sulphur solubility should decrease eventually, in keeping with the Henrian behaviour shown above. In support of this are the experiments of Katsura & Nagashima (1974), which show that a rhyodacite melt (66.7 wt % SiO₂) equilibrated at 1 bar and 1250°C, with an H–C–S–O gas mixture over an *f*O₂ range of NNO–5 to NNO+5, displays sulphur concentration always below 50 ppm. Thus, given that natural silicic magmas have eruption temperatures that are 300–400°C lower, and that they have higher silica contents than the rhyodacite experimental melts, it can be anticipated that their equilibrium sulphur concentration at near atmospheric conditions is close to the ppm level. That, at 1 bar, silicic compositions have much lower sulphur solubility than mafic ones is related to the fact that, under anhydrous conditions, the proportion of O²⁻ is significantly higher in the latter than in the former.

Application of the empirical model

The empirical model can be tested by applying it to eruption products for which common pre-eruptive parameters are well known (*P*, *T*, *f*O₂) in addition to *f*S₂ and the melt-sulphur concentration. These include the Bishop Tuff (0.76 Ma), Krakatau (1885), Katmai (1912), Mt St Helens (1980) and El Chichón (1982) eruptions. Pre-eruptive parameters, as well as calculated melt-sulphur fugacities, are listed in Table 6. Equation (25) reproduces observed sulphur fugacities to within 1 log unit, except for Mt St Helens. The origin of the disagreement for Mt St Helens might be because of the relatively low sulphur content measured, close to the detection limit

of EMPA: if, for instance, a pre-eruptive melt-sulphur content of 100 ppm is assumed, then the calculated ΔFFS is 7.78, or within 0.2 log units of the measured one by the pyrrhotite composition (Whitney, 1984). As the experimental melt compositions are metaluminous and because of the empirical nature of equation (25), the model should not be applied to either peraluminous or peralkaline rhyolites.

SUMMARY

(1) Experiments with controlled $f\text{O}_2$ and $f\text{S}_2$ have been performed in the temperature range 800–1000°C on rhyolite-melt compositions, to establish the effect of those variables on the solubility of sulphur. The $f\text{S}_2$ was measured either by the pyrrhotite ($f\text{O}_2 < \text{NNO}+1$) or by the fluid phase ($f\text{O}_2 > \text{NNO}+1.5$) compositions, using for the latter an MRK equation of state.

(2) At $f\text{O}_2 < \text{NNO}+1$ and temperatures of <1000°C, the addition of iron to a sulphur-bearing rhyolite melt promotes massive crystallization of pyrrhotite, which consequently removes most of the sulphur from the melt.

(3) At constant $f\text{O}_2$, the sulphur solubility increases with $f\text{S}_2$. Below $\text{NNO}+1$, there is a linear relationship between the melt-sulphur concentration and $f\text{H}_2\text{S}$, whereas above $\text{NNO}+1$, $f\text{SO}_2$ seems the dominant controlling factor.

(4) The sulphur solubility in rhyolite melts is a result of the dissolution of both H_2S and SO_2 fluid species. The strong positive dependence of solubility on temperature observed at $f\text{O}_2 < \text{NNO}+1$ suggests that the abundance of dissolved molecular H_2S is low and that most of the sulphur dissolves by chemical reaction with the aluminosilicate framework. SO_2 possibly behaves similarly but this needs to be checked by additional experimental work, with a more precise control of $f\text{S}_2$ under oxidizing conditions than that achieved in the present work.

(5) A simple thermodynamic model can be developed, adopting an approach similar to that used for modelling the solubilities of both H_2O and CO_2 in silicate melts. This model reproduces first-order observations concerning the sulphur solubility and speciation in silicate melts. Yet, there is an urgent need for rigorously determining the nature and abundance of different sulphur species dissolved in melts and their dependence on melt composition.

(6) An empirical model for metaluminous rhyolites that is easier to handle has been developed also. It may be used primarily to determine the $f\text{S}_2$ of common metaluminous silicic arc magmas.

ACKNOWLEDGEMENTS

This paper is a part of the PhD thesis of the first author. The experimental work has been partly financed by the

ENV4-CT96-0259 CE programme. Helpful comments were provided by Malcolm Rutherford on an early version of the manuscript. Thorough and helpful reviews of Jake Lowenstern, Leslie Baker and an anonymous reviewer are gratefully acknowledged.

REFERENCES

- Abraham, K. P. & Richardson, F. D. (1960). Sulphide capacities of silicate melts. Part II. *Journal of the Iron & Steel Institute* **196**, 313–317.
- Abraham, K. P., Davis, M. W. & Richardson, F. D. (1960). Sulphide capacities of silicate melts. Part I. *Journal of the Iron & Steel Institute* **196**, 309–312.
- Ancey, M., Bastenaire, F. & Tixier, R. (1978). Application des méthodes statistiques en microanalyse. In: Physique, E. D. (ed.) *Microanalyse, microscope électronique à balayage*. Orsay: Edition de physique, Université d'Orsay, pp. 323–347.
- Anderson, A. T., Newman, S., Williams, S. N., Druitt, T. H., Skirius, C. & Stolper, E. (1989). H_2O , CO_2 , Cl gas in Plinian and ash-flow Bishop rhyolite. *Geology* **17**, 221–225.
- Arnold, R. G. & Reicher, L. E. (1962). Measurement of the metal content of naturally occurring, metal deficient, hexagonal pyrrhotite by an X-Ray spacing method. *American Mineralogist* **47**, 105–111.
- Baker, L. I. & Rutherford, M. J. (1996a). Sulphur diffusion in rhyolite melts. *Contributions to Mineralogy and Petrology* **123**, 335–344.
- Baker, L. I. & Rutherford, M. J. (1996b). The effect of dissolved water on the oxidation state of silicic melts. *Geochimica et Cosmochimica Acta* **60**, 2179–2187.
- Blank, J., Stolper, E. M. & Carroll, M. (1993). Solubilities of carbon dioxide and water in rhyolitic melt at 850°C and 750 bars. *Earth and Planetary Science Letters* **119**, 27–36.
- Boorman, R. S. (1967). Subsolidus studies in the ZnS–FeS–FeS₂ system. *Economic Geology* **62**, 614–631.
- Breedveld, G. J. F. & Prausnitz, J. M. (1973). Thermodynamic properties of supercritical fluids and their mixture at very high pressures. *AIChE Journal* **19**, 783–796.
- Buchanan, D. L. & Nolan, J. (1979). Solubility of sulphur and sulphide immiscibility in synthetic tholeiitic melts and their relevance to Bushveld-complex rocks. *Canadian Mineralogist* **17**, 483–494.
- Buchanan, D. L., Nolan, J., Wilkinson, N. & De Villiers, J. P. R. (1983). An experimental investigation of sulfur solubility as a function of temperature in silicate melts. *Special Publication of the Geological Society of South Africa* **7**, 383–391.
- Burnham, C. W. (1979). The importance of volatiles constituents. In: Yoder, H. S. (ed.) *The Evolution of the Igneous Rocks: Fiftieth Anniversary Perspectives*. Princeton, NJ: Princeton University Press, pp. 439–482.
- Burnham, C. W., Holloway, J. R. & Davis, N. F. (1969). Thermodynamic properties of water to 1000°C and 10 000 bars. *Geological Society of America, Special Paper* **132**, 1–96.
- Carroll, M. R. & Rutherford, M. J. (1985). Sulphide and sulfate saturation in hydrous silicate melts. *Journal of Geophysical Research* **90**, C601–C612.
- Carroll, M. R. & Rutherford, M. J. (1987). The stability of igneous anhydrite: experimental results and implications for sulphur behavior in the 1982 El Chichón trachyandesite and other evolved magmas. *Journal of Petrology* **28**, 781–801.
- Carroll, M. R. & Rutherford, M. J. (1988). Sulphur speciation in hydrous experimental glasses of varying oxidation state: results from measured wavelength shifts of sulphur X-rays. *American Mineralogist* **73**, 845–849.

- Carroll, M. R. & Webster, J. D. (1994). Solubilities of sulfur, noble gases, nitrogen, chlorine, and fluorine in magmas. In: Holloway, J. R. & Carroll, M. R. (eds) *Volatiles in Magmas. Mineralogical Society of America, Reviews in Mineralogy* **30**, 231–280.
- Connolly, J. A. & Cesare, B. (1993). C–O–H–S fluid composition and oxygen fugacity in graphitic metapelites. *Journal of Metamorphic Geology* **11**, 379–388.
- Devine, J. D., Gardner, J. E., Brack, H. P., Layne, G. D. & Rutherford, M. J. (1995). Comparison of microanalytical methods for estimation of H₂O contents of silicic volcanic glasses. *American Mineralogist* **80**, 319–328.
- Ferry, J. M. & Baumgartner, L. (1987). Thermodynamic models of molecular fluids at the elevated pressures and temperatures of crustal metamorphism. In: Carmichael, I. S. E. & Eugster, H. P. (eds) *Reviews in Mineralogy. Mineralogical society of America*, 323–365.
- Fincham, C. J. B. & Richardson, F. D. (1954). Behaviour of sulphur in silicate and aluminate melts. *Proceedings of the Royal Society of London* **223**, 40–62.
- Fogel, R. A. & Rutherford, M. J. (1990). The solubility of carbon dioxide in rhyolitic melts: a quantitative FTIR study. *American Mineralogist* **75**, 1311–1326.
- Froese, E. & Gunter, A. E. (1976). A note on the pyrrhotite-sulphur vapor equilibrium. *Economic Geology* **71**, 1589–1594.
- Gaillard, F., Scaillet, B., Pichavant, M. & Bény, J. M. (2001). The effect of water and *f*O₂ on the ferric–ferrous ratio of silicic melts. *Chemical Geology* **174**, 255–273.
- Haughton, D. R., Roeder, P. L. & Skinner, B. J. (1974). Solubility of sulphur in mafic magmas. *Economic Geology* **69**, 451–466.
- Hildreth, W. (1979). The Bishop Tuff: evidence for the origin of compositional zonation in silicic magma chambers. In: Chapin, C. E. & Elston, W. E. (eds) *Ash-flow Tuffs. Geological Society of America, Special Paper* **180**, 43–75.
- Hildreth, W. (1983). The compositionally zoned eruption of 1912 in the Valley of Ten Thousand Smokes, Katmai National Park, Alaska. *Journal of Volcanology and Geothermal Research* **18**, 1–56.
- Holloway, J. R. (1977). Fugacity and activity of molecular species in supercritical fluids. In: Fraser, D. (ed.) *Thermodynamics in Geology*. Boston, MA: Kluwer Academic, pp. 161–181.
- Holloway, J. R. (1981). Volatile interactions in magmas. In: Newton, R. C., Navrotsky, A. & Wood, B. J. (eds) *Thermodynamics of Minerals and Melts. Advances in Physical Geochemistry* **1**, 273–283.
- Holloway, J. R. (1987). Igneous fluids. In: *Thermodynamic Modelling of Geological Materials: Minerals, Fluids and Melts. Mineralogical society of America, Reviews in Mineralogy* **17**, 211–232.
- Holloway, J. R. & Blank, J. (1994). Application of experimental results to C–O–H species in natural melts. In: Holloway, J. R. & Carroll, M. R. (eds) *Volatiles in Magmas. Mineralogical society of America, Reviews in Mineralogy* **30**, 187–230.
- Holloway, J. R., Pan, V. & Gudmundsson, G. (1992). High-pressure fluid-absent melting experiments in the presence of graphite: oxygen fugacity, ferric/ferrous ratio and dissolved CO₂. *European Journal of Mineralogy* **4**, 105–114.
- Holzheid, A. & Grove, T. L. (2002). Sulfur saturation limits in silicate melts and their implications for core formation scenarios for terrestrial planets. *American Mineralogist* **87**, 227–237.
- Katsura, T. & Nagashima, S. (1974). Solubility of sulphur in some magmas at 1 atmosphere. *Geochimica et Cosmochimica Acta* **38**, 517–531.
- Keppler, H. (1999). Experimental evidence for the source of excess sulfur in explosive volcanic eruptions. *Science* **284**, 1652–1654.
- Kucha, H., Wouters, R. & Arken, O. (1989). Determination of sulphur and iron valence by microprobe. *Scanning Microscopy* **3**, 89–97.
- Kullerud, G. (1971). Experimental techniques in dry sulphide research. In: Ulmer, G. C. (ed.) *Research Techniques for High Pressure and High Temperature*. New York: Springer Verlag, pp. 288–315.
- Lowenstern, J. B. (1993). Evidence for a copper-bearing fluid in magma erupted at the Valley of Ten Thousand Smokes, Alaska. *Contributions to Mineralogy and Petrology* **114**, 409–421.
- Luhr, J. F. (1990). Experimental phase relations of water- and sulphur-saturated arc magmas and the 1982 eruptions of El Chichón Volcano. *Journal of Petrology* **31**, 1071–1114.
- Luhr, J. F., Carmichael, I. S. E. & Varekamp, J. C. (1984). The 1982 eruptions of El Chichón Volcano, Chiapas, Mexico: mineralogy and petrology of the anhydrite-bearing pumices. *Journal of Volcanology and Geothermal Research* **23**, 69–108.
- Mandeville, C. W., Carey, S. & Sigurdsson, H. (1996). Magma mixing, fractional crystallisation and volatile degassing during the 1883 eruption of Krakatau volcano, Indonesia. *Journal of Volcanology and Geothermal Research* **74**, 243–274.
- Mandeville, C. W., Sasaki, A., Saito, G., Faure, K., King, R. & Hauri, E. (1998). Open-system degassing of sulphur from Krakatau 1883 magma. *Earth and Planetary Science Letters* **160**, 709–722.
- Martel, C., Pichavant, M., Bourdier, J. L., Traineau, H., Holtz, F. & Scaillet, B. (1998). Magma storage conditions and control of eruption regime in silicic volcanoes: experimental evidence from Mt. Pelée. *Earth and Planetary Science Letters* **156**, 89–99.
- Mavrogenes, J. A. & O'Neill, H. S. C. (1999). The relative effects of pressure, temperature and oxygen fugacity on the solubility of sulfide in mafic magmas. *Geochimica et Cosmochimica Acta* **63**, 1173–1180.
- Mysen, B. & Popp, R. K. (1980). Solubility of sulfur in CaMgSi₂O₆ and NaAlSi₃O₈ melts at high pressure and temperature with controlled *f*O₂ and *f*S₂. *American Journal of Science* **280**, 78–92.
- Nagashima, S. & Katsura, T. (1973). The solubility of sulfur in Na₂O–SiO₂ melts under various oxygen partial pressures at 1200, 1250 and 1300°C. *Bulletin of Chemistry of the Japanese Society* **46**, 3099–3103.
- Ohmoto, H. & Kerrick, D. M. (1977). Devolatilization equilibria in graphitic systems. *American Journal of Science* **277**, 1013–1044.
- O'Neill, H. S. C. & Mavrogenes, J. A. (2002). The sulfide capacity and the sulfur content at sulfide saturation of silicate melts at 1400°C and 1 bar. *Journal of Petrology* **43**, 1049–1087.
- Pichavant, M. (1987). Effects of B and H₂O on liquidus phase relations in the haplogranite system at 1 kbar. *American Mineralogist* **72**, 1056–1070.
- Pitzer, K. S. & Sterner, S. M. (1994). Equation of state valid continuously from zero to extreme pressures for H₂O and CO₂. *Journal of Chemical Physics* **101**, 3111–3116.
- Pownceby, M. I. & O'Neill, H. S. C. (1994). Thermodynamic data from redox reactions at high temperatures. III: Activity–composition relations in Ni–Pd alloys from EMF measurements at 850–1250 K, and calibration of the NiO+Ni–Pd assemblages as a redox sensor. *Contributions to Mineralogy and Petrology* **116**, 327–339.
- Richardson, F. D. & Fincham, C. J. B. (1954). Sulphur in silicate and aluminate slags. *Journal of the Iron & Steel Institute* **178**, 4–15.
- Richardson, F. D. & Withers, G. (1950). Thermodynamic aspects of the movement of sulphur between gas and slag in the basic open-hearth process. *Journal of the Iron & Steel Institute* **165**, 66–71.
- Ricke, W. (1960). Ein Beitrag zur Geochemie des Schwefels. *Geochimica et Cosmochimica Acta* **21**, 35–80.
- Robie, R. A., Hemingway, B. S. & Fisher, J. R. (1979). Thermodynamic properties of minerals and related substances at 298.15 K and 1 bar (10⁵ Pa) pressure and at higher temperatures. *US Geological Survey Bulletin* 1452.
- Roux, J. & Volfinger, M. (1996). Mesures précises à l'aide d'un détecteur courbe. *Journal de Physique IV* **6**, C4-127–C4-134.

- Rutherford, M. J., Sigurdsson, H., Carey, S. & Davis, A. (1985). The May 18, 1980, eruption of Mount St. Helens 1. Melt composition and experimental phase equilibria. *Journal of Geophysical Research* **90**, 2929–2947.
- St Pierre, G. R. & Chipman, J. (1956). Sulfur equilibria between gases and slags containing FeO. *Transactions of the Metallurgical Society of AIME* **206**, 1474–1483.
- Scaillet, B. & Evans, B. W. (1999). The June 15, 1991 eruption of Mount Pinatubo. I. Phase equilibria and pre-eruption P – T – $f\text{O}_2$ – $f\text{H}_2\text{O}$ conditions of the dacite magma. *Journal of Petrology* **40**, 381–411.
- Scaillet, B., Pichavant, M., Roux, J., Humbert, G. & Lefevre, A. (1992). Improvements of the Shaw membrane technique for measurement and control of $f\text{H}_2$ at high temperatures and pressures. *American Mineralogist* **77**, 647–655.
- Scaillet, B., Clemente, B., Evans, B. W. & Pichavant, M. (1998). Redox control of sulfur degassing in silicic magmas. *Journal of Geophysical Research* **103**, 23937–23949.
- Schneider, A. (1970). The sulphur isotopic composition of basaltic rocks. *Contributions to Mineralogy and Petrology* **25**, 95–124.
- Shaw, H. R. (1963). Hydrogen–water vapour mixtures: control of hydrothermal experiments by hydrogen osmosis. *Science* **139**, 1220–1222.
- Shi, P. & Saxena, S. K. (1992). Thermodynamic modeling of the C–O–H–S fluid system. *American Mineralogist* **77**, 1220–1222.
- Shima, H. & Naldrett, A. J. (1975). Solubility of sulphur in an ultramafic melt and relevance of the system Fe–S–O. *Economic Geology* **70**, 960–967.
- Spera, F. J. & Bergman, S. C. (1980). Carbon dioxide in igneous petrogenesis: I aspects of the dissolution of CO_2 in silicate liquids. *Contributions to Mineralogy and Petrology* **74**, 55–66.
- Stolper, E. M. (1982). Water in silicate glasses: an infrared spectroscopic study. *Contributions to Mineralogy and Petrology* **81**, 1–17.
- Stolper, E. M., Fine, G. J., Johnson, T. & Newman, S. (1987). The solubility of carbon dioxide in albitic melts. *American Mineralogist* **72**, 1071–1085.
- Taylor, J. R., Wall, V. J. & Pownceby, M. I. (1992). The calibration and application of accurate redox sensors. *American Mineralogist* **77**, 284–295.
- Toulmin, P. I. & Barton, P. B. J. (1964). A thermodynamic study of pyrite and pyrrhotite. *Geochimica et Cosmochimica Acta* **28**, 641–671.
- Turkdogan, E. T. & Pearce, M. L. (1963). Kinetics of sulphur reaction in oxide melt–gas system. *Transactions of the Metallurgical Society of AIME* **227**, 940–949.
- Wendlandt, R. F. (1982). Sulfide saturation of basalt and andesite melts at high pressures and temperatures. *American Mineralogist* **67**, 877–885.
- Westrich, H. R., Eichelberger, J. C. & Hervig, R. L. (1991). Degassing of the 1912 Katmai magmas. *Geophysical Research Letters* **18**, 1561–1564.
- Whitney, J. A. (1984). Fugacities of sulphurous gases in pyrrhotite-bearing silicic magmas. *American Mineralogist* **69**, 69–78.
- Winther, K. T., Watson, E. B. & Korenowski, G. M. (1998). Magmatic sulphur compounds and sulphur diffusion in albite melt at 1 GPa and 1300–1500°C. *American Mineralogist* **83**, 1141–1151.
- Yund, R. A. & Hall, H. T. (1969). Hexagonal and monoclinic pyrrhotites. *Economic Geology* **64**, 420–423.
- Zhang, Y. (1999). H_2O in rhyolitic glasses and melts: measurement, speciation, solubility, and diffusion. *Reviews of Geophysics* **37**, 493–516.