

## Nomenclature of amphiboles: Additions and revisions to the International Mineralogical Association's amphibole nomenclature

BERNARD E. LEAKE<sup>1,†</sup> (CHAIRMAN), ALAN R. WOOLLEY<sup>2</sup> (SECRETARY), WILLIAM D. BIRCH,<sup>\*,3</sup>  
ERNST A.J. BURKE,<sup>\*,4</sup> GIOVANNI FERRARIS,<sup>\*,5</sup> JOEL D. GRICE,<sup>6</sup> FRANK C. HAWTHORNE,<sup>7</sup>  
HANAN J. KISCH,<sup>8</sup> VLADIMIR G. KRIVOVICHEV,<sup>9</sup> JOHN C. SCHUMACHER,<sup>10</sup>  
NICHOLAS C.N. STEPHENSON,<sup>11</sup> AND ERIC J.W. WHITTAKER<sup>12</sup>

<sup>1</sup>School of Earth, Ocean and Planetary Sciences, Cardiff University, P.O. Box 914, Cardiff CF10 3YE, U.K.

<sup>2</sup>Department of Mineralogy, Natural History Museum, Cromwell Road, London SW7 5BD, U.K.

<sup>3</sup>Department of Mineralogy and Petrology, Museum Victoria, 328 Swanston Street, Melbourne, Victoria 3000, Australia

<sup>4</sup>Faculty of Earth and Life Sciences, Vrije Universiteit, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands

<sup>5</sup>Department of Scienze Mineralogiche e Petrologiche, Via Valperga Caluso 35, I-10125 Torino, Italy

<sup>6</sup>Mineral Sciences Division, Canadian Museum of Nature, P.O. Box 3443, Station D, Ottawa, Ontario K1P 6P4, Canada

<sup>7</sup>Department of Geological Sciences, University of Manitoba, Winnipeg, Manitoba R3T 2N2, Canada

<sup>8</sup>Department of Geology and Mineralogy, Ben Gurion University of the Negev, Beer Sheva 84105, P.O. Box 653, Israel

<sup>9</sup>Faculty of Geology, St. Petersburg University, University Em. 7/9, 199034, St. Petersburg, Russia

<sup>10</sup>Department of Earth Sciences, Wills Memorial Building, University of Bristol, Bristol BS8 1RJ, U.K.

<sup>11</sup>Earth Sciences, University of New England, Armidale, NSW 2351, Australia

<sup>12</sup>60 Exeter Road, Kidlington, Oxford OX5 2DZ, U.K.

### ABSTRACT

The introduction of a fifth amphibole group, the Na-Ca-Mg-Fe-Mn-Li group, defined by  $0.50 < {}^B(\text{Mg}, \text{Fe}^{2+}, \text{Mn}^{2+}, \text{Li}) < 1.50$  and  $0.50 \leq {}^B(\text{Ca}, \text{Na}) \leq 1.50$  apfu (atoms per formula unit), with members whittakerite and ottoliniite, has been required by recent discoveries of  ${}^B(\text{LiNa})$  amphiboles. This, and other new discoveries, such as sodicpedrizite (which is herein slightly, but significantly changed from the original idealized formula), necessitate amendments to the IMA 1997 definitions of the Mg-Fe-Mn-Li, calcic, sodic-calcic and sodic groups. The discovery of obertiite and the finding of an incompatibility in the IMA 1997 subdivision of the sodic group, requires further amendments within the sodic group. All these changes, which have IMA approval, are summarized.

### INTRODUCTION

After 13 years of conferring, the Subcommittee on Amphiboles of the International Mineralogical Association's Commission on New Minerals and Mineral Names (IMA CNMMN) first published a standard internationally agreed amphibole nomenclature in 1978 (IMA78; Leake 1978). After 9 years of further work, the current scheme was agreed and published in 1997 (IMA97; Leake et al. 1997). This retained the same four major amphibole groups and classifying parameters as in the 1978 scheme. Since 1997, several new amphibole root names have been approved by IMA, including various Li-rich amphiboles, some of which (e.g., ferriwhittakerite) have made it necessary to institute a new (fifth) amphibole group and slightly revise the definitions of the previous four groups as formulated in IMA97. In addition, application of IMA97 to several thousand analyzed natural amphiboles has shown the need to slightly revise the parameters used to subdivide the sodic amphiboles (Mogessie et al. 2001). This account summarizes the IMA-approved additions and revisions to IMA97 but does not repeat the main unchanged parts of IMA97, so this paper needs to be read in conjunction with IMA97.

† Email: LeakeB@cardiff.ac.uk

\* Indicates a non-voting official of the CNMMN

### GENERAL CLASSIFICATION OF THE AMPHIBOLES

The standard amphibole formula  $A_{0-1} B_2 C_5 T_8 O_{22}(\text{OH})_2$  remains unchanged, but the order of allocation of the cations has become even more critical than previously. In addition, some of the cations not considered in IMA97 have since been found to occur occasionally in amphiboles and so a new list follows; the full explanation of procedure is given in IMA97:

- (1) Sum T to 8.00 using Si, then Al, then Ti.
- (2) Sum C to 5.00 using any excess Al and Ti from (1) and then successively  $\text{Fe}^{3+}$ , V, Cr,  $\text{Mn}^{3+}$ , Zr, Mg, Zn, Ni, Co,  $\text{Fe}^{2+}$ ,  $\text{Mn}^{2+}$ , Li
- (3) Sum B to 2.00 using first any excess above 5.00 from C, in the reverse order of (2) starting with Li and then  $\text{Mn}^{2+}$  etc, and then follow with Ca, Sr, Ba and Na.
- (4) Excess above 2.00 in B is assigned to A in the reverse order of (3), starting with Na and then finally all the K is allocated to A.

Until recently, Li was only rarely determined in amphiboles and all Li-rich varieties known belonged either to the orthorhombic holmquistite or the monoclinic clinoholmquistite series [ $\square$ Li<sub>2</sub>(Mg<sub>3</sub>Al<sub>2</sub>)Si<sub>8</sub>O<sub>22</sub>(OH)<sub>2</sub>], being part of the Mg-Fe-Mn-Li group, or they were in the sodic amphibole group, being members of the leakeite [ $\text{Na Na}_2(\text{LiMg}_2\text{Fe}^{3+})\text{Si}_8\text{O}_{22}(\text{OH})_2$ ] or kornite [(NaK)Na<sub>2</sub>(LiMg<sub>2</sub>Mn<sup>3+</sup>)Si<sub>8</sub>O<sub>22</sub>(OH)<sub>2</sub>] species. Other,

less Li-rich Li-bearing amphiboles ( $\text{Li} > 0.25$  apfu in the standard formula) were denoted by the optional modifier “lithian”, preceding a species name which was determined by the rules formulating and subdividing the four amphibole groups. With the introduction of ion probes, the determination of Li, which electron microprobes cannot do, is now possible. Consequently, Li-rich amphiboles of a new type have been discovered which the IMA97 classification cannot accommodate satisfactorily as they fall between the existing groups.

The 1997 scheme divided the amphiboles into the following four groups, depending on the occupancy of the B sites as calculated arithmetically, i.e., there is no consideration of where ions really are because, for most amphibole analyses this is unknown, as structural studies are rare, although the arithmetic places ions in their most likely sites.

(1) The Mg-Fe-Mn-Li group defined as  ${}^B(\text{Ca} + \text{Na}) < 1.00$  and  ${}^B(\text{Mg}, \text{Fe}^{2+}, \text{Mn}^{2+}, \text{Li}) \geq 1.00$  apfu.

(2) The calcic group with  ${}^B(\text{Ca} + \text{Na}) \geq 1.00$  and  ${}^B\text{Na} < 0.50$  apfu.

(3) The sodic-calcic group with  ${}^B(\text{Ca} + \text{Na}) \geq 1.00$ , and  $0.50 \leq {}^B\text{Na} < 1.50$  apfu.

(4) The sodic group with  ${}^B\text{Na} \geq 1.50$  apfu.

At that time, nearly all known members of Group 1 had  ${}^B(\text{Mg}, \text{Fe}^{2+}, \text{Mn}^{2+}, \text{Li}) > 1.50$  apfu and the calcic and sodic-calcic members had  $(\text{Ca} + \text{Na}) > 1.50$  apfu. This led to the implicit assumption that the B sites of Group 1 contain only a small minority of Ca + Na ions and that there is only a small minority of (Mg + Fe + Mn + Li) ions at the B sites of the remaining groups.

The discovery of more or less continuous ranges of compositions, involving Li-rich amphiboles, between the Mg-Fe-Mn-Li group on the one hand and the calcic, sodic-calcic and sodic groups on the other hand, makes nonsense of the above limits. Thus an amphibole with  ${}^B(\text{Na}_{1.01}\text{Li}_{0.99})$  falls in the sodic-calcic group, although it contains no Ca, and one with  ${}^B(\text{Ca}_{0.52}\text{Na}_{0.49}\text{Li}_{0.99})$  is classified into the calcic group although Ca is not its dominant B cation. If the unchanged IMA97 rules were to be applied to the recently discovered ferriwhittakerite, ideally  $\text{Na}(\text{NaLi})(\text{LiMg}_2\text{Fe}_2^{3+})\text{Si}_8\text{O}_{22}(\text{OH})_2$  (Oberti et al. 2004), it would sit exactly on the boundary between the Mg-Fe-Mn-Li group and the sodic-calcic group, whereby an analysis with  $\text{Li} > 1.00$  apfu falls into the first group, but if  $\text{Li} < 1.00$  apfu, it belongs in the second. This sort of naming was explicitly avoided in IMA97. Consequently, the IMA97 definitions need to be modified to take into account the complete solid-solution between Li and Na at B sites (Oberti et al. 2003a). New definitions of the amphibole groups follow.

**Group 1.** Where the sum of the L-type ions  ${}^B(\text{Mg}, \text{Fe}, \text{Mn}, \text{Li}) \geq 1.50$  apfu then the amphibole is a member of the magnesium-iron-manganese-lithium group. (L-type ions are divalent Mg, Fe, Mn, Zn, Ni, Co etc and Li, as described in IMA97).

**Group 2.** Where  ${}^B(\text{Mg}, \text{Fe}^{2+}, \text{Mn}^{2+}, \text{Li}) \leq 0.50$ ,  ${}^B(\text{Ca}, \text{Na}) \geq 1.00$  and  ${}^B\text{Na} < 0.50$  apfu, then the amphibole is a member of the calcic group.

**Group 3.** Where  ${}^B(\text{Mg}, \text{Fe}^{2+}, \text{Mn}^{2+}, \text{Li}) \leq 0.50$ ,  ${}^B(\text{Ca}, \text{Na}) \geq 1.00$ , and  $0.50 \leq {}^B\text{Na} < 1.50$  apfu, then the amphibole is a member of the sodic-calcic group.

**Group 4.** Where  ${}^B(\text{Mg}, \text{Fe}^{2+}, \text{Mn}^{2+}, \text{Li}) \leq 0.50$ , and  ${}^B\text{Na} \geq 1.50$  apfu, then the amphibole is a member of the sodic group.


**Group 5.** A new amphibole group is defined as “Where  $0.50 < {}^B(\text{Mg}, \text{Fe}^{2+}, \text{Mn}^{2+}, \text{Li}) < 1.50$  and  $0.50 \leq {}^B(\text{Ca}, \text{Na}) \leq 1.50$  apfu, then the amphibole is a member of the sodium-calcium-magnesium-iron-manganese-lithium group.” This cumbersome title is clearer than any alternative and may be referred to as “Group 5”. The definitions of the prefixes and modifiers given in IMA97 that are stated to apply to all groups apply to the new group with the addition that the prefix alumino, where  ${}^C\text{Al} > 1.00$  apfu, (note not = 1.00 apfu) also applies.

## APPLICATIONS OF THE REVISED NOMENCLATURE

In general, no change is made to the nomenclature of any composition defined as a species in IMA97, but there are regrettable changes to some more recently described species, mainly because a special adjustment was initially approved to the IMA97 use of the prefix “sodic” in Group 1 in an attempt to avoid instituting a fifth amphibole group. Later however, as still more Li-rich amphiboles were discovered, the recognition of a Group 5 became unavoidable, and the use of sodic in Group 1 reverted to its IMA97 meaning of total  $\text{Na} > 0.50$  apfu. The IMA rule forbidding validity of root names until they have been found in Nature, when combined with the short time to respond to proposals for new species, makes any coherent systematic approach to a series of closely related species, proposed over several years, difficult. Ideally one would wait a few years until the range of variation was apparent.

### Group 1: Mg-Fe-Mn-Li amphiboles

Holmquistite and clinoholmquistite,  $\square \text{Li}_2(\text{Mg}_3\text{Al}_2)\text{Si}_8\text{O}_{22}(\text{OH})_2$ , have  ${}^B\text{Li} \geq 1.00$  apfu but no significant  ${}^C\text{Li}$ . The discovery of sodic-ferripedrizite  ${}^A\text{Na}{}^B\text{Li}_2{}^C(\text{LiMg}_2\text{Fe}^{3+}_2)\text{Si}_8\text{O}_{22}(\text{OH})_2$  (Oberti et al. 2000; Caballero et al. 2002), and the equivalent



**FIGURE 1.** Classification of the Group 1 amphiboles; Mg-Fe-Mn-Li amphiboles.

sodic-ferri-ferropedrizite (Oberti et al. 2003) species in which it is critical to have  ${}^C\text{Li} \geq 0.50$  apfu, to distinguish them from holmquistite and clinoholmquistite, justify the following names:

Sodicpedrizite  $\text{Na Li}_2 (\text{LiMg}_2\text{Fe}^{3+}\text{Al}) \text{Si}_8 \text{O}_{22} (\text{OH})_2$

Sodic-ferropedrizite  $\text{Na Li}_2 (\text{LiFe}^{2+}_2\text{Fe}^{3+}\text{Al}) \text{Si}_8 \text{O}_{22} (\text{OH})_2$ .

The prefix sodic is derived from the IMA97 rules for the use of prefixes in which sodic applies in Group 1 only, and then only to samples with total Na > 0.50 apfu. Those with total Na  $\leq 0.50$  apfu are pedrizite unless K > 0.50 apfu which is potassicpedrizite. Likewise the prefix ferri for  $\text{Fe}^{3+} > 1.00$  apfu (not = 1.00), applies to all groups except the sodic amphiboles. All the prefix rules listed in IMA97 remain unchanged. It is recognized that a few samples that only marginally fell into IMA97 Group 1, may now belong to Group 5. The revised classification is shown in Figure 1. The only substantive change is to Li-rich amphiboles.

### Group 2: Calcic amphiboles

There are no changes, or additions, of names in this group. Application of the prefix parvo, as described below, necessitates over-riding the new  ${}^B(\text{Mg}, \text{Fe}^{2+}, \text{Mn}^{2+}, \text{Li}) \leq 0.50$  apfu rule.

### Group 3: Sodic-calcic amphiboles

Again there are no changes or additions in this group and the same provision applies to the use of parvo as in the calcic amphiboles.

### Group 4: Sodic amphiboles

There are no changes deriving from the introduction of Group 5 except for the use of parvo as in the previous groups. However, since IMA97 was published (Leake et al. 1997), extensive tests have been carried out on published amphibole analyses to see how robust the nomenclature is, especially with respect to compositions marginal to the four groups. Mogessie et al. (2001) calculated the names of over 500 analyzed amphiboles of all types as listed in Deer et al. (1997) and they found about four sodic amphibole analyses that could not be classified, as although they were clearly sodic amphiboles, they did not fall

into any of the sodic nomenclature diagrams. Consequently two amendments were made to avoid this. First, the rarely known oxidation state of Mn is made much less crucial than in IMA97 by grouping  $\text{Mn}^{2+} + \text{Mn}^{3+}$  together for the initial classification, although  $\text{Mn}^{3+}$  contents are still required to distinguish kornite from leakeite. Secondly, the classification parameter  $(\text{Mg} + \text{Fe}^{2+} + \text{Mn}^{2+}) < \text{or} > 2.5$  apfu is completely discarded as it conflicts with other conditions. The result of these changes gives a new sodic amphibole nomenclature diagram. (Fig. 2).

In addition, the new anhydrous sodic species obertiite  $\text{Na Na}_2 (\text{Mg}_3\text{Fe}^{3+}\text{Ti}) \text{Si}_8 \text{O}_{22} \text{O}_2$  which is characterized by  $\text{Ti} > 0.50$  and  $(\text{OH} + \text{F} + \text{Cl}) < 1.00$  apfu, has recently been described by Hawthorne et al. (2000). As this can be derived from magnesio-arfvedsonite by substitution of Ti and O for Mg and (OH), it appears in the same box as magnesio-arfvedsonite in the Group 4 nomenclature diagram. As  $\text{Ti} > 0.50$  apfu is an essential aspect of the root name, the prefix titano is not used, nor is the modifier oxygenian because  $(\text{OH} + \text{F} + \text{Cl}) < 1.00$  apfu by definition.

### Group 5: Na-Ca-Mg-Fe-Mn-Li amphiboles

(1)  ${}^B\text{Li} > 0.50$  apfu. The members of this new group that have  ${}^B\text{Li} > 0.50$  apfu derive from Oberti et al. (2004) finding ferriwhittakerite,  ${}^A\text{Na} {}^B(\text{NaLi}) {}^C(\text{LiMg}_2\text{Fe}_2^{3+}) \text{Si}_8 \text{O}_{22} (\text{OH})_2$  with  $A \geq 0.50$  apfu and ferri-ottoliniite,  ${}^A\Box {}^B(\text{NaLi}) {}^C(\text{Mg}_3\text{Fe}_3^{3+}) \text{Si}_8 \text{O}_{22} (\text{OH})_2$ , with  $A < 0.50$  apfu. Both must have  ${}^B\text{Li} > 0.50$  apfu. Usually there is also significant  ${}^C\text{Li}$ . The classification is shown in Figure 3 and the end member compositions are listed below.

Whittakerite  $\text{Na} (\text{NaLi}) (\text{LiMg}_2\text{Fe}^{3+}\text{Al}) \text{Si}_8 \text{O}_{22} (\text{OH})_2$  and “Ferrowhittakerite”  $\text{Na} (\text{NaLi}) (\text{LiFe}_2^{3+}\text{Fe}^{3+}\text{Al}) \text{Si}_8 \text{O}_{22} (\text{OH})_2$

Ottoliniite  $\Box (\text{NaLi}) (\text{Mg}_3\text{Fe}^{3+}\text{Al}) \text{Si}_8 \text{O}_{22} (\text{OH})_2$  and “Ferro-ottoliniite”  $\Box (\text{NaLi}) (\text{Fe}_3^{3+}\text{Fe}^{3+}\text{Al}) \text{Si}_8 \text{O}_{22} (\text{OH})_2$ .

All the prefixes and modifiers listed and defined in IMA97 apply in the new group, and use of the prefix alumino for  ${}^C\text{Al} > 1.00$  apfu is extended to this group. In accordance with IMA97, the modifier lithian is not used in species defined by the presence of Li.

(2)  ${}^B\text{Li} \leq 0.50$  apfu. It is well known that in the calcic and sodic-calcic groups, it is usual for the total of the cations initially





FIGURE 2. Classification of the Group 4 amphiboles; sodic amphiboles.



**FIGURE 3.** Classification of the Group 5 amphiboles; Na-Ca-Mg-Fe-Mn-Li amphiboles.

allocated to C to exceed the limit of 5.00 apfu and the excess of the larger ions overspill into B. This is particularly common where “cummingtonite solid solution” in calcic amphiboles occurs. Occasionally, the total (Mg + Fe<sup>2+</sup> + Mn<sup>2+</sup> + Li) at B slightly exceeds 0.50 apfu although <sup>B</sup>Li < 0.50 apfu. This is usually due to erroneously low (OH + F + Cl), giving too large a cation total, but there are a few reliable analyses with such excesses above <sup>B</sup>0.50 and with <sup>B</sup>Li ≤ 0.50 apfu. Such samples only slightly exceed <sup>B</sup>0.50 apfu because the presently known miscibility limit of such amphiboles is only slightly greater than <sup>B</sup>0.50 apfu. In order to avoid a rash of new root names for such compositions of small cations in B (and small or zero Li), the prefix “parvo” from the Latin for “small” should be prefixed to the root name derived for a species in the calcic, sodic-calcic or sodic groups, as if Group 5 did not exist, i.e., over-riding in this instance only, the requirement in Groups 2, 3 and 4 for <sup>B</sup>(Mg,Fe<sup>2+</sup>,Mn<sup>2+</sup>,Li) ≤ 0.50 apfu. This preserves familiar IMA97 names, makes no distinction as to what the dominant L-type ion is in B, and avoids making new names as a consequence of creating Group 5 to deal with certain Li-rich amphiboles. Thus such analyses are named exactly as they were before Group 5 existed, but with the added prefix “parvo”. An unusually large <sup>B</sup>(Mg,Fe,Mn,Li) value of <sup>B</sup>0.74 apfu occurs in the following rare composition, which was in Group 3, but which now falls into Group 5: <sup>A</sup>(K<sub>0.03</sub>Na<sub>0.30</sub>)<sup>B</sup>(Na<sub>0.87</sub>Ca<sub>0.39</sub>Mn<sub>0.68</sub>Li<sub>0.06</sub>)<sup>C</sup>(Mn<sub>0.27</sub>Mg<sub>4.01</sub>Fe<sub>0.72</sub>)<sup>D</sup>Si<sub>7.88</sub>Al<sub>0.12</sub>O<sub>22</sub>F<sub>0.40</sub>(OH)<sub>1.60</sub> which was reported by Oberti and Ghose (1993). According to IMA97, this is a winchite, or with modifiers, a fluorian manganian winchite. It now becomes parvowinchite or fluorian manganian parvowinchite. As the authors interpret it as “on the verge of exsolution” it may be near to the extreme value of high <sup>B</sup>(Mg,Fe<sup>2+</sup>,Mn<sup>2+</sup>) in Li-poor Group 5. Similarly, there are rare <sup>B</sup>Li ≤ 0.50 apfu amphiboles which were previously in Group 2, but now fall into Group 5. Thus a most extraordinary calcic amphibole, in its high <sup>B</sup>(Mn), is <sup>A</sup>(K<sub>0.02</sub>Na<sub>0.65</sub>)<sup>B</sup>(Ca<sub>1.17</sub>Mn<sub>0.83</sub>)<sup>C</sup>(Mn<sub>0.27</sub>Fe<sub>0.09</sub>Mg<sub>4.46</sub>Ti<sub>0.01</sub>Al<sub>0.16</sub>)<sup>D</sup>Si<sub>7.18</sub>Al<sub>0.82</sub>O<sub>22</sub>(OH<sub>1.91</sub>F<sub>0.09</sub>) (Skogby and Rossman 1991). If this is not a mixture of phases, then it was per IMA97, a manganio-edenite and it now becomes parvo-manganio-edenite. More usually, any excess of <sup>B</sup>(Mg,Fe<sup>2+</sup>,Mn<sup>2+</sup>) above 0.50 apfu in Li-poor (Li < 0.50) calcic amphiboles is trivial.

Equally, to avoid new root names proliferating for rare amphiboles with compositions of <sup>B</sup>Li ≤ 0.50 apfu which were in Group

1, but now fall into Group 5, because of significant <sup>B</sup>(Ca,Na), these should be named as previously with Group 1 root names and prefixes but prefixed “magno” to indicate the entry of large cations at B position in Group 5.

It should be noted that in the first published use of sodic-ferripedrizite, (Oberti et al. 2000), one of the analyzed crystals is sodic-ferripedrizite as above, whereas the second is now ferriwhittakerite, because the ideal formula of pedrizite was later changed to contain <sup>B</sup>Li<sub>2</sub>, while <sup>B</sup>(NaLi) became whittakerite or ottoliniite. The second published use of ferripedrizite (Caballero et al 2000) had <sup>B</sup>Li<sub>2</sub> and is only changed by now being prefixed with sodic as above.

**Erratum.** In IMA97, Table 2, the modifier ferrian, which applies in all but the sodic group, should have been stated as  $0.75 < \text{Fe}^{3+} \leq 1.00$  and not 0.99 apfu.

### ACKNOWLEDGMENTS

We particularly thank R. Oberti for very substantial help, patience, and cooperation.

### REFERENCES CITED

- Caballero, J.M., Oberti, R., and Ottolini, L. (2002) Ferripedrizite, a new monoclinic <sup>B</sup>Li amphibole end-member from the Eastern Pedriza Massif, Sierra de Guadarrama, Spain, and a restatement of the nomenclature of the Mg-Fe-Mn-Li amphiboles. *American Mineralogist*, 87, 976–982.
- Deer, W.A., Howie, R.A., and Zussman, J. (1997) *Rock-forming Minerals*, 2B, Double-Chain Silicates, (2nd ed.), 764 p. Geological Society, London.
- Hawthorne, F.C., Cooper, M.A., Grice, J.D., and Ottolini, L. (2000) A new anhydrous amphibole from the Eifel region, Germany: description and crystal structure of obertiite, NaNa<sub>3</sub>(Mg<sub>3</sub>Fe<sup>3+</sup>Ti<sup>4+</sup>)Si<sub>8</sub>O<sub>22</sub>O<sub>2</sub>. *American Mineralogist*, 85, 236–241.
- Leake, B.E. (1978) Nomenclature of amphiboles. *American Mineralogist*, 63, 1023–1052.
- Leake, B.E., Woolley, A.R., Arps, C.E.S., Birch, W.D., Gilbert, M.C., Grice, J.D., Hawthorne, F.C., Kato, A., Kisch, H.J., Krivovichev, V.G., Linthout, K., Laird, J., Mandarino, J.A., Maresch, W.V., Nickel, E.H., Rock, N.M.S., Schumacher, J.C., Smith, D.C., Stephenson, N.C.N., Ungaretti, L., Whittaker, E.J.W., and Youzhi, G. (1997) Nomenclature of amphiboles: Report of the Subcommittee on Amphiboles of the International Mineralogical Association, Commission on New Minerals and Mineral Names. *American Mineralogist*, 82, 1019–1037.
- Mogessie, A., Ettinger, K., Leake, B.E., and Tessadri, R. (2001) AMPH-IMA97: a hypercard program to determine the name of an amphibole from electron microprobe and wet analyses. *Computers and Geosciences*, 27, 1169–1178.
- Oberti, R. and Ghose, S. (1993) Crystal-chemistry of a complex Mn-bearing alkali amphibole (“tirodite”) on the verge of exsolution. *European Journal of Mineralogy*, 5, 1153–1160.
- Oberti, R., Caballero, J.M., Ottolini, L., López-Andrés, S., and Herreros, V. (2000) Sodicferripedrizite, a new monoclinic amphibole bridging the magnesium-iron-manganese-lithium and the sodium-calcium groups. *American Mineralogist*, 85, 578–585.
- Oberti, R., Cámara, F., Ottolini, L., and Caballero, J.M. (2003a) Lithium in amphiboles: detection, quantification and incorporation mechanisms in the compositional space bridging sodic and <sup>B</sup>Li amphibole. *European Journal of Mineralogy*, 15, 309–319.
- Oberti, R., Cámara, F., Caballero, J.M., and Ottolini, L. (2003b) Sodic-ferri-ferropedrizite and ferri-clinoferroholmquistite. Mineral data and ordering of A-cations in Li-rich amphiboles. *Canadian Mineralogist*, 41, 1345–1354.
- Oberti, R., Cámara, F., and Caballero, J.M. (2004) Ferri-ottoliniite and ferriwhittakerite, two new end-members in the new Group 5 for amphiboles. *American Mineralogist*, 89, 888–893.
- Skogby, H. and Rossman, G.R. (1991) The intensity of amphibole OH bands in the infrared absorption spectrum. *Physics and Chemistry of Minerals*, 18, 64–68.

### APPENDIX

Information concerning the etymology, the type locality and the unit-cell parameters of amphibole root name end-members described since IMA97.

**Obertiite**

Named for R. Oberti of Pavia, Italy.

Type locality: Bellerberg, Laccher See district, Eifel, Germany.

X-ray data: a 9.776 Å, b 17.919 Å, c 5.292 Å.  $\beta$  104.05°.

Reference: Hawthorne et al. (2000).

**Ottoliniite**

Named for Luisa Ottolini of Pavia, Italy.

Type locality: Fuente Grande, Eastern Pedriza Massif, Sierra de Guadarrama, Central System, Spain.

X-ray data: a 9.535 Å, b 17.876 Å, c 5.234 Å.  $\beta$  102.54°.

Reference: Oberti et al. (2004).

**Pedrizite**

Named for locality.

Type locality: Fuente Grande, Eastern Pedriza Massif, Sierra de Guadarrama, Central System, Spain.

X-ray data: a 9.501 Å, b 17.866 Å, c 5.292 Å.  $\beta$  102.17°.

Reference: Caballero et al. (2002); name first used by Oberti et al. (2000). Ferropedrizite is described in Oberti et al. (2003b).

**Whittakerite**

Named for Eric J. W. Whittaker of Oxford, England.

Type locality: Fuente Grande, Eastern Pedriza Massif, Sierra de Guadarrama, Central System, Spain.

X-ray data: a 9.712 Å, b 17.851 Å, c 5.297 Å.  $\beta$  103.63°. Reference: Oberti et al. (2004), first used in Oberti et al. (2003a).