

Gamma rays from cosmic radioactivities

Roland DIEHL^{1*}, Dieter H. HARTMANN², and Nikos PRANTZOS³

¹Max-Planck-Institut für extraterrestrische Physik, D-85748 Garching, Germany

²Department of Physics and Astronomy, Clemson University, Clemson, South Carolina 29634, USA

³Institut d'Astrophysique, F-75014 Paris, France

*Corresponding author. E-mail: rod@mpe.mpg.de

(Received 14 October 2006; revision accepted 19 March 2007)

Abstract—Gamma rays from radioactive byproducts of cosmic nucleosynthesis are direct messengers from nuclear processes taking place in various cosmic sites, and can be measured with telescopes operated in space. Due to low detector sensitivity, up until now, only a handful of sources have been detected in that electromagnetic window. Cobalt lines from SN1987A and ⁴⁴Ti lines from the Cassiopeia A (Cas A) supernova remnant offer unique constraints on the properties of the innermost regions of core collapse supernovae. Diffuse gamma-ray lines from the decay of radioactive ²⁶Al and the annihilation of positrons are bright enough for mapping the Milky Way in the MeV regime, and are both measured by recent spaceborne spectrometers with unprecedented precision. This constrains the sources of Al production and the state of interstellar gas in the vicinity of these sites: the total mass of ²⁶Al produced by stellar sources throughout the Galaxy is estimated to be ~3 M_⊙ per Myr, and the interstellar medium near those sources appears to be characterized by velocities of ~100 km s^{−1}. Positron annihilation must occur in a modestly ionized, warm phase of the interstellar medium, but at present the major positron production site(s) remain unknown. The spatial distribution of the annihilation gamma-ray emission constrains positron production sites and positron propagation in the Galaxy. ⁶⁰Fe radioactivity has been clearly detected recently; the flux ratio relative to ²⁶Al of about 15% is on the lower side of predictions from massive star and supernova nucleosynthesis models. Those views at nuclear and astrophysical processes in and around cosmic sources by space-based gamma-ray telescopes offer invaluable information on cosmic nucleosynthesis.

INTRODUCTION

Branches of astronomy are often defined by their observational band or technique, such as “optical,” “radio,” or “high-energy” astronomy. Ernst Zinner has been one of the pioneers of a completely different and alternative “astronomy” rooted in terrestrial laboratories: high-precision mass spectrometry of the isotopic composition of presolar grains (included in meteorites as direct and material probes of cosmic nucleosynthesis), which is now established as an astronomy discipline in its own right (Anders and Zinner 1993; Zinner et al. 1987; Clayton and Nittler 2004; Zinner 1998). Much of the information that links such isotopic measurements to cosmic nucleosynthesis processes is based on radioactive decays of unstable isotopes. Characteristic daughter-product isotopes generate a unique isotope abundance signature, which is distinct from common isotopic varieties in meteoritic material. The astrophysical challenge that connects Ernst Zinner’s investigations of meteorites and

presolar grains to our work, which is based on radioactivity gamma rays (Fig. 1), concerns the physics underlying the formation of fresh nuclei in the universe. The relevant processes occur inside stars undergoing either hydrostatic or explosive nuclear burning, but also in interstellar space via high-energy particle interactions (Wallerstein et al. 1997). Theories for cosmic nucleosynthesis continue to be developed following the pioneering works of Cameron and Burbidge, Burbidge, Fowler, and Hoyle in 1957 (Cameron 1970; Burbidge et al. 1957; Arnett 1969; Truran and Cameron 1971; Clayton 1968). Observations of radioactive isotopes through gamma rays ensued from pioneering gamma-ray experiments following the Apollo program of the 1970s and balloon experiments (Johnson and Haymes 1973). Parallel to this experimental development, the emergence of isotopic analyses of stardust in the laboratory (after Ca-Al-rich inclusions of meteorites had been recognized as carriers of presolar signatures [Zinner et al. 1987]) has been consolidated by Ernst Zinner and his colleagues (Zinner 1998). A direct




Fig. 1. Radioactive decay of unstable isotopes leads to gamma-ray emission, which can be measured from telescopes aboard satellites above the atmosphere of the Earth. Here the case of ^{26}Al is shown, decaying with a characteristic lifetime of 1.04 Myr into an excited state of ^{26}Mg , which de-excites into the Mg ground state by gamma-ray emission.

connection between theories of nucleosynthesis and observation of its products is possible with the detection of trace isotopes, which is feasible with current technology both in space and in terrestrial laboratories. In comparison, the vast amount of data on stellar spectroscopy (Cayrel and Cayrel de Strobel 1966; Cayrel de Strobel et al. 2001; Gratton et al. 2000; Edvardsson et al. 1993) offers complementary, but less direct, information: indeed, theories for element mixing and radiative transfer in stellar atmospheres must be employed to convert observed spectral features to information on nucleosynthesis. As part of the common theme of astronomy with radioactivities, we present here the experimental techniques and astrophysical achievements in astronomy obtained with gamma-ray telescopes. We also point to a series of workshops on this topic, where over the past ten years we have enjoyed frequent discussions with Ernst Zinner and his colleagues on those common themes (Diehl and Hartmann 2000; Diehl et al. 2002; Diehl et al. 2004; Hartmann et al. 2006).

GAMMA-RAY TELESCOPES

Gamma rays typically penetrate matter of substantial density (on the order of a few grams cm^{-2})—this allows observers to probe far-away regions of the Galaxy even if they are obscured by molecular clouds, or the interiors of supernova or nova explosions. The other downside of this penetrating power is the difficulty to detect and measure

cosmic gamma rays, and even more to derive the imaging information needed for many astronomical purposes. Gamma-ray telescopes (Fig. 2) consist of massive detector elements with typical thickness of several centimeters (much larger than the micrometers-thick CCD cameras used for optical imaging). Moreover, many of those detectors have to be operated simultaneously with electronic systems to capture imaging information and to reduce instrumental backgrounds (for more details see Diehl et al. 2006a). Instrumental backgrounds are large in space environments due to the irradiation of telescopes and satellites from cosmic ray particles, which leads to nuclear excitations of atomic nuclei in those materials producing gamma rays locally, just as the remote cosmic sites of interest do. Massive shields of gamma-ray cameras through particle detectors are commonly used to provide a veto signal when cosmic-ray particles hit the instrument. In comparison with other wavelength regions, these experimental difficulties have slowed progress of astrophysical studies in the gamma-ray band. But the unique information from nuclear processes in cosmic sites makes these efforts worthwhile (Diehl et al. 2006a; Prantzos and Diehl 1996; Diehl and Timmes 1998).

After pioneering work with detectors flown on stratospheric balloons, rockets and satellites in the 1960s and 1970s, NASA's Compton Gamma Ray Observatory mission of 1991–2000 performed the first sky survey of bright gamma-ray line emission, with a limiting sensitivity of about 10^{-5} $\text{ph cm}^{-2} \text{s}^{-1}$ and spatial resolution of a few degrees (Gehrels et al.

Table 1. The isotopes that lead to astronomically relevant gamma-ray lines from their radioactive decay. Mean decay times are given for the slowest decay in the chain. Isotopes which have been detected in gamma-rays show their gamma-ray energies marked in green. The underlying nucleosynthesis processes are indicated, with H = hydrogen burning, NSE = nuclear statistical equilibrium, meaning nucleosynthesis in equilibrium among all strong nuclear reactions, α -NSE = α -rich freeze-out from NSE in core-collapse supernovae, Ne = explosive neon burning, ν = neutrino process that decomposes nuclei generating excess free nucleons by inelastic scattering.

Isotope	Mean decay time	Decay chain	γ -ray energy (keV)	Detected source	Source type	Nuclear process
^7Be	77 d	$^7\text{Be} \rightarrow ^7\text{Li}^*$	478	None	Novae	H (explosive)
^{56}Ni	111 d	$^{56}\text{Ni} \rightarrow ^{56}\text{Co}^* \rightarrow ^{56}\text{Fe}^* + e^*$	158, 812, 847, 1238	SN1987A, SN1991T	Supernovae	NSE
^{57}Ni	390 d	$^{57}\text{Co} \rightarrow ^{57}\text{Fe}^*$	122	SN1987A	Supernovae	NSE
^{22}Na	3.8 yr	$^{22}\text{Na} \rightarrow ^{22}\text{Ne}^* + e^*$	1275	None	Novae	H (explosive)
^{44}Ti	85 yr	$^{44}\text{Ti} \rightarrow ^{44}\text{Sc}^* \rightarrow ^{44}\text{Ca}^* + e^*$	78, 68, 1157	SNR Cas A	Supernovae	α -NSE
^{26}Al	$10.4 \cdot 10^6$ yr	$^{26}\text{Al} \rightarrow ^{26}\text{Mg}^* + e^*$	1809	Galactic plane regions	Stars, novae, supernovae	H, Ne (hydrost., explosive), ν
^{60}Fe	$2.2 \cdot 10^6$ yr	$^{60}\text{Fe} \rightarrow ^{60}\text{Co}^* \rightarrow ^{60}\text{Ni}^*$	59, 1173, 1332	Galaxy	Supernovae, stars	n capture
e^+	10^5 – 10^7 yr	$e^+ + e^- \rightarrow \text{Ps} \rightarrow \gamma\gamma$	511, <511	Galactic bulge	Supernovae, novae	β^+ decay




Fig. 2. Satellites such as ESA's INTEGRAL (top) carry gamma-ray telescopes above the atmosphere of the Earth. Gamma-ray telescopes consist of arrays of photon detectors, which image the sky either through a measurement of Compton scattering (left; the COMPTEL telescope consists of two planes of photon detectors operating in coincidence), or through a "coded" telescope aperture (right; the SPI imaging spectrometer (Vedrenne et al. 2003; Roques et al. 2003) is built around a 19-element camera of Ge detectors; incident gamma-rays will cast a characteristic shadow onto this camera due to the coded mask, which allows to discriminate sources against the instrumental background).

1993). From these early observations, it was clearly established that the inner Galaxy is a bright source of positron annihilation gamma rays at 511 keV (Purcell et al. 1997), the Galaxy glows in ^{26}Al radioactivity (Diehl et al. 1995), and supernova radioactivity is observable in special circumstances, such as ^{44}Ti from the 340-year-old Cassiopeia A (Cas A) supernova remnant (Iyudin et al. 1994), and Ni decay gamma rays from SN1987A (Kurfess et al. 1992) and the bright type Ia supernova called SN1991T (Morris et al. 1997). However, several puzzles remained: nova radioactivity had not been seen directly (Iyudin et al. 1995), supernova objects were found to be less luminous than expected (Georgii et al. 2002), and the tantalizing reports of nuclear excitation lines from star-forming regions such as Orion (Bloemen et al. 1994), the 2.2 MeV gamma rays (McConnell et al. 1997) possibly from a nearby white dwarf, and a second ^{44}Ti supernova remnant in the Vela region (Iyudin et al. 1998) were too uncertain for realistic astrophysical interpretations (Bloemen et al. 1999; Shönfelder et al. 2000). It should be noted that most of these gamma-ray line studies were made with modest spectral resolution and thus could not really resolve and autonomously identify the nuclear lines (see Table 1).

Recently established observatories with high-resolution solid-state detectors such as the Reuven Ramaty High-Energy Solar Spectroscopic Imager (RHESSI) (Lin et al. 2002) and the International Gamma-Ray Astrophysics Laboratory (INTEGRAL)/SPI spectrometer (Winkler et al. 2003; Vedrenne et al. 2003; Roques et al. 2003) (Fig. 2) added a new quality to this field: spectroscopy of these nuclear lines allows the identification of cosmic lines against reference lines from instrumental background and is capable of constraining the kinematics of cosmic isotopes in the gamma-ray emission source region. The astrophysical importance of such velocity information for the study of particle acceleration in solar flares has been demonstrated and will not be further addressed here (Murphy and Share 2005; Gros et al. 2004; Hurford et al. 2006). Below we discuss how this information is exploited for cosmic sources, i.e., for supernova nucleosynthesis ejecta and for the turbulent interstellar medium in the vicinity of massive stars. The special processes involved in positron annihilation gamma-ray production (Guessoum et al. 2005) shape the corresponding gamma-ray line at 511 keV. This can be exploited to constrain the ionization state and temperature of the interstellar gas in the region where positrons annihilate, which may be quite far from the locales where the positrons were generated in the first place.

Candidate sources of characteristic gamma-ray lines are supernovae and novae, but also the winds from massive stars (Diehl and Timmes 1998; Diehl et al. 2006a). In these objects, freshly produced nuclei from explosive layers near the surface of compact stars and from stellar interiors are ejected into interstellar space, where their decay can be observed directly

in gamma rays. Once ejected, long-lived radioisotopes will decay in interstellar space; once merged with interstellar gas, they provide radioactive tracers that reflect the state of circumstellar and interstellar gas around the source. The kinematics of inner-ejecta material (rather than outer envelope and swept-up matter) can be studied in young supernova remnants; more or less complete ionization of matter will lead to modified decay histories for the total of ^{44}Ti produced in the explosion (mean decay time of 85 years [Ahmad et al. 2004] by electron capture only). Longer-lived isotopes such as ^{26}Al ($\tau \sim 1.04$ Myr, see review by Prantzos and Diehl 1996) will reflect interstellar-gas kinematics through Doppler shift or broadening of decay gamma-ray line(s); this interstellar medium around massive star sources is otherwise hard to study, as it is hot and tenuous in nature. Positrons from decays of isotopes on the proton-rich side of the valley of stability will typically be produced at MeV energies; their propagation before their annihilation is more complex, being controlled by density and magnetic field morphologies near the sources (Guessoum et al. 2005; Murphy and Share 2005); high-energy emission from annihilation-in-flight would be the “smoking gun” near the source, while the 511 keV line and lower-energy continuum arise from thermalized positrons with therefore unknown origin.

GAMMA-RAY LINES FROM SUPERNOVAE

Cobalt Gamma-Ray Lines from Radioactivity Powering Supernova Light Curves

Supernovae may shine as bright as entire galaxies for weeks. The energy source providing this luminosity cannot be the explosion itself, as this would result in much more rapid fading; radioactive release of energy over those longer time scales is required and most likely can be attributed to the decay of freshly synthesized Ni and Co isotopes, with characteristic decay lifetimes of 8 and 111 days for ^{56}Ni and ^{56}Co , respectively. The amount of synthesized radioactive Ni provides one of the key parameters of supernova explosion models. Its determination is indirect, in general, involving models of energy conversion from radioactivity (gamma-rays and positrons) to optical light (Höflich 2006; Röpke et al. 2006). Gamma-ray lines from ^{56}Co and ^{57}Co decays have been detected from supernova SN1987A with the Gamma-Ray Spectrometer on the Solar Maximum Mission (Matz et al. 1988) and the Oriented Scintillation Spectrometer Experiment on the Compton Gamma Ray Observatory (Kurfess et al. 1992), respectively. These were the first and only direct confirmations that Ni decay and its products power supernova light curves. However, the accuracy of absolute flux determination was insufficient for direct calibration of the corresponding yields due to instrumental background and the embedding of those radioactivities in a massive and semi-opaque supernova envelope: SN1987A




Fig. 3. Gamma-ray flux from Cas A from radioactive decay of ^{44}Ti has been measured by several X/gamma-ray instruments. In spite of instrumental uncertainties, results consistently suggest $\sim 10^{-4} M_{\odot}$ of ^{44}Ti production and make this supernova a unique study object for core-collapse models (see text).

was the core-collapse termination of a massive star's evolution, and its explosion produced $\sim 0.075 M_{\odot}$ of ^{56}Ni and $0.003 M_{\odot}$ of ^{57}Co largely buried under $\sim 16 M_{\odot}$ of ejecta. These amounts of radioactive material have been determined from the measurement of the fading supernova light assuming full deposit and conversion of the radioactive energy, and are consistent with precision gamma ray flux results (Woosley et al. 1989).

A calibration of ^{56}Ni masses is particularly important for supernovae of type Ia because of their cosmological “standard candle” applications (Leibundgut 2000). These events are attributed to thermonuclear explosions of white dwarfs in binary systems (Hoeftlich 2006), they produce about $0.5 M_{\odot}$ of ^{56}Ni and have small overlying envelopes. Their remarkably homogeneous light curves ask for a thorough physical understanding of that kind of explosions so that the spectacular cosmological implications on the long-term expansion behavior of the universe derived from SNIa are put on more than empirical grounds (Röpke et al. 2006; Leibundgut 2000). Unfortunately, such events are rare in the nearby universe accessible to current-day gamma-ray telescopes. INTEGRAL can, in principle, detect SNIa up to 5 Mpc and, after four years of operation, is still waiting for such an opportunity (Isern et al. 2004). The results from the Compton observatory for two relatively nearby events were controversial: SN1991T at a distance of 13 Mpc was an exceptionally bright SNIa, but in order to explain COMPTEL's marginal gamma-ray line signal (Morris et al. 1997; Morris et al., unpublished data) $1.65 M_{\odot}$ of ^{56}Ni is required; this is on the high side of indirect determinations, but too uncertain for far-reaching interpretations. SN1998bu, at a distance of 11.6 Mpc, was somewhat closer; however, no gamma-ray lines were observed from the estimated $0.7\text{--}0.8 M_{\odot}$ of ^{56}Ni , which puts tight constraints on some models of SNIa (Georgii et al. 2002).

It is clear that gamma-ray telescopes will fulfill their promise of solving one of the key issues in astrophysical models of supernova explosions, namely their synthesis of

^{56}Ni , only with substantially more sensitive experiments that are able to reach SNIa from beyond the Virgo cluster and hence measure a substantial statistical sample of such events. Two experiments exemplify what is being studied for this purpose: the European Gamma-Ray Imager (Knödlseider 2006) and the U.S. Advanced Compton Telescope (Boggs et al. 2006); they await announcements of mission opportunities by space agencies.

^{44}Ti from Cassiopeia A: Probing Asymmetries of Core Collapses?

The Cas A supernova remnant has an estimated age of 340 years and provides us with an excellent opportunity to study the recent explosion of a relatively nearby massive star. Ejecta kinematics has been studied in detail through recombination lines, and the emission morphology of the remnant imaged in X-rays through optical to infrared emission suggests substantial 3-D effects and deviations from spherical symmetry (Hughes et al. 2000). A pioneering discovery of ^{44}Ti radioactivity gamma rays at 1.156 MeV was made with the Imaging Compton Telescope on NASA's Compton observatory (Iyudin et al. 1994). Several other instruments have confirmed ^{44}Ti decay in this young remnant (Fig. 3): from Beppo-Sax measurements, the 68 and 78 keV lines from ^{44}Sc de-excitation were found (Vink et al. 2001), and INTEGRAL/IBIS recently detected the same ^{44}Sc hard X-ray lines with improved significance (Renaud et al. 2006). The combined gamma-ray measurements indicate a flux of $\sim 2.5 (\pm 0.5) 10^{-5} \text{ ph cm}^{-2} \text{ s}^{-1}$ (Nittler et al. 1996; Vink 2005). With the distance ($\sim 3.4 \text{ kpc}$ [Reed et al. 1995]), a radioactive lifetime of ^{44}Ti (85.2 years [Ahmad et al. 2006]), and the ionization state for ^{44}Ti (Motizuki et al. 1999), the γ -ray flux translates into a ^{44}Ti yield of $0.8\text{--}2.5 10^{-4} M_{\odot}$ of ^{44}Ti produced in this supernova. This places Cas A's ^{44}Ti yield somewhat on the high side when compared to the predictions of spherically symmetric models (Prantzos 2004a; Diehl et al. 2006a) (Fig. 5).




Fig. 5. The SPI measurement of the ^{26}Al line from the inner Galaxy finds the line broadening from astrophysical effects (galactic rotation, ISM turbulence) to be rather small (Diehl et al. 2006b).




Fig. 4. The image obtained by the COMPTEL telescope through the 9-year Compton observatory mission in the 1808.65 keV line from ^{26}Al shows the nucleosynthesis traces from massive stars in the galaxy over the last several million years (^{26}Al decays with a radioactive lifetime of 1.04 Myr).

Increased ^{44}Ti production in Cas A due to deviations from spherical symmetry in the explosion has been suggested (e.g., Prantzos 2004a), combining the evidence from observations of the Cas A jet (see e.g., Laming et al. 2006) and fast-moving knots (see e.g. Fesen et al. 2006) with supernova explosion models where polar regions show higher entropy caused by presupernova stellar rotation (Maeda and Nomoto 2003). This is reinforced by comparisons of ^{56}Ni and ^{44}Ti yields for SN1987A and by interpreting the solar abundances of ^{56}Fe and ^{44}Ca in terms of their radioactive parents of ^{56}Ni and ^{44}Ti , respectively (The et al. 2006;

Prantzos 2004a). In all cases, the ^{44}Ti yields of 1-D models for supernova production relative to ^{56}Ni appear lower than indicated by these measurements by a factor up to three. Note, however, that a recent redetermination of the rate of the key production reaction $^{40}\text{Ca}(\alpha, \gamma)^{44}\text{Ti}$ (Nassar et al. 2006) suggests a factor of about 2 increase in model yields, which may weaken such reasoning.

No ^{44}Ti radioactivity has been seen clearly from any other core-collapse supernova, either in the Compton observatory survey or in INTEGRAL's mission up to now (The et al. 2006; Renaud et al. 2006). This is remarkable, as it implies that ^{44}Ti




Fig. 6. New precision spectroscopy with the imaging spectrometer SPI on INTEGRAL allows to trace line characteristics over different regions along the plane of the Galaxy. The first results on the centroid of the line (data points) are consistent with the expectations from Doppler shifts due to ^{26}Al rotating with the bulk gas throughout the inner Galaxy (simulations shown in color/contours, e.g., Diehl et al. 2006c).

ejection is not common among core-collapse supernovae and that Cas A was an exceptional supernova explosion rather than a typical one. It adds to the mounting evidence from hypernova and gamma-ray burst research, which suggests that variety of explosion types and 3-D effects are typical for core-collapse events, and we may have sampled only a small part of the parameter range spanned by such terminations of the lives of massive stars (Woosley and Bloom 2006). Presolar grains, which are attributed to a core-collapse supernova origin, have been observed to show traces of ^{44}Ti inclusions, sometimes in identified subgrains (Nittler et al. 1996; Zinner 1998). This is yet another indication that supernovae from massive stars do eject ^{44}Ti , at least for some events.

The non-detection of the 1157 keV gamma-ray line with the SPI instrument on INTEGRAL also fits into this picture: if ^{44}Ti ejection occurs mostly from the rapid (rather than slow-moving) inner part of ejecta, velocities could be as high as $7000\text{--}8000\text{ km s}^{-1}$, producing lines too broad to be detectable by SPI (Vink 2005).

^{26}Al IN THE GALAXY: DIAGNOSTICS OF RECENT GALACTIC NUCLEOSYNTHESIS

Diffuse gamma-ray emission at 1.8 MeV from the decay of ^{26}Al in the Galaxy (Prantzos and Diehl 1996) has been mapped by COMPTEL from 9 years of observations with NASA's Compton observatory (1991–2000) (Plüschke et al. 2001) (Fig. 4). A disk-like distribution of the gamma-ray emission with irregularities and prominent intensity in the Cygnus region argues for predominant production of galactic ^{26}Al by massive stars (Prantzos 1993). In comparison, the contributions from AGB stars or novae (which are plausible sources of ^{26}Al found in presolar grains), should be small in the Galaxy. The large numbers of such long-lived objects (each one with small contributions to the total) would result in a smooth glow along the plane of the Galaxy, washing out, for example, the spiral-arm tangential viewing directions which can be seen in the ^{26}Al image (Prantzos and Diehl 1996).

^{26}Al gamma-rays reflect the integrated nucleosynthesis production from many individual sources in the Galaxy. Due to the long decay time, most of the gamma rays are generated while ^{26}Al travels through interstellar space along with other nucleosynthesis ejecta while mixing with ambient interstellar gas. Around clusters of massive stars, we expect a hot and tenuous phase of interstellar gas, which is otherwise hard to observe; radioactivity and ^{26}Al gamma rays are able to convey information about the kinematics of this phase. A balloon-borne experiment reported the ^{26}Al gamma-ray line to be as broad as 6.4 keV, which would correspond to interstellar velocities above 500 km s^{-1} and would imply the existence of large interstellar cavities for such velocities to be maintained over the ^{26}Al decay time (Prantzos and Diehl 1996). Years of data from INTEGRAL/SPI have now resulted in a high-quality line shape measurement of the ^{26}Al line with a similar high-resolution Ge detector (Zinner et al. 1991; Diehl et al. 2006b) (Fig. 5). From this measurement, any astrophysical broadening is found to be on the order of $\sim 1\text{ keV}$, with a firm upper limit (2σ) of 2.8 keV. Therefore, line broadening as expected from large-scale galactic rotation and from interstellar turbulence in massive-star regions is fully compatible with these recent measurements.

With fine spectroscopy and deep exposure of the inner regions of the Galaxy, systematic centroid shifts of the ^{26}Al gamma-ray line were seen, consistent with what is in principle expected from the differential rotation of the galactic disk (Zinner et al. 1991; Diehl et al. 2006b) (Fig. 6). Line centroid offsets are a few tenths of keV only; hence they require the maintenance of fine energy resolution over these multi-year exposures through annealing and careful accounting of the time-variable spectral resolution due to cosmic ray damage. From this signature of galactic rotation, one may conclude that the observed source regions indeed are located in the inner regions of the Galaxy, and that ^{26}Al production is a large-scale, Galaxy-wide phenomenon that is not significantly distorted by localized sources. Therefore, the

observed ^{26}Al gamma-ray flux represents ^{26}Al emission from all sources integrated over the entire Galaxy, and geometrical source distribution models can be used to assign distances. With a model for the massive star population size distribution in the Galaxy, the observed ^{26}Al line flux then translates into a total galactic amount of $2.8 \pm 0.9 M_{\odot}$, where the uncertainty mainly arises from different spatial distribution models for the ^{26}Al sources (Diehl et al. 2006c). Because the ^{26}Al origin is predominantly attributed to massive stars, this ^{26}Al amount can be converted into a core-collapse supernova rate, assuming a steady-state situation and using the ^{26}Al production per stellar mass from nucleosynthesis models with the standard mass distribution function for massive stars (for details, see Zinner et al. 1991; Diehl et al. 2006c). Therefore, a new and systematically different measurement of the rate of stellar births and deaths in our galaxy can be obtained from penetrating gamma rays from the otherwise occulted massive star population in the inner regions of our galaxy. This method relies on the stellar mass distribution for massive stars (which is rather well established; see Kroupa 2002) and nucleosynthetic yields of ^{26}Al (which may be uncertain according to Zinner et al. 1991; see Diehl et al. 2006c); unlike the other methods, no corrections for unobserved distant sources nor Galaxy morphology arguments are important in this case, thus providing a useful independent view on the galactic SN rate. The INTEGRAL/SPI-derived value is $1.9 (\pm 1.1)$ supernovae $(100 \text{ yr})^{-1}$, corresponding to a conversion of $\sim 4 M_{\odot}$ of interstellar gas into stars per year on average in the Galaxy.

Another interesting quantity may be derived from this measurement: the $^{26}\text{Al}/^{27}\text{Al}$ isotopic ratio of the average interstellar medium, which is 8.4×10^{-6} (Zinner et al. 1991); here we assume an interstellar gas mass of $4.95 \times 10^9 M_{\odot}$ and an abundance by number of $\log N(^{27}\text{Al}) = 6.4$ on the astronomical abundance scale with $\log N(\text{H}) = 12$. Note that this $^{26}\text{Al}/^{27}\text{Al}$ ratio is about one order of magnitude lower than the solar nebula value of $\sim 4.5 \times 10^{-5}$ (Lee et al. 1998; Young et al. 2005). This seems plausible from continued production of Al since the birth of the solar system 4.6 Gyr ago and from several other hints for special enrichment of the solar nebula at birth from a nearby nucleosynthesis source (Wasserburg 1998; Clayton and Jin 1995; Goswami et al. 2001; Meyer and Clayton 2000), or from locally produced ^{26}Al enrichments made possible by jet outflows in the early stage of solar nebula accretion onto the central Sun (Gounelle et al. 2006).

^{60}Fe PRODUCTION BY SUPERNOVAE: ANOTHER LONG-LIVED TRACER

Stellar nucleosynthesis models have always predicted that the convective envelopes of massive stars are plausible production sites of long-lived ^{60}Fe isotopes ($\tau \sim 2.2 \text{ Myr}$),

which would be ejected in the terminal core-collapse supernova explosion (Prantzos 1989; Limongi and Chieffi 2005; Woosley and Heger 2007; Limongi and Chieffi 2006). Because of the dominant origin of ^{26}Al in massive stars, the gamma-ray brightness ratio $R_{\text{Fe-Al}} = I_{\gamma}(^{60}\text{Fe})/I_{\gamma}(^{26}\text{Al})$ from the emission of diffuse radioactivities in steady state over the last several million years is a suitable test of the models. ^{60}Fe has been found to be part of meteorites formed in the early solar system (Shukolyukov and Lugmair 1993). The inferred $^{60}\text{Fe}/^{56}\text{Fe}$ ratio for these meteorites exceeded the interstellar medium estimates from nucleosynthesis models, which led to the suggestion of late supernova injection before formation of the solar system (Tachibana and Huss 2003). As another proof for cosmic ^{60}Fe production, accelerator mass spectroscopy of seafloor crust material from the southern Pacific Ocean revealed an ^{60}Fe excess in a crust depth corresponding to an age of 2–3 Myr (Knie et al. 2004). From this measurement, it was concluded that a supernova must have occurred very nearby at this time, depositing some of its debris directly in the Earth's atmosphere. Both of these measurements demonstrate that ^{60}Fe nucleosynthesis does occur; it is now interesting to search for current ^{60}Fe production in the Galaxy through radioactive-decay gamma rays.

A hint for ^{60}Fe gamma-rays was recognized for the first time in 2003 with RHESSI (Smith 2003). The weak signal was found to correspond to $16 (\pm 5)\%$ of the ^{26}Al brightness (Smith 2005). This seemed to confirm earlier predictions of $\sim 15\%$ (Timmes et al. 1995) very well, but it was cast in doubt by later studies of nucleosynthesis in massive stars which predicted much larger ratios (see Prantzos 2004b). In fact, substantial uncertainties both from stellar structure (establishment of suitable convective-burning regions, mass loss) and from nuclear cross sections (neutron capture on unstable ^{59}Fe is difficult to measure and its β decay lifetime is uncertain) still persist.

Confirmation of the RHESSI ^{60}Fe signal was reported from the INTEGRAL/SPI data collected during the first year of the mission (Harris et al. 2005), while a recent analysis with more data finds a significant ^{60}Fe signal (Wang et al. 2007) (at 5σ , Fig. 7). The SPI team finds the $^{60}\text{Fe}/^{26}\text{Al}$ gamma-ray brightness ratio $R_{\text{Fe-Al}}$ of 14%, with an uncertainty of a few percent (Knie et al. 2004; Wang et al. 2007). At first glance, this seems to be a nice confirmation of the original theoretical estimates. However, in view of the substantial revisions to nucleosynthesis models for massive stars and of nuclear reaction rates, this result should be carefully scrutinized. Massive-star nucleosynthesis models are the integrated (and evolving) result of what we know from all observations of massive stars' evolution and their nucleosynthesis in the presupernova stage, and from explosive nucleosynthesis reactions (e.g., Prantzos 1989; Gounelle et al. 2006; Limongi and Chieffi 2006; Woosley and Heger 2007).




Fig. 7. INTEGRAL/SPI ^{60}Fe signal from the inner Galaxy (Knödlseder et al. 2004). Data from both lines have been superimposed, referencing to the laboratory line energies of 1173.23 keV and 1332.49 keV, respectively. The solid line represents a Gaussian fit with its width fixed from nearby instrumental background lines.

ANNIHILATION OF POSITRONS AND OTHER TRACES OF RADIOACTIVITY

Nuclear processes in stars, supernovae, and novae are expected to release positrons from radioactive decays of isotopes on the proton-rich side of the valley of stability. Positrons do annihilate in our galaxy, as revealed through their characteristic annihilation gamma rays at 511 keV: this was the first gamma-ray line detected from cosmic sources in the Galaxy (for a recent review, see Diehl et al. 2006a). In recent years, INTEGRAL/SPI made it possible to map this gamma-ray emission over large regions of the sky (Knödlseder et al. 2003; Knödlseder et al. 2005). But the map of annihilation emission resulted in a surprise that puzzles a broad community of astrophysicists (see detailed discussion and references in Smith 2005): rather than following the plausible spatial distribution of nucleosynthesis along the disk of the Galaxy, annihilation emission is bright in an extended yet surprisingly spherically symmetric region, broadly coinciding with the galactic bulge. Although electron-positron pair production is expected from compact stars (strong magnetic fields of rapidly rotating neutron stars result in particle-accelerating electric potential gaps; accretion flow onto a compact star in a binary system results in plasma jets such as observed in microquasars), it was expected that nucleosynthesis sources would dominate. Annihilation emission appears to arise from the bulge region predominantly, outshining the weak galactic disk. The observed annihilation map (Smith 2003; Smith 2005; Knödlseder et al. 2003; Knödlseder et al. 2005) prompted several interesting ideas on cosmic-ray production and propagation (Prantzos 2006), other positron source types (Guessoum et al. 2006), and even dark matter physics

(Hooper and Silk 2005; Ascascibar et al. 2006). At present, it is not clear whether nucleosynthesis is the major source of galactic positrons. It should be noted, however, that nucleosynthesis positrons are released at low energies (MeV) and do not produce high-energy gamma rays, contrary to positrons released in most of other (non-thermal) processes; nucleosynthesis then naturally satisfies constraints imposed from the observed galactic emissivity at a few tens of MeV (Beacom and Yüksel 2006).

Novae are the most common variety of explosive events in the Galaxy. They are powered by nuclear reactions under runaway conditions on the surfaces of white dwarfs in binary systems (Gehrtz et al. 1998). Nuclear energy generation allows the overlying envelope to expand, extinguishing and partially uncovering the nuclear burning region. In these events, production of proton-rich isotopes is expected under the conditions of explosive hydrogen burning, and thus transient positron annihilation gamma-ray emission should be bright (Jose and Hernanz 1998). The key isotopes are too short-lived, however, and the gamma-ray flash occurs before the nova brightens enough in optical light to be discovered; sky surveys with large field-of-view gamma-ray telescopes are required. So far, none of these flashes has been discovered (Harris et al. 2000). Other longer-lived radioactive isotopes could lead to observable gamma-ray line emission: ^7Be with a line at 478 keV and a decay lifetime of 78 days may still be substantially occulted, burried under the expanding envelope. But ^{22}Na should be sufficiently long-lived (lifetime 3.8 years), such that it would decay in the diluted nova wind and be observable from novae occurring out to distances of approximately one kiloparsec (Prantzos 2006; Jose and Hernanz 1998). None of these have been clearly observed, but there are a few tantalizing reports (Iyudin et al. 1999); yet upper limits are consistent with expectations from current nova nucleosynthesis models (Iyudin et al. 1995; Jose et al. 2006).

SUMMARY AND PROSPECTS

The Compton Gamma-Ray Observatory mission opened the era of cosmic gamma-ray line astronomy in the 1990s, after experimental efforts through a variety of balloon and rocket campaigns had shown the promise of this field. The new high-resolution spectroscopy of cosmic gamma-rays with spaceborne Ge detectors has demonstrated new aspects of gamma-ray lines from nuclear and high-energy non-thermal processes: ^{44}Ti , ^{26}Al and ^{60}Fe radioactive decays have been measured from cosmic sources providing information on nuclear reactions inside massive stars. The width and centroid of ^{44}Ti lines constrain the kinematics of inner ejecta from core-collapse supernovae. Similarly, the line shape and centroid of the ^{26}Al line reflects nucleosynthesis yields and their dynamics in otherwise hardly accessible hot and tenuous gas regions near nucleosynthesis sources. With ^{60}Fe gamma-

rays and their intensity ratio to those of ^{26}Al , the models for massive-star nucleosynthesis can be tested in a global sense. Although it is difficult to combine imaging and spectroscopy in an astronomical instrument for gamma-rays, the results from recent years demonstrate that such combination may open new frontiers, as in the case of the positron annihilation gamma-ray line with its puzzling morphology, dominated by an extended inner-Galaxy source.

INTEGRAL will continue to operate well into the next decade and is expected to provide a gamma-ray line survey of the Galaxy down to the level of 10^{-5} ph cm $^{-2}$ s $^{-1}$. This will reveal only the brightest sources of nuclear gamma-ray lines, but it will prepare the ground for deeper exploration of cosmic sources. We hope that it will be succeeded by future experiments with higher sensitivity, which are presently planned by the international space science community. We are confident that a deeper exploration of the nuclear universe will greatly improve our understanding of cosmic element production.

Acknowledgments—This contribution results from the collaborative work with numerous colleagues in the gamma-ray and nuclear astrophysics community, and our colleagues who construct theoretical models for the sources of nucleosynthesis. We are grateful for the lively discussions with all of them that we enjoyed in the past and for an active decade of gamma-ray line astrophysics. We acknowledge the fine observational efforts in nuclear astronomy, especially with NASA's Compton observatory and ESA's INTEGRAL mission.

Editorial Handling—Dr. Ian Lyon

REFERENCES

- Ahmad I. P., Greene E. F., Moore S., Ghelberg A., Ofan M., Paul M., and Kutschera W. Improved measurement of the ^{44}Ti half-life from a 14-year long study. *Physical Review C* 74:065803.
- Anders E. and Zinner E. 1993. Interstellar grains in primitive meteorites—Diamond, silicon carbide, and graphite. *Meteoritics* 28:490–514.
- Arnett D. 1969. A possible model of supernovae: Detonation of ^{12}C . *Astrophysics and Space Science* 5:180.
- Ascasibar Y., Jean P., Boehm C., and Knödlseeder J. 2006. Constraints on dark matter and the shape of the Milky Way dark halo from the 511 keV line. *Monthly Notices of the Royal Astronomical Society* 368:1695–1705.
- Beacom J. and Yüksel H. 2006. Stringent constraint on galactic positron production. *Physical Review Letters* 97:7.
- Bloemen H., Wijnands R., Bennett K., Diehl R., Hermsen W., Lichti G., Morris D., Ryan J., Schonfelder V., Strong A. W., Swanenburg B. N., de Vries C., and Winkler C. 1994. COMPTEL observations of the Orion complex: Evidence for cosmic-ray induced gamma-ray lines. *Astronomy & Astrophysics* 281:L5–L8.
- Bloemen H., Morris D., Knödlseeder J., Bennett K., Diehl R., Hermsen W., Lichti G., van der Meulen R. D., Oberlack U., Ryan J., Schönfelder V., Strong A. W., de Vries C., and Winkler C. The revised COMPTEL Orion results. *The Astrophysical Journal* 521:L137–L140.
- Boggs S. 2006. The Advanced Compton Telescope mission. *New Astronomy Reviews* 50:604–607.
- Burbidge M. E., Burbidge G. R., Fowler W. A., and Hoyle F. 1957. Synthesis of the elements in stars. *Review of Modern Physics* 29:547–650.
- Cayrel R. and Cayrel de Strobel G. 1966. Abundance determinations from stellar spectra. *Annual Review of Astronomy and Astrophysics* 4:1.
- Cayrel de Strobel G., Soubiran C., and Ralite N. 2001. Catalogue of (Fe/H) determinations for FGK stars: 2001 edition. *Astronomy & Astrophysics* 373:159–163.
- Cameron A. G. W. 1970. Extinct radioactivities. *Comments on Astrophysics & Space Physics* 12:18.
- Clayton D. D. 1968. *Principles of stellar evolution and nucleosynthesis*. Chicago: University of Chicago Press. 634 p.
- Clayton D. D. and Jin J. 1995. Gamma rays, cosmic rays, and extinct radioactivity in molecular clouds. *The Astrophysical Journal* 451:681.
- Clayton D. D. and Nittler L. 2004. Astrophysics with presolar stardust. *Annual Review of Astronomy and Astrophysics* 42:39–78.
- Diehl R., Dupraz C., Bennett K., Bloemen H., Hermsen W., Knödlseeder J., Lichti G., Morris D., Ryan J., Schoenfelder V., Steinle H., Strong A., Swanenburg B., Varendorff M., and Winkler C. 1995. COMPTEL observations of galactic ^{26}Al emission. *Astronomy & Astrophysics* 298:445.
- Diehl R. and Timmes F. X. 1998. Gamma-ray line emission from radioactive isotopes in stars and galaxies. *Publications of the Astronomical Society of the Pacific* 110:637–659.
- Diehl R. and Hartmann D. H. 2000. Astronomy with radioactivities. *Publications of the Astronomical Society of the Pacific* 112:1278–1279.
- Diehl R., Hartmann D. H., Hoppe P., and Prantzos N. 2002. Astronomy with radioactivities. *New Astronomy Reviews* 46:8–10.
- Diehl R., Hartmann D. H., Kanbach G., Korschinek G., Knödlseeder J., Ott U., Prantzos N., and Schönfelder V. 2004. Preface to “Astronomy with Radioactivities IV and Filling the Sensitivity Gap in MeV Astronomy.” *New Astronomy Reviews* 48:1–2.
- Diehl R., Prantzos N., and Ballmoos P. von. 2006a. Astrophysical constraints from gamma-ray spectroscopy. *Nuclear Physics A* 777:70–97.
- Diehl R., Halloin H., Kretschmer K., Strong A. W., Wang W., Jean P., Lichti G. G., Knödlseeder J., Roques J. P., Schanne S., Schönfelder V., Kienlin A. von, Weidenspointner G., Winkler C., and Wunderer C. 2006b. ^{26}Al in the inner Galaxy: Large-scale characteristics derived with INTEGRAL/SPI. *Astronomy & Astrophysics* 449:1025–1031.
- Diehl R., Halloin H., Kretschmer K., Lichti, Giseler G., Schönfelder V., Strong A. W., Kienlin A. von, Wang W., Jean P., Knödlseeder J., Roques J. P., Weidenspointner G., Schanne S., Hartmann D. H., Winkler C., and Wunderer C. 2006c. Radioactive ^{26}Al from massive stars in the Galaxy. *Nature* 439:45–47.
- Edvardsson B., Andersen J., Gustafsson B., Lambert D. L., Nissen P. E., and Tomkin J. 1993. The chemical evolution of the galactic disk. I—Analysis and results. *Astronomy & Astrophysics* 275:101.
- Fesen R. A., Hammell M. C., Morse J., Chevalier R. A., Borkowski K. J., Dopita M. A., Gerardy C. L., Lawrence S. S., Raymond J. C., and van den Bergh S.. 2006. Discovery of outlying high-velocity oxygen-rich ejecta in Cassiopeia A. *The Astrophysical Journal* 636:859–872.
- Gehrels N., Fichtel C. E., Fishman G. J., Kurfess J. D., and Schonfelder V. 1993. The Compton Gamma-Ray Observatory. *Scientific American* 269:68.

- Gehrz R. D., Truran J. W., Williams R. E., and Starrfield S. 1998. Nucleosynthesis in classical novae and its contribution to the interstellar medium. *Publications of the Astronomical Society of the Pacific* 110:3–26.
- Georgii R., Plüschke S., Diehl R., Lichti G. G., Schönfelder V., Bloemen H., Hermesen W., Ryan J., and Bennett K. 2002. COMPTEL upper limits for the ^{56}Co gamma ray emission from SN1998bu. *Astronomy & Astrophysics* 394:517–523.
- Goswami J. N., Marhas K. K., and Sahijpal S. 2001. Did solar energetic particles produce the short-lived nuclides present in the early solar system? *The Astrophysical Journal* 549:1151–1159.
- Gounelle M., Shu F. H., Shang H., Glassgold A. E., Rehm K. E., and Lee T. 2006. The irradiation origin of beryllium radioisotopes and other short-lived radionuclides. *The Astrophysical Journal* 640:1163–1170.
- Gratton R., Caretta E., Matteucci F., and Sneden C. 2000. Abundances of light elements in metal-poor stars. IV. (Fe/O) and (Fe/Mg) ratios and the history of star formation in the solar neighborhood. *Astronomy & Astrophysics* 358:671–681.
- Gros M., Tatischeff V., Kiener J., Cordier B., Chapuis C., Weidenspointner G., Vedrenne G., Kienlin A. von, Diehl R., Bykov A., and Méndez M. 2004. INTEGRAL/SPI observation of the 2003 Oct 28 solar flare. ESA Special Publication #552. pp. 669.
- Guessoum N., Jean P., and Gillard W. 2005. The lives and deaths of positrons in the interstellar medium. *Astronomy & Astrophysics* 436:171.
- Guessoum N., Jean P., and Prantzos N. 2006. Microquasars as sources of positron annihilation radiation. *Astronomy & Astrophysics* 457:753.
- Harris M. J., Teegarden B. J., Cline T. L., Gehrels N., Palmer D. M., Ramaty R., and Seifert H. 2000. Transient Gamma-Ray Spectrometer observations of gamma-ray lines from novae. II. Constraining the galactic nova rate from a survey of the southern sky during 1995–1997. *The Astrophysical Journal* 542:1057–1063.
- Harris M. J., Knödseder J., Jean P., Cisana E., Diehl R., Lichti G. G., Roques J.-P., Schanne S., and Weidenspointner G. 2005. Detection of gamma-ray lines from interstellar ^{60}Fe by the high-resolution spectrometer SPI. *Astronomy & Astrophysics* 433: L49–L52.
- Hartmann D. H., Diehl R., Prantzos N., and Zinner E. 2006. Astronomy with radioactivities. *New Astronomy Reviews* 50:7–8, 469–447.
- Höflich P. 2006. Physics of type Ia supernovae. *Nuclear Physics A* 777:579–600.
- Hooper D. and Silk J. 2005. Searching for dark matter with future cosmic positron experiments. *Physical Review D* 71.
- Hughes J. P., Rakowski C. E., Burrows D. N., and Slane P. O. 2000. Nucleosynthesis and mixing in Cassiopeia A. *The Astrophysical Journal* 528:L109–L113.
- Hurford G. J., Krucker S., Lin R. P., Schwartz R. A., Share G. H., and Smith D. M. 2006. Gamma-ray imaging of the 2003 Oct/Nov solar flares. *The Astrophysical Journal* 644:L93–L96.
- Isern J., Bravo E., Hirschmann A., and García-Senz D. 2004. Gamma-ray emission from type Ia supernovae. *New Astronomy Reviews* 48:31–33.
- Iyudin A. F., Diehl R., Bloemen H., Hermesen W., Lichti G. G., Morris D., Ryan J., Schoenfelder V., Steinle H., Varendorff M., de Vries C., and Winkler C. 1994. COMPTEL observations of ^{44}Ti gamma-ray line emission from Cas A. *Astronomy & Astrophysics* 284:L1–L4.
- Iyudin A. F., Bennett K., Bloemen H., Diehl R., Hermesen W., Lichti G. G., Morris D., Ryan J., Schoenfelder V., Steinle H., Strong A., Varendorff M., and Winkler C. 1995. COMPTEL search for ^{22}Na line emission from recent novae. *Astronomy & Astrophysics* 300: 422.
- Iyudin A. F., Schönfelder V., Bennett K., Bloemen H., Diehl R., Hermesen W., Lichti G. G., van der Meulen R. D., Ryan J., and Winkler C. 1998. Emission from ^{44}Ti associated with a previously unknown galactic supernova. *Nature* 369:142–144.
- Iyudin A. F., Bennett K., Bloemen H., Diehl R., Hermesen W., Knödseder J., Lichti G. G., Ryan J., Schönfelder V., Strong W. A., and Winkler C. 1999. COMPTEL constraints on nova-produced ^{22}Na . *Astrophysical Letters and Communications* 38:371.
- Johnson W. N. and Haymes R. C. 1973. Detection of a gamma-ray spectral line from the galactic center region. *The Astrophysical Journal* 184:103–126.
- Jose J. and Hernanz M. 1998. Nucleosynthesis in classical novae: CO versus ONe white dwarfs. *The Astrophysical Journal* 494:680.
- Jose J., Hernanz M., and Iliadis C. 2006. Nucleosynthesis in classical novae. *Nuclear Physics A* 777:550.
- Knie K., Korschinek G., Faestermann T., Dorfi E. A., Rugel G., and Wallner A. 2004. ^{60}Fe anomaly in a deep-sea manganese crust and its implications for a nearby supernova source. *Physical Review Letters* 93:171103.
- Knödseder J., Lonjou V., Jean P., Allain M., Mandrou P., Roques J.-P., Skinner G. K., Vedrenne G., Ballmoos P. von, Weidenspointner G., Caraveo P., Cordier B., Schönfelder V., and Teegarden B. J. 2003. Early SPI/INTEGRAL constraints on the morphology of the 511 keV line emission in the 4th galactic quadrant. *Astronomy & Astrophysics* 411:L457–L460.
- Knödseder J., Jean P., Lonjou V., Weidenspointner G., Guessoum N., Gillard W., Skinner G., von Ballmoos P., Vedrenne G., Roques J.-P., Schanne S., Teegarden B., Schönfelder V., and Winkler C. 2005. The all-sky distribution of 511 keV electron-positron annihilation emission. *Astronomy & Astrophysics* 441:513–532.
- Knödseder J. 2006. GRI: The Gamma-Ray Imager mission. *Proceedings of SPIE* 6266:626623.
- Kroupa P. 2002. The initial mass function of stars: Evidence for uniformity in variable systems. *Science* 295:82–91.
- Kurfess J. D., Johnson W. N., Kinzer R. L., Kroeger R. A., Strickman M. S., Grove J. E., Leising M. D., Clayton D. D., Grabelsky D. A., Purcell W. R., Ulmer M. P., Cameron R. A., and Jung G. V. 1992. Oriented Scintillation Spectrometer Experiment observations of ^{57}Co in SN 1987A. *The Astrophysical Journal* 399:L137–L140.
- Laming J. M., Hwang U., Radics B., Lekli G., and Takács E. 2006. The polar regions of Cassiopeia A: The aftermath of a gamma-ray burst? *The Astrophysical Journal* 644:260–273.
- Lee T., Shu F. H., Shang H., Glassgold A. E., and Rehm K. E. 1998. Protostellar cosmic rays and extinct radioactivities in meteorites. *The Astrophysical Journal* 506:898–912.
- Leibundgut B. 2000. Type Ia supernovae. *Astronomy & Astrophysics Review* 10:179–209.
- Limongi M. and Chieffi A. 2005. ^{26}Al and ^{60}Fe from massive stars. *Nuclear Physics A* 758:11.
- Limongi M. and Chieffi A. 2006. The nucleosynthesis of ^{26}Al and ^{60}Fe in solar metallicity stars extending in mass from 11 to 120 $M_{\odot}$: The hydrostatic and explosive contributions. *The Astrophysical Journal* 647:483–500.
- Lewis M., Madden N., van Beek H. F., Appleby M., Raudorf T., McTiernan J., Ramaty R., Schmahl E., Schwartz R., Krucker S., Abiad R., Quinn T., Berg P., Hashii M., Sterling R., Jackson R., Pratt R., Campbell R. D., Malone D., Landis D., Barrington-Leigh C. P., Slassi-Sennou S., Cork C., Clark D., Amato D., Orwig L., Boyle R., Banks I. S., Shirey K., Tolbert A. K., Zarro D., Snow F., Thomsen K., Henneck R., McHedlishvili A., Ming P., Fivian M., Jordan J., Wanner R., Crubb J., Preble J., Matranga M., Benz A., Hudson H., Canfield R. C., Holman G. D.,

- Crannell C., Kosugi T., Emslie A. G., Vilmer N., Brown J. C., Johns-Krull C., Aschwanden M., Metcalf T., and Conway A. 2002. The Reuven Ramaty High-Energy Solar Spectroscopic Imager (RHESSI). *Solar Physics* 210:3–32.
- Maeda K. and Nomoto K. 2003. Bipolar supernova explosions: Nucleosynthesis and implications for abundances in extremely metal-poor stars. *The Astrophysical Journal* 598:1163–1200.
- Matz S. M., Share G. H., Leising M. D., Chupp E. L., and Vestrand W. T. 1988. Gamma-ray line emission from SN1987A. *Nature* 331:416–418.
- McConnell M., Fletcher S., Bennett K., Bloemen H., Diehl R., Hermsen W., Ryan J., Schonfelder V., Strong A., and van Dijk R. 1997. COMPTEL all-sky imaging at 2.2 MeV. *AIP Conference Proceedings* 410:1099.
- Meyer B. S. and Clayton D. D. 2000. Short-lived radioactivities and the birth of the sun. *Space Science Reviews* 92:133–152.
- Morris D. J., Bennett K., Bloemen H., Diehl R., Hermsen W., Lichti G. G., McConnell M. L., Ryan J. M., and Schonfelder V. 1997. Reassessment of the ^{56}Co emission from SN 1991T. *AIP Conference Proceedings* 410:1084.
- Mochizuki Y., Takahashi K., Janka H.-Th., Hillebrandt W., Diehl R. 1999. ^{44}Ti : Its effective decay rate in young supernova remnants, and its abundance in Cassiopeia A. *Astronomy & Astrophysics* 346:831–842.
- Murphy R. J. and Share G. H. 2005. What gamma-ray deexcitation lines reveal about solar-flares. *Advances in Space Research* 35: 1825–1832.
- Nassar H., Paul M., Ahmad I., Ben-Dov Y., Caggiano J., Ghelberg S., Goriely S., Greene J. P., Hass M., Heger A., Heinz A., Henderson D. J., Janssens R. V., Jiang C. L., Kashiv Y., Nara Singh B. S., Ofan A., Pardo R. C., Pennington T., Rehm K. E., Savard G., Scott R., and Vondrasek R. 2006. $^{40}\text{Ca}(\alpha, \gamma)^{44}\text{Ti}$ reaction in the energy regime of supernova nucleosynthesis. *Physical Review Letters* 96:4.
- Nittler L., Amari S., Zinner E., Woosley S. E., and Lewis R. S. 1996. Extinct ^{44}Ti in presolar graphite and SiC: Proof of a supernova origin. *The Astrophysical Journal* 462:L31–L34.
- Plüschke S., Diehl R., Schönfelder V., Bloemen H., Hermsen W., Bennett K., Winkler C., McConnell M., Ryan J., Oberlack U., and Knödseder J. 2001. The COMPTEL 1.809 MeV survey. ESA Special Publication #459. pp. 55–58.
- Prantzos N. 1989. Gamma-ray lines from radioactive nuclei produced in hydrostatic stellar burning phases. *Astronomy & Astrophysics* 223:136.
- Prantzos N. 1993. Radioactive ^{26}Al from massive stars: Production and distribution in the Galaxy. *The Astrophysical Journal* 405: L55.
- Prantzos N. and Diehl R. 1996. Radioactive ^{26}Al in the Galaxy: Observations versus theory. *Physics Reports* 267:1.
- Prantzos N. 2004a. Astrophysical-ray lines: A probe of stellar nucleosynthesis and star formation. ESA Special Publication #552. p. 15.
- Prantzos N. 2004b. Radioactive ^{26}Al and ^{60}Fe in the Milky Way: Implications of the RHESSI detection of ^{60}Fe . *Astronomy & Astrophysics* 420:1033.
- Prantzos N. 2006. On the intensity and spatial morphology of the 511 keV emission in the Milky Way. *Astronomy & Astrophysics* 449:869–878.
- Purcell W. R., Cheng L.-X., Dixon D. D., Kinzer R. L., Kurfess J. D., Leventhal M., Saunders M. A., Skibo J. G., Smith D. M., and Tueller J. 1997. OSSE mapping of galactic 511 keV positron annihilation line emission. *The Astrophysical Journal* 491:725.
- Reed J. E., Hester J. J., Fabian A. C., and Winkler P. F. 1995. The three-dimensional structure of the Cassiopeia A supernova remnant. I. The spherical shell. *The Astrophysical Journal* 440: 706.
- Renaud M., Vink J., Decourchelle A., Lebrun F., den Hartog P. R., Terrier R., Couvreur C., Knödseder J., Martin P., Prantzos N., Bykov A. M., and Bloemen H. 2006. The signature of ^{44}Ti in Cassiopeia A revealed by IBIS/ISGRI on INTEGRAL. *The Astrophysical Journal* 647:L41–L44.
- Röpke F. K., Hillebrandt W., and Blinnikov S. I. 2006. On the mechanism of Type Ia supernovae. astro-ph/0609631.
- Roques J. P., Schanne S., Kienlin A. von, Knödseder J., Briet R., Bouchet L., Paul Ph., Boggs S., Caraveo P., Cassé M., Cordier B., Diehl R., Durouchoux P., Jean P., Leleux P., Lichti G., Mandrou P., Matteson J., Sanchez F., Schönfelder V., Skinner G., Strong A., Teegarden B., Vedrenne G., von Ballmoos P., and Wunderer C. 2003. SPI/INTEGRAL in-flight performance. *Astronomy & Astrophysics* 411:L91–L100.
- Schönfelder V., Bloemen H., Collmar W., Diehl R., Hermsen W., Knödseder J., Lichti G. G., Plüschke S., Ryan J., Strong A., and Winkler C. 2000. ^{44}Ti gamma-ray line emission from Cas A and RXJ0852-4622/GROJ0852-4642. *AIP Conference Proceedings* 510:54.
- Shukolyukov A. and Lugmair G. W. 1993. Live ^{60}Fe in the early solar system. *Science* 259:598–1138.
- Smith D. 2003. The Reuven Ramaty High-Energy Solar Spectroscopic Imager observation of the 1809 keV line from galactic ^{26}Al . *The Astrophysical Journal* 589:L55.
- Smith D. 2005. Gamma-ray line observations with RHESSI. ESA Special Publication #552. p. 45.
- Tachibana S. and Huss G. R. 2003. The initial abundance of ^{60}Fe in the solar system. *The Astrophysical Journal* 588:L41.
- Timmes F. X., Woosley S. E., Hartmann D. H., Hoffman R. D., Weaver T. A., and Matteucci F. 1995. ^{26}Al and ^{60}Fe from supernova explosions. *The Astrophysical Journal* 449:204.
- The L.-S., Clayton D. D., Diehl R., Hartmann D. H., Iyudin A. F., Leising M. D., Meyer B. S., Motizuki Y., and Schönfelder V. 2006. Are ^{44}Ti -producing supernovae exceptional? *Astronomy & Astrophysics* 450:1037–1050.
- Truran J. W. and Cameron A. G. W. 1971. Evolutionary models of nucleosynthesis in the Galaxy. *Astrophysics & Space Science* 14: 179.
- Vedrenne G., Roques J.-P., Schönfelder V., Mandrou P., Lichti G. G., von Kienlin A., Cordier B., Schanne S., Knödseder J., Skinner G., Jean P., Sanchez F., Caraveo P., Teegarden B., von Ballmoos P., Bouchet L., Paul P., Matteson J., Boggs S., Wunderer C., Leleux P., Weidenspointner G., Durouchoux Ph., Diehl R., Strong A., Cassé M., Clair M. A., and André Y. 2003. SPI: The spectrometer aboard INTEGRAL. *Astronomy & Astrophysics* 411:L63–L70.
- Vink J., Laming J. M., Kaastra J. S., Bleeker J. A. M., Bloemen H., and Oberlack U. 2001. Detection of the 67.9 and 78.4 keV lines associated with the radioactive decay of ^{44}Ti in Cassiopeia A. *The Astrophysical Journal* 560:L79–L82.
- Vink J. 2005. Gamma-ray observations of explosive nucleosynthesis products. *Advances in Space Research* 35:976.
- Wallerstein G., Iben I., Jr., Parker P., Boesgaard A. M., Hale G. M., Champagne A. E., Barnes C. A., Käppeler F., Smith V. V., Hoffman R. D., Timmes F. X., Sneden C., Boyd R. N., Meyer B. S., and Lambert D. L. 1997. Synthesis of the elements in stars: Forty years of progress. *Review of Modern Physics* 69:995–1084.
- Wang W., Harris M., Diehl R., Halloin H., Cordier B., Strong A. W., Kretschmer K., Knödseder J., Jean P., Lichti G. G., Roques J. P., Schanne S., Kienlin A. von, Weidenspointner G., and Wunderer C. 2006. SPI observations of the diffuse ^{60}Fe emission in the Galaxy. *Astronomy & Astrophysics* 469:1004–1012.
- Wasserburg G. J., Gallino R., and Busso M. 1998. A test of the supernova trigger hypothesis with ^{26}Al and ^{60}Fe . *The Astrophysical Journal* 500:L189.

- Winkler C., Courvoisier T. J.-L., Di Cocco G., Gehrels N., Giménez A., Grebenev S., Hermsen W., Mas-Hesse J. M., Lebrun F., Lund N., Palumbo G. G. C., Paul J., Roques J. P., Schnopper H., Schönfelder V., Sunyaev R., Teegarden B., Ubertini P., Vedrenne G., and Dean A. J. 2003. The INTEGRAL mission. *Astronomy & Astrophysics* 411:L1–L6.
- Woosley S. E., Hartmann D. H., and Pinto P. 1989. Hard emission at late times from SN 1987A. *The Astrophysical Journal* 346:395.
- Woosley S. E. and Bloom J. S. 2006. The supernova gamma-ray burst connection. *Annual Review of Astronomy & Astrophysics* 44:507.
- Woosley S. E. and Heger A. 2007. Nucleosynthesis and remnants in massive stars of solar metallicity. Hans A. Bethe Prize lecture, American Physical Society Meeting. *Physics Reports* 442:269–283.
- Young E. D., Simon J. I., Galy A., Russell S. S., Tonui E., and Lovera O. 2005. Supra-canonical $^{26}\text{Al}/^{27}\text{Al}$ and the residence time of CAIs in the solar protoplanetary disk. *Science* 308:223–227.
- Zinner E., Tang M., and Anders E. 1987. Large isotopic anomalies of Si, C, N, and noble gases in interstellar silicon carbide from the Murray meteorite. *Nature* 330:730.
- Zinner E., Amari S., Anders E., and Lewis R. S. 1991. Large amounts of extinct ^{26}Al in interstellar grains from the Murchison meteorite. *Nature* 349:51–54.
- Zinner E. 1998. Stellar nucleosynthesis and the isotopic composition of presolar grains from meteorites. *Annual Review of Earth and Planetary Sciences* 26:147.
- Zinner E. 1998. Stellar nucleosynthesis and the isotopic composition of presolar grains from primitive meteorites. *Annual Review of Earth and Planetary Sciences* 26:147.
-