

This article was downloaded by: [Stanford University Libraries]

On: 15 May 2012, At: 04:12

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## International Geology Review

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tigr20>

### Metamorphic Conditions in Orogenic Belts: A Record of Secular Change

M. Brown<sup>a</sup>

<sup>a</sup> University of Maryland, College Park

Available online: 06 Aug 2010

To cite this article: M. Brown (2007): Metamorphic Conditions in Orogenic Belts: A Record of Secular Change, International Geology Review, 49:3, 193-234

To link to this article: <http://dx.doi.org/10.2747/0020-6814.49.3.193>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## Metamorphic Conditions in Orogenic Belts: A Record of Secular Change

M. BROWN<sup>1</sup>

*Laboratory for Crustal Petrology, Department of Geology, University of Maryland, College Park, Maryland 20742*

### Abstract

In general, Archean rocks exhibit rather ordinary moderate-P–high-T facies series metamorphism; neither blueschists nor any record of deep continental subduction and return are documented. However, the abundance and scale of ultrahigh-temperature (UHT) metamorphic belts from the Neoproterozoic to the Cambrian imply a significant change in geodynamics during the Neoproterozoic Era, after which transient sites of high heat flow were available at intervals throughout this period of Earth evolution. Many Neoproterozoic–Cambrian UHT metamorphic belts appear to have developed in settings analogous to modern backarcs that were closed and inverted during crustal aggregation and formation of the Gondwana supercontinent. If backarcs were the general setting for UHT metamorphism, then on a hotter Earth the cyclic formation of supercratons (in the Neoproterozoic Era) and supercontinents (in the Proterozoic Eon) required the destruction of oceans floored by thinner lithosphere that may have generated hotter backarcs than those associated with the current destruction of the Pacific Ocean on the modern Earth. The inherent weakness of the lithosphere in a hotter thermal regime inevitably localized magmatism and deformation at these sites contemporaneously with UHT metamorphism. Medium-temperature eclogites of crustal derivation and associated high-pressure granulites are also first recognized in the Neoproterozoic, for example within the Belomorian Mobile Belt, and they occur at intervals throughout the Proterozoic, for example in the Orosirian Usagaran Orogen and in the Grenvillian belts of the Proto-Atlantic region, and Paleozoic, for example in the circum-North Atlantic Caledonides and European Variscides. Eclogite–high-pressure granulite (E-HPG) metamorphism is predominantly a Proterozoic–Paleozoic phenomenon—complementary to but sparser than UHT metamorphism to begin with, but extending further into the Paleozoic than does UHT metamorphism—that is inferred to record subduction-to-collision orogenesis.

Blueschists appear in the Neoproterozoic Era, becoming common through the Phanerozoic Eon; they record the low thermal gradients associated with modern subduction. Lawsonite-bearing blueschists and eclogites, and ultrahigh-pressure (UHP) metamorphism characterized by coesite or diamond are predominantly Phanerozoic phenomena related to deep subduction within subduction-to-collision orogens. In general, UHP metamorphic belts in subduction-to-collision orogens are not associated with a contemporary magmatic arc in the hanging wall; this suggests that deep subduction of continental crust may inhibit the generation of calc-alkaline magmas, a feature that may have enabled preservation during exhumation of the mineralogical evidence for extreme pressures. However, an enigma concerning UHP metamorphism is the first evidence of deep subduction of continental crust in the rock record. At issue is the recycling implied by the geochemistry of Archean and Proterozoic diamonds (since entrained from mantle to upper crust in younger magmatic events), which requires a supracrustal component. Are these diamonds evidence of deep subduction of continental crust and an early record of UHP metamorphism, or was some other mechanism (e.g., delamination of underthrust lithosphere, some form of slab break-off) responsible for taking a supracrustal component deep into the mantle source?

The Archean and Proterozoic eons were characterized by higher but decreasing mean mantle temperatures and radioactive heat production (RHP), and a thinner thermal boundary layer (TBL) with a shorter residence time than modern Earth. Modeling the effect of increased RHP on the thermal evolution of crust instantaneously doubled in thickness predicts that metamorphic rocks in Archean collisional orogens should have experienced maximum temperatures several hundreds of degrees Celsius higher than those recorded by metamorphic rocks in modern collisional orogens.

<sup>1</sup>Email: mbrown@geol.umd.edu

However, there is no evidence of this—the Archean record is dominated by rather ordinary P–T conditions and crustal melting at relatively low temperatures probably fluxed by water. I argue that a duality of metamorphic belts—reflecting a duality of thermal environments—is the characteristic metamorphic imprint of plate tectonics in the rock record, and it appears only since the Neoproterozoic transition. Based on the occurrence of both UHT and E-HPG metamorphic belts since the Archean-to-Proterozoic transition, I suggest this transition records the onset of a “*Proterozoic plate tectonics regime*,” although the total ridge length and the number of plates undoubtedly were larger in the beginning than on modern Earth and the style of tectonics likely involved some differences. The Neoproterozoic transition to the “*modern plate tectonics regime*” registers a change to subduction of continental crust deeper into the mantle and its (partial) return from depths of up to 300 km, a change perhaps related to whole mantle convection as oceanic lithosphere became thicker with decreased thermal gradients. Although there was overlap in time and space between E-HPG and UHP metamorphism during the Paleozoic, since the Carboniferous UHP metamorphism has dominated, and extreme thermal metamorphism is rare.

### Dedication

This review is dedicated to honoring the career contributions of J.G. (Louie) Liou. Of course, this is not the end of Louie’s contributions, and the timing of the Liou Symposium in his honor at Stanford University on December 10–11, 2005 was determined more by Louie’s change in status to Emeritus Professor than any expectation that he will slow down in research. However, I do anticipate that the greater flexibility associated with an Emeritus faculty rank will give Louie more time to travel, and to contribute research and write papers—his research career is but partway through! An occasion like the Liou Symposium provides an opportunity for retrospection, for example, to review our current understanding of the thermal conditions and depth of burial implied by the mineralogic record of exhumed deep crust now exposed in eroded remnants of metamorphic belts, and for prospection, for example, by asking how much our view of metamorphic conditions might change within the next decade or two!

### Introduction

METAMORPHISM IS CONCERNED with the interpretation of information recorded by the mineralogy, microstructure, and chemistry of rocks and the application of this information to understand the burial and thermal history of the crust and underlying mantle. Metamorphic processes occur at scales that range from atomic (in minerals) to global (interactions between tectonic plates). On the modern Earth, these processes occur at: (1) convergent plate boundaries, where subduction (e.g., the Andes) or continental collision after a phase of subduction (e.g., the Himalayas) takes crustal rocks

deep into Earth’s mantle during orogeny; (2) at plate boundaries that involve predominantly lateral displacement or that also may involve transpression or transtension (e.g., the San Andreas fault system and associated structures in the Transverse Ranges of the western United States, and the Kermadec Trench and the Alpine fault of New Zealand); and (3) at divergent plate boundaries, where newly generated oceanic crust is metamorphosed following hydrothermal circulation of sea water (ocean-floor metamorphism alongside the mid-ocean ridge system). These diverse tectonic environments generate metamorphic signatures in rocks that span a range in temperature from several hundred to more than a thousand degrees Celsius and in depth from near-surface to several hundred kilometers (Brown, 2001).

Field relations, rock fabrics, mineral assemblages, reaction microstructures, and mineral and whole-rock compositional variation provide a record of orogenic evolution. Our ability to interpret the evidence recorded by metamorphic rocks is critical to understanding their history and to constraining models of orogenesis. The metamorphic cycle involving burial (increasing pressure and temperature) and exhumation (decreasing pressure and temperature) is referred to as a pressure-temperature-time-deformation (P–T–t–d) path. When viewed in a metamorphic P–T diagram, these paths may be clockwise or counterclockwise in sense, according to the relative rates of burial and heating versus cooling and exhumation.

We are concerned with two scales of interaction. At the global scale, similarities in type of metamorphism are likely to reflect geodynamic processes, and changes in type with time are likely to reflect changes in geodynamic processes. At the regional

scale, differences in type of metamorphism from terrane to terrane enable us to characterize the tectonic settings involved and the sequence of events that have produced a particular orogenic collage.

### Why Is This Topical Now?

My objective in this paper is to review the record of metamorphism back through Earth history to evaluate secular change in thermal regimes and assess the link between the metamorphic record and global tectonic regimes. One may question whether there is anything new to say in reviewing metamorphic conditions, particularly since this year is the 50th Anniversary of de Roever's landmark paper "Some Differences between Post-Paleozoic and Older Regional Metamorphism" (de Roever, 1956) and the 10th Anniversary of the massive "Blueschists and Eclogites of the World and their Exhumation" by Maruyama, Liou and Terabayashi (Maruyama et al., 1996). Furthermore, the relationship between secular change in Earth's thermal structure as it loses heat and the metamorphic conditions recorded by exhumed and exposed metamorphic belts, and the relationship between these and global tectonic regimes have generated a succession of contributions during the past five decades. These contributions include those by de Roever, Miyashiro, and Ernst in the early days of the paradigm shift in Earth Sciences with the plate tectonics revolution (de Roever, 1956, 1965; Miyashiro, 1961, 1973; Ernst, 1972), to the more recent contributions by Maruyama and Liou (Maruyama et al., 1996; Maruyama and Liou, 1998, 2005), and by Stern (2005), with only the single paper by Grambling (1981) in the intervening period. However, although the topic of metamorphic conditions and secular change has a 50-year history, several advances have led to: (1) a dramatic increases in the maximum calculated pressures (and implied depths of burial; e.g., Chopin, 2003) and highest calculated temperatures (Harley, 2004) recorded by terrane-scale metamorphic rocks in orogens; and (2) a higher level of confidence in the absolute values of retrieved maximum pressures and temperatures (e.g., Green et al., 2000; Powell et al., 2005), and a better understanding of the relationship between this P-T culmination in the tectonic evolution, and retrieved age data that may relate to P-T points along the prograde, around the peak, or along the retrograde segments of the P-T

path (e.g., Rubatto, 2002; Hodges, 2003; Müller, 2003).

These advances include: (1) the ability to identify, analyze and date indirectly (if contained in zircon) micro-inclusions and assemblages of micro-inclusions in rock-forming minerals, such as garnet and pyroxene (e.g., Chopin, 2003), and accessory minerals, such as zircon (F. Liu et al., 2002, 2003; McClelland et al., 2005); (2) significant improvements in thermobarometry (e.g., Pattison et al., 2003; Ravna and Terry, 2004) and in the range of P-T space accessible using comprehensive internally-consistent thermodynamic datasets (e.g., Holland and Powell, 1998), particularly with reference to activity-composition models for minerals characteristic of extreme P-T conditions (e.g., Holland and Powell, 1996a, 1996b, 2003; Powell and Holland, 1999; Wei and Powell, 2003, 2004, 2006; Kelsey et al., 2004; Dale et al., 2005); (3) the development of new trace element-based thermometers (e.g., Watson et al., 2006); and (4) improvements in the precision of U-Pb age data both by TIMS and SIMS (e.g., Baldwin et al., 2004; Collins et al., 2004), and the refinement of monazite dating by EPMA (e.g., Williams and Jercinovic, 2002; Dahl et al., 2005; Baldwin et al., 2006).

For these reasons, it is a worthwhile exercise to review types of metamorphic belt in relation to tectonic setting, compile the most recent estimates of P-T conditions and age of peak metamorphism for selected metamorphic belts, consider the implications of any change in thermal environment of metamorphism inferred from these data, and to relate any changes to possible non-linear switches in the pattern of mantle convection and the tectonics of the TBL. It is through this process of periodic review and integration of data from different specialties in the Earth sciences that we identify unanswered questions that may direct future research.

### Metamorphism at Extreme P-T Conditions—Definitions

It is only a little more than 25 years since our community in metamorphic geology was introduced to the spectacular mineral assemblages reported from lower crustal rocks exposed on a regional scale in Antarctica that yielded estimates of peak temperature of  $>1000^{\circ}\text{C}$  (e.g., Ellis, 1980; Ellis et al., 1980; Harley and Green, 1982), following earlier reports of assemblages with sapphirine + quartz (Dallwitz, 1968). Only a few years later, the exciting

realm of ultrahigh-pressure metamorphism was revealed through the near simultaneous discovery of coesite in the Western Alps (Chopin, 1984) and Norway (Smith, 1984).

For the purposes of this paper, I follow the proposal by Harley (1998) to use 900°C as an arbitrary lower temperature limit for ultrahigh-temperature (UHT) metamorphism. Therefore, UHT metamorphism is:

*metamorphism of crustal rocks in which peak temperature exceeds 900°C, recognized either by robust thermobarometry or by the presence of a diagnostic mineral assemblage in an appropriate bulk composition and oxidation state, such as assemblages with orthopyroxene + sillimanite + quartz, sapphirine + quartz or spinel + quartz, generally at pressure conditions of sillimanite stability in metapelites.*

Additionally, corundum + quartz has been argued to characterize UHT conditions (e.g., Mouri et al., 2003), but since this mineral pair is metastable relative to sillimanite (e.g., Harlov and Milke, 2002) this mineral pair is not included as part of the definition of UHT metamorphism herein. The current state-of-the-art in robust thermobarometry for granulite metamorphism in general for a variety of bulk compositions is either the method proposed by Pattison et al. (2003) or the average P–T mode in THERMOCALC (Powell et al., 1998) with the internally-consistent thermodynamic dataset of Holland and Powell (1998, most recent update). Also, I define ultrahigh-pressure metamorphism as:

*metamorphism of crustal rocks under conditions of pressure that exceed the stability field of quartz, recognized either by the presence of coesite or diamond, or by equivalent P–T conditions determined using robust thermobarometry.*

Examples of robust thermobarometry currently in use, as recently evaluated by Hacker (2006), include the method of Ravna and Terry (2004) and the average P–T mode in THERMOCALC (Powell et al., 1998) with the internally consistent thermodynamic dataset of Holland and Powell (1998, most recent update).

In between these two extreme types of metamorphism, the transition in pressure from granulite to high-pressure granulite facies conditions is not well defined, but it may be taken to correspond approximately to conditions of the change from sillimanite

to kyanite stability in metapelitic rocks. Furthermore, the field of high-pressure granulite-facies metamorphism overlaps the transition from amphibolite to eclogite in basaltic rocks, which is a transition over more than 1 GPa according to chemical composition of the protolith (O'Brien and Rötzler, 2003).

## The Metamorphic Realm

The metamorphic realm traditionally has been divided into facies, each represented by a set of mineral assemblages repeatedly associated in space and time that reflect equilibration within a range of P–T conditions (e.g., Turner, 1968). The standard metamorphic facies are well established; ignoring very low grade and contact metamorphism, these are the blueschist, greenschist, eclogite, amphibolite, and granulite facies (Fig. 1). Furthermore, the idea that orogenic belts might be characterized by different metamorphic facies series is well understood (Miyashiro, 1961; Zwart, 1967), although the concept of “paired” metamorphic belts is a result of terrane accretion (Brown, 1998a, 1998b, 2002b; and implicit in Miyashiro, 1961) rather than *in situ* generation, as has sometimes been implied.

New discoveries in the last 25 years (discussed above) have led us to realize that conditions of crustal metamorphism have been more extreme than we previously imagined. Thus, the granulite facies now extends to temperatures above 1,000°C, and at the extreme is only limited by the liquidus for crustal rocks (UHT metamorphism; e.g., Harley, 1998), and the eclogite facies extends to pressures of at least 6 GPa, and possibly as much as 10 GPa, for crustal terranes returned to the surface, and extends to at least these pressures for xenolithic materials (UHP metamorphism; e.g., Chopin, 2003; L. Liu et al., 2005), but this facies does not have a defined upper pressure limit anyway. There is a transition between the two referred to as medium-temperature eclogite–high-pressure granulite (E-HPG) metamorphism (e.g., O'Brien and Rötzler, 2003). These extremes in the metamorphic realm reflect a range of thermal regimes characterized by high dT/dP to low dT/dP gradients—essentially the paleo-thermal gradients (Fig. 1).

## Caveats

Metamorphic rocks record evidence of the pressure (from which we infer depth) and temperature




FIG. 1. P-T diagram illustrating conditions of selected metamorphic facies and P-T ranges of different types of metamorphism. HP-UHP metamorphism includes the following: BS = blueschist; AEE = amphibole-epidote eclogite facies; ALE = amphibole-lawsonite eclogite facies; LE = lawsonite-eclogite facies; AE = amphibole-eclogite facies; UHPM = ultrahigh-pressure metamorphism; E-HPG = eclogite-high-pressure granulite metamorphism; G = granulite facies, whereas UHTM = ultrahigh-temperature metamorphic part of the granulite facies.

evolution with time—the P-T-t path—which reflects the thermal environment under which the constituent mineral assemblage(s) equilibrated. The map distribution of peak P-T conditions provides information about the spatial variation in thermal regime(s). Our task is to piece together the changes in depth, temperature, and kinematics with time from the mineralogical, microstructural, and chemical evidence preserved in the rock. This evidence may be incomplete, it may not be clear how one piece relates to another, evidence from two or more different segments of the path may overprint to create ambiguity in interpretation, and connections may be made between segments that do not correctly reflect the actual path. However, as a community during the past decade, we have increased confidence in our ability correctly to interpret this fragmentary record of evidence (e.g., O'Brien, 1997, 1999; Brown, 2001, 2002a; White et al., 2002; O'Brien and Rötzler, 2003; Johnson and Brown, 2004; J. A. Baldwin et al., 2005; Powell et al.,

2005); this confidence was lacking in the early days of studies of metamorphism under extreme P-T conditions (e.g., Green, 2005).

The age of metamorphic events (including closure of minerals to diffusion) and pinning these dates to the depth of metamorphism is critical, since this yields rates of burial and exhumation (e.g., Hodges, 2003; Müller, 2003). When this information is combined with structural data to constrain the kinematics, the dataset may allow the particular tectonic setting in which the metamorphism occurred to be identified. Measured time scales for both thermal and burial-exhumation events have become shorter as precision in dating has increased, and as a result we now find acceptable rates for tectonic processes that would have been considered too fast only a few years ago (e.g., Hermann, 2003; Camacho et al., 2005).

The P-T-t paths recovered by this detective work on rocks are generally described as clockwise (in P-T space; CW) or counter-clockwise (CCW),




FIG. 2. Internally consistent thermobarometry of single rock samples of granulite and UHT metamorphism (Grt–Opx thermobarometry by Pattison et al., 2003 using Al-solubility in Opx corrected for retrograde Fe–Mg exchange).

although the P–T loops may vary from hairpin to open. CW paths may include a segment with significant isentropic decompression at high T, CCW paths may exhibit approximately isobaric cooling at moderate P, and portions of either type may be stepped; however, these features alone are not diagnostic of the tectonic setting. Without correct determination of the P–T–t evolution, including unraveling the effects of overprinting of multiple orogenic events (polymetamorphism) where present or suspected (e.g., Hensen and Zhou, 1995; Hensen et al., 1995), any evaluation of secular change is potentially flawed.

Fortunately, as discussed above, the last decade has seen significant advances in our ability to interrogate rocks. Also, the wider availability of high-precision ages on accessory phases linked to specific microstructural sites and metamorphic P–T conditions, commonly through the identification of mineral assemblages preserved as micro-inclusions in the core, mantle, or rim of zircon grains, has increased confidence that we know the true age of peak metamorphism in metamorphic belts. Thus, we may now assess how the metamorphic facies are dis-

tributed through Earth history and evaluate changes in thermal regime over time.

### Metamorphic Conditions Determined by Thermobarometry

One approach to the record of crustal metamorphism is to calculate P and T using an internally consistent thermodynamic dataset applied to individual samples that have been well characterized in terms of mineralogy, microstructure, and mineral chemistry (e.g., Powell and Holland, 1994). Although uncertainty associated with thermobarometry is large when calculating absolute P and T—10% or more (e.g., Worley and Powell, 2000)—when used for comparative purposes among a suite of samples, thermobarometry yields a good estimate of the range of P–T conditions for different types of metamorphism. I illustrate this point with two examples.

Figure 2 shows a plot of P–T conditions for individual samples of granulite and ultrahigh-temperature metamorphic belts calculated by Pattison et al. (2003, their Appendix 2), using Al-solubility in




FIG. 3. Internally consistent thermobarometry of single rock samples for HP and UHP metamorphism [Grt-Cpx-Ky-Pag-Qtz/Coe thermobarometry using the calibration of Ravna and Terry (2004), with additional data from Hacker (2006) and Tsujimori et al. (2006b)].

Opx, corrected for retrograde Fe-Mg exchange. These data for individual samples define a limited range of  $P$ —about 1 GPa—across more than 400°C, and reflect thermal gradients with high  $dT/dP$ .

For metamorphic belts that exhibit mineralogical characteristics indicative of high-pressure and ultrahigh-pressure metamorphism, Figure 3 shows the range of  $P$ - $T$  conditions calculated using the methodology proposed by Ravna and Terry (2004), based on Grt-Cpx-Ky-Phe-Qtz/Coe thermobarometry, and reported by these authors. Additional data are included from Hacker (2006) and Tsujimori et al. (2006b). The spread of  $P$ - $T$  conditions recorded from HP and UHP metamorphic belts suggests a range of depths of burial are recorded by individual samples, together with some variation in thermal gradient, although in all cases the metamorphism is characterized by low values of  $dT/dP$ .

### Types of Metamorphism Classified by Terrane

I follow Miyashiro (1961) in believing that it is useful in terms of understanding crustal-scale geo-

dynamics to classify metamorphic belts by type. However, I am cognizant of the advice offered by Francis J. Turner in his Presidential Address to the Mineralogical Society of America on November 11, 1969:

... much of petrological effort has been concentrated on attempts to discern, define and interpret regularly recurring patterns of ... metamorphic petrogenesis ... I do not wish to deny the existence of discernable order in petrological phenomena ... but ... every geological event ... is unique. If we concentrate too heavily on discernment of order and pattern, may we perhaps overlook or underrate the unique quality of each ... metamorphic episode? Possibly uniqueness may have as great a significance as conformity to pattern when we attempt to fit the phenomena of petrology into the broad framework of geology. (Turner, 1970)

Accepting this caveat, for the purpose of this paper, I classify metamorphic belts into three types.

TABLE 1. Peak P-T Conditions and Age of Peak Metamorphism for Granulite (G) Metamorphic Belts

| Age, Ma   | Location                                                                     | P (GPa) <sup>1</sup> | T (°C) <sup>1</sup> | Original references                                                                                                                                             |
|-----------|------------------------------------------------------------------------------|----------------------|---------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|
| ~2700     | Minto block, Superior Province, Canada (average 7 samples)                   | 6.7                  | 840                 | Bégin and Pattison, 1994; Pattison and Bégin, 1994                                                                                                              |
| ~2700     | English River domain, Superior Province, Canada (average 7 samples)          | 6.5                  | 810                 | Perkins and Chipera, 1985                                                                                                                                       |
| 2690–2670 | Quetico metasedimentary belt, Superior Province, Canada (average 6 samples)  | 8.0                  | 880                 | Percival and McGrath, 1986                                                                                                                                      |
| 2685–2650 | Ashuanipi Complex, Superior Province, Canada (average 10 samples)            | 7.8                  | 880                 | Percival, 1991                                                                                                                                                  |
| ~2530     | Karnataka, Dharwar craton, India (average 7 samples)                         | 9.3                  | 840                 | Janardhan et al., 1982; Hansen et al., 1987                                                                                                                     |
| ~2530     | Nilgiri Hills, India (average 51 samples)                                    | 9.9                  | 800                 | Raith et al., 1990                                                                                                                                              |
| 2590–2510 | Napier Complex, Enderby Land, Antarctica (average 25 samples)                | 8.9                  | 950                 | Harley, 1985; Harley et al., 1990                                                                                                                               |
| ~2500     | Kasai craton, Zaire (average 6 samples)                                      | 7.2                  | 810                 | Bingen et al., 1988                                                                                                                                             |
| 1950–1930 | Taltson magmatic zone, Canada (average 15 samples)                           | 6.7                  | 930                 | Berman and Bostock, 1997                                                                                                                                        |
| ~1850     | Ketilidian Belt, Greenland (average 9 samples)                               | 5.8                  | 940                 | Dempster et al., 1991                                                                                                                                           |
| ~1100     | Adirondack Highlands, Grenville Province, North America (average 11 samples) | 9.2                  | 820                 | Bohlen and Essene, 1979, 1980                                                                                                                                   |
| 640–630   | Furua Complex, Tanzania (average 16 samples)                                 | 12.6                 | 900                 | Coolen, 1980                                                                                                                                                    |
| 610–550   | Sri Lanka Highlands, Sri Lanka (average 10 samples)                          | 7.7                  | 800                 | Schumacher et al., 1990                                                                                                                                         |
| 530–520   | Kerala Khondalite Belt, India (average 34 samples)                           | 7.9                  | 920                 | Harris et al., 1982; Srikantappa et al., 1985; Chacko et al., 1987, 1996; Hansen et al., 1987; Santosh, 1987; Santosh et al., 1990; Nandakumar and Harley, 2000 |
| ~510      | Prydz Bay, Princes Elizabeth Land, Antarctica (average 16 samples)           | 7.2                  | 890                 | Fitzsimons and Harley, 1994                                                                                                                                     |
| ~510      | Madras Granulites, India (average 8 samples)                                 | 8.5                  | 860                 | Sen and Bhattacharya, 1984                                                                                                                                      |

<sup>1</sup>From Pattison et al., 2003.

Figures 4–6 show plots of metamorphic belts classified by type,<sup>2</sup> based on a judgment of the current best estimate of representative peak P–T

conditions and age associated with these conditions for each metamorphic belt (listed in Tables 1–5). Different thermal gradients are implied by the metamorphic grade evaluated by terrane for each type of metamorphism (Fig. 7); one implication of these different thermal gradients is that each type of metamorphism is likely to have a characteristic tectonic setting with which it was associated.

<sup>2</sup>Granulite–ultrahigh temperature (G-UHT) metamorphism, medium-temperature eclogite–high-pressure granulite (E-HPG) metamorphism, and high-pressure to ultrahigh-pressure (HP-UHP) metamorphism.




FIG. 4. Peak metamorphic P-T conditions of metamorphic belts: granulite-ultrahigh temperature (G-UHT) metamorphism (lighter grey symbols are granulite belts from Table 1 and darker grey symbols are UHT metamorphic belts from Table 2).

### *Granulite and ultrahigh-temperature (G-UHT) metamorphism*

Common granulite-facies metamorphism (Fig. 4) is ubiquitous throughout Earth history, and granulites are represented widely in the Archean rock record (e.g., Harley, 1989; Pattison et al., 2003). On modern Earth, granulites of Cretaceous age occur in the deeper parts of oceanic plateaux (e.g., Gregoire et al., 1994; Shafer et al., 2005) and granulites apparently are being generated today under orogenic plateaux such as Tibet, based on the inference of mica breakdown melting derived from interpretations of multiple geophysical datasets (e.g., Nelson et al., 1996; Unsworth et al., 2005).

In contrast, UHT metamorphism (Fig. 4; metamorphic conditions of  $>900^{\circ}\text{C}$  and from  $\sim 0.5$  to  $\sim 1.5$  GPa) is predominantly a Proterozoic phenomenon, probably associated with crustal aggregation into supercontinents, although lower crustal UHT granulites of Cenozoic age are sampled as metapelitic xenoliths retrieved from Quaternary volcanoes in central Mexico (e.g., Hayob et al., 1989), which may relate to the Mesozoic–Cenozoic eastward migration of the Antilles arc from the Pacific to the Atlantic

and the formation of the Caribbean plate. A common diagnostic mineral assemblage in metaclastic rocks is Al-rich orthopyroxene-sillimanite-quartz, but more exotic assemblages including sapphirine-quartz and spinel-quartz also are characteristic.

Common granulite and UHT metamorphic mineral assemblages imply thermal regimes with high  $dT/dP$  of  $>>750^{\circ}\text{C/GPa}$ ; this is approximately equivalent to the more familiar form of the thermal gradient expressed with reference to depth of  $>>20^{\circ}\text{C/km}$ , assuming a density for the lithostatic load equivalent to average continental crust and no significant tectonic overpressure, both of which are reasonable assumptions in this case. P-T paths may be CW or CCW, but the limited range of pressures for common granulite and UHT metamorphism leads to an element of near-isobaric cooling from peak P-T in many cases, followed by decompression during exhumation.

### *Medium-temperature eclogite–high-pressure granulite (E-HPG) metamorphism*

E-HPG eclogites and granulites (Fig. 5) occur in the Neoproterozoic Belomorian Mobile Belt of the Fennoscandian Shield (Earth's oldest crustal




FIG. 5. Peak metamorphic P-T conditions of metamorphic belts: eclogite–high-pressure granulite (E-HPG) metamorphic belts (data from Table 3).

omphacite-garnet rocks at ~2.72 Ga; Volodichev et al., 2004), in the Proterozoic Orosirian Usagaran Orogen of Tanzania (~2.0 Ga; Collins et al., 2004) and in the Grenvillian belts of the Proto-Atlantic region and in the Paleozoic Caledonian and Variscan belts. E-HPG metamorphism (metamorphic conditions of ~750 to >1000°C and from ~1.5 to >2.5 GPa) is predominantly a Proterozoic–Phanerozoic phenomenon related to subduction and collision. The E-HPG field of metamorphism for rocks of quartz–tholeiitic and high-Al basaltic compositions is characterized by garnet and Ca–Tschermarks clinopyroxene-bearing assemblages, whereas metapelitic and quartzofeldspathic rocks are characterized by garnet–kyanite–K-feldspar assemblages.

These observations translate to an environment for E-HPG metamorphism with thermal gradients of 350–750°C/GPa, which is approximately equivalent to the more familiar form of the thermal gradient expressed with reference to depth of 10–20°C/km, assuming a density for the lithostatic load equivalent to average continental crust and no significant tectonic overpressure—assumptions that may result in an underestimate of the thermal gradient in this

case. The higher pressures of metamorphism and colder thermal regime of E-HPG metamorphism in comparison with common granulite and UHT metamorphism are consistent with subduction of crustal rocks into the mantle. P-T paths are more commonly CW than CCW, with peak T after peak P, which suggests that burial to maximum pressures was followed by thermal relaxation as subduction was blocked by collision and erosion or orogenic collapse facilitated exhumation.

#### *Blueschist (BS) and high-pressure to ultrahigh-pressure (HP-UHP) metamorphism*

BS metamorphic belts appear for the first time in the Neoproterozoic Era and become increasingly common through the Phanerozoic Eon; they record low to very low thermal gradients due to subduction at convergent plate margins during the pre-collisional accretionary phase of the tectonic cycle. BS rocks may be associated with low-temperature eclogites.

Lawsonite and inferred pseudomorphs after lawsonite are reported in low-temperature HP blueschist facies rocks world-wide (e.g., Ballèvre et al., 2003), where glaucophane-lawsonite is a lower-




FIG. 6. Peak metamorphic P-T conditions of metamorphic belts: high pressure–ultrahigh pressure (HP–UHP) metamorphism (lighter grey symbols are lawsonite blueschists and eclogites from Table 4, lines join prograde and peak P-T conditions for some localities, and darker grey symbols are UHP metamorphic belts from Table 5).

temperature association compared to glaucophane-epidote; more rarely lawsonite occurs in low-temperature HP eclogite-facies rocks found in subduction-to-collision orogens and young active convergent margins (Table 4; see also Tsujimori et al., 2006b). Lawsonite is stable over a broad range of very low temperature–high-pressure conditions developed during subduction (Fig. 6), although high water content likely is required to stabilize lawsonite-bearing assemblages and cold, possibly rapid exhumation likely is required to preserve them. The oldest lawsonite eclogites are Ordovician in age (Table 4), which suggests that subduction-zone thermal regimes evolved over a few hundred million years from cold enough to generate ordinary blueschists during the Neoproterozoic to colder still during the early Paleozoic Era. This is confirmed by the recent discovery of Fe–Mg carpholite in supracrustal rocks of the lower unit on Motalafjella, in the Svalbard Caledonides, which is by far the oldest known occurrence of this mineral that is indicative of very low thermal gradients (Agard et al., 2005).

UHP metamorphism (Fig. 6; metamorphic conditions of ~600 to >900°C and from ~2.6 to >10.0 GPa) is predominantly a Phanerozoic phenomenon related to deep subduction of continental crust and its (partial) return to normal crustal depths. Approximately 20 HP–UHP metamorphic belts have been documented within major continental plate collision belts. The record of HP–UHP metamorphism is preserved in pods and slabs of eclogite and garnet peridotite within gneisses (e.g., Cuthbert et al., 2000); where present, ophiolitic eclogites are older than continental crustal eclogites. A few UHP rocks contain minute inclusions of coesite in zircon, garnet, and omphacite, and microdiamond in garnet and zircon (e.g., Chopin, 2003), although the microdiamond in UHP metamorphic rocks most likely was precipitated from a fluid phase rather than being simply recrystallized graphite (Dobrzhi-netskaya et al., 2004, 2005). Recent discoveries of pyroxene exsolution in garnet and coesite exsolution in titanite from some HP–UHP metamorphic belts indicate pressures on the order of 6 GPa (e.g., Chopin, 2003), but the inference of stishovite from




FIG. 7. Metamorphic patterns by belt in relation to thermal gradients and age (filled symbols are belts where peak P-T conditions were recorded at < 488 Ma, grey symbols are belts where peak P-T conditions were recorded at < 850 but > 488 Ma, and open symbols are belts where peak P-T conditions were recorded at > 850 Ma).

at least one locality in the Altyn Tagh indicates pressures >10 GPa (L. Liu et al., 2005) and suggests that under the right circumstances, supracrustal rocks may be returned from depths of approximately 300 km, from around the “point of no return” along the descent into the mantle.

The environment of BS and HP-UHP metamorphism suggests thermal gradients of 150–350°C/GPa; this would be approximately equivalent to 4–10°C/km, assuming a density for the lithostatic load equivalent to average continental crust and no significant tectonic overpressure, but the increases in density due to phase changes with depth means natural gradients will be lower. These very cold thermal gradients require subduction of crustal rocks into the mantle, but the younger granulite-facies overprint that is common in some UHP metamorphic terranes suggests thermal relaxation as subduction was blocked by collision, and evolution of thermal gradients to values closer to Barrovian thermal gradients. However, the extreme pressures attributed to some UHP metamorphic belts may be misleading if tectonic overpressure was significant and if it was dynamically maintained (e.g., Manck-

telow, 1995; Petrini and Podladchikov, 2000; Hartz et al., 2005), although tectonic overpressure remains a contentious issue (e.g., Green, 2005).

### Orogenic Systems and Metamorphism

There are two principal styles of orogenic system on modern Earth (e.g., Wilson and Burke, 1972; Miyashiro, 1973; Murphy and Nance, 1991; Ernst, 2005): (1) accretionary orogenic systems [sometimes called “Pacific-type” (e.g., Matsuda and Uyeda, 1971; Maruyama, 1997) or “Cordilleran-type” (e.g., Coney et al., 1980)], which form during ongoing plate convergence, and dominate in the period after supercontinent break-up and continent dispersal; and (2) collisional orogenic systems (sometimes called “Himalayan-type”), in which an ocean is closed, commonly during continent assembly and formation of supercontinents. Collisional orogenic systems may be superimposed on accretionary orogenic systems (e.g., Liou et al., 2004; herein termed “subduction-to-collision orogenesis”), although some accretionary orogenic systems exist for hundreds of millions of years without disruption

by collision whereas others have the continuity of subduction interrupted—but not necessarily terminated if a new trench is formed outboard of the collider—by minor collision due to subduction of ocean-floor debris, including oceanic plateaus (e.g., Nur and Ben-Avraham, 1989; Cloos, 1993; Mann and Taira, 2004), or more moderate collision if an oceanic arc is involved. These alternatives generate the terminal and non-terminal sutures of Dewey and Kidd (1977; see also Burke et al., 1977), respectively.

Accretionary orogenic systems may be subdivided into two types. In extensional-contractional accretionary orogenic systems, continental growth is related to mantle-derived magmatism in dominantly extensional arc systems (e.g., Collins, 2002). In terrane-accretion orogenic systems, allochthonous, commonly oceanic elements are accreted to continental margins and modified into continental lithosphere in arc systems that involve oblique to highly oblique relative plate motion vectors (e.g., Cowan et al., 1997; Monger, 1997; Johnston, 2001).

In extensional-contractional accretionary orogenic systems, the metamorphic pattern is dominated by low-pressure–high-temperature (LP-HT) metamorphism, with looping or ambiguous (CW or CCW) P-T-t paths. Peak metamorphic mineral growth occurred late in relation to tectonic fabrics, and there is an absence at surface outcrop of granulites (especially of UHT granulites), HP granulites, and UHP metamorphic rocks, although blueschists may occur sporadically early in the orogenic cycle (e.g., Spaggiari et al., 2002a, 2002b). Short-lived compressional-contractional phases of orogenesis probably relate to interruptions in the continuity of subduction by topographic features on the ocean plate, particularly oceanic plateaus (e.g., Nur and Ben-Avraham, 1989; Cloos, 1993; Collins, 2002). Granite magmatism is an extensive accompaniment to metamorphism. The rapid loading and unloading P-T-t paths of some Mesozoic metamorphic belts may reflect intra-arc overthickening of extended arc lithosphere (e.g., Hiroi et al., 1998; Whitney et al., 1999).

In terrane-accretion orogenic systems, coeval “paired” metamorphic belts have been argued to characterize the metamorphic pattern (Miyashiro, 1961, 1973). However, this is a false construct that results from failure to recognize orogen-parallel terrane migration and the limitations of the chronological datasets (Brown, 2002b), even though tectonic contacts between the belts were widely

recognized (e.g., Miyashiro, 1961). Commonly a HP-LT (blueschist-eclogite) metamorphic belt occurs outboard and is separated from a LP-HT (andalusite-sillimanite type) metamorphic belt by a terrane boundary. In some of these systems an important additional feature of the orogenic process is ridge subduction (Brown, 1998a), which is reflected in both the pattern of LP-HT metamorphism and the associated magmatism normal to the orogen, which may have adakitic characteristics (Brown, 1998b). Extensive granite magmatism is common, but thin lithosphere suggests mantle delamination. Granulites may occur at the highest grade of metamorphism in the LP-HT belt, and are implied to occur at depth, but so far UHP metamorphic rocks appear to be rare in the outboard HP-LT belt (but see Ota et al., 2004b).

Collisional orogenic systems, as represented in the Phanerozoic geologic record, generally produce characteristic CW metamorphic P-T-t paths. Due to progress made during the past 20 years, we now know that on the modern Earth, initial collision between a subducting continental plate and an overriding plate may generate extreme UHP metamorphism of subducted continental crust, which may melt and if subducted past the “point of no return” may be transported into the deep mantle (e.g., Irifune et al., 1994; Domanik and Holloway, 2000; Searle et al., 2001; Dobrzhenetskaya and Green, 2005; L. Liu et al., 2005; Hacker, 2006). Subduction of continental crust inhibits generation of calc-alkaline magmas, because the high-pressure stability of micas in this thermal regime inhibits liberation of the volatile phase necessary to flux melting (e.g., Ernst, 2005). It is not yet clear whether it is this factor that enables preservation and exhumation of UHP metamorphic rocks and their return back to normal crustal depths. The complementary thermal environment to the cold subduction zone environment is the backarc, and Hyndman et al. (2005a) have argued that because subduction zone backarcs have hot, thin, and consequently weak lithosphere over a considerable distance, these are likely sites of crustal thickening and orogenesis during continental collision. Collision could generate CW or CCW metamorphic P-T-t paths, according to the relative rates of thickening and heat transfer.

An important question relates to the styles of orogenesis earlier in Earth history. Next I review metamorphic conditions of ancient orogenic systems. The pattern of crustal metamorphism in the geological record indicates secular change, and




FIG. 8. Plot of thermal gradients versus age for the three main types of extreme metamorphic belt. Symbols: circles = G-UHT metamorphism; diamonds = E-HPG metamorphism; squares = HP-UHP metamorphism; data from Tables 1–5.

records two major discontinuous modifications in the thermal structure of orogenic systems that separate three tectonic regimes in Earth history. However, I avoid the sterile semantics of what constitutes plate tectonics—it is possible that some form of thermal boundary layer tectonics has operated throughout most of Earth history (Davies, 2006).

**Metamorphic Regimes Through Time**

Change in the metamorphic record broadly coincides with the change from the Archean to Proterozoic and Proterozoic to Phanerozoic Eons (Fig. 8).

*Ultrahigh-temperature metamorphism*

UHT metamorphism has been recognized during at least four distinct periods in Earth history (see Table 2 and references therein): (1) the Neoarchean (e.g., Kaapvaal craton, southern Africa, at ~2.72 Ga; and the Napier Complex, East Antarctica and the Andriamena Unit, Madagascar, at ~2.50–2.46 Ga); (2) the Orosirian (e.g., the Hoggar, northern Africa, and the Lewisian Complex, Scotland, at ~2.0–1.88 Ga); (3) the Ectasian to Stenian (e.g., the Eastern

Ghats Belt, India and the Mollendo-Camana Block, Peru, at ~1.4–1.0 Ga); and (4) the Ediacaran to Early Cambrian (e.g., the Brazilian and Pan-African belts of the southern continents at ~630–510 Ma). There is only one confirmed example of UHT metamorphism >>2.5Ga in age, which is the Voronezh Crystalline Massif of Sarmetia (~3.2 Ga in age). UHT metamorphism is rare in the post-Cambrian rock record. In the Phanerozoic Eon, parts of the European Variscides evolve from E-HPG metamorphism to UHT metamorphism, but there is only one confirmed example of UHT metamorphism younger than Paleozoic, which is the Gruf Unit in the central Alps (~33 Ma). The four main periods of UHT metamorphism show a remarkable coincidence with the first formation of supercratons (Superior/Scavia) and then the supercontinent cycle (Nuna, Rodinia, and Gondwana).

*Medium-temperature eclogite–high-pressure granulite metamorphism*

There are about 50 documented terranes of this type, with approximately equal numbers of Proterozoic and Phanerozoic age, although the Phanerozoic examples are dominated by various segments of the

TABLE 2. Peak P-T Conditions and Age of Peak Metamorphism for Ultrahigh-Temperature (UHT) Metamorphic Belts

| Age (Ma)   | Location                                                               | P (GPa) | T (°C)    | References                                                                                                                                                            |
|------------|------------------------------------------------------------------------|---------|-----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 3178 ± 38  | Voronezh Crystalline Massif, Sarmatia                                  | 1.0–1.1 | 1,000     | Fonarev et al., 2006                                                                                                                                                  |
| 2720–2,715 | Kaapvaal craton, Southern Africa                                       | 0.9–1.1 | 1000–1100 | Dawson and Smith, 1987; Dawson et al., 1997; Schmitz and Bowring, 2003                                                                                                |
| ~2700      | Sutam Block, Central Aldan Superterrane, Russia                        | 1.05    | 1070      | Perchuk et al., 1985; Sal'nikova et al., 2004                                                                                                                         |
| ~2690      | Southern marginal zone, Limpopo Belt, southern Africa                  | 0.9     | 950       | Kreissig, et al., 2000; Tsunogae et al., 2004                                                                                                                         |
| ~2680      | Sipiwesk Lake, Pikwitonei Granulite domain, NW Superior craton, Canada | 1.0–1.1 | 1060–1100 | Arima and Barnett, 1984; Böhm, 2004                                                                                                                                   |
| ~2675      | Layered gneiss unit, Teton–Wind River domain, Wyoming, USA             | >1.1    | >950      | Frost et al., 2005; Hughes et al., 2005                                                                                                                               |
| 2590–2510  | Napier Complex, eastern Antarctica                                     | 0.9–1.1 | 1050–1100 | Ellis et al., 1980; Harley, 1985; Harley and Motoyoshi, 2000; Hokada, 2001; Hokada and Harley, 2004; Hokada et al., 2004; Kelly and Harley, 2005; Suzuki et al., 2006 |
| ~2500      | Andriamena Mafic Unit, north-central Madagascar                        | 1.05    | 1000      | Goncalves et al., 2004; Paquette et al., 2004                                                                                                                         |
| ~2480      | Badcall Bay, Assynt terrane, Lewisian Complex, Scotland                | 1.4     | 1000      | Cartwright and Barnicoat, 1989; Love et al., 2004                                                                                                                     |
| 2090–2040  | Bhandara-Balaghat granulite domain, Central Indian tectonic zone       | 0.9     | 950       | Bhowmik et al., 2005                                                                                                                                                  |
| ~2070      | Jequié Block, Bahia, Brazil                                            | 0.5–0.7 | 1000–1100 | Barbosa et al., 2004                                                                                                                                                  |
| 2027 ± 6   | Central Zone, Limpopo Belt, southern Africa                            | 1.0     | 1000      | Harris and Holland, 1984; Jaekel et al., 1997; Droop, 1989; Hisada and Miyano, 1996; Kröner et al., 1999                                                              |
| ~2000      | InOuzzal Terrain, NW Hoggar, Algeria                                   | 1.0–1.1 | 1000–1050 | Kienast and Ouzegane, 1987; Bertrand et al., 1992; Ouzegane and Boumaza, 1996; Ouzegane et al., 2003                                                                  |
| 1950–1930  | Taltson magmatic zone, northwestern Canada                             | 0.6–0.8 | >962      | Farquhar et al., 1996                                                                                                                                                 |
| ~1880      | South Harris, Lewisian Complex, Scotland                               | 1.0     | 955       | Baba, 1998, 1999, 2003; Friend and Kinny, 2001; Mason et al., 2004; Hollis et al., 2006                                                                               |
| ~1650      | Wilson Lake, Grenville Province, Canada                                | 1.0–1.1 | 1060–1100 | Morse and Talley, 1971; Leong and Moore, 1972; Arima et al., 1986; James et al., 2002; Korhonen and Stout, 2004                                                       |
| 1525–1510  | Epembe Unit, NW Namibia                                                | 0.95    | 970       | Brandt et al., 2003; Seth et al., 2003                                                                                                                                |
| ~1100      | Eastern Ghats Province, India                                          | 0.9–1.0 | 985–1085  | Dasgupta et al., 1992, 1994; Jarick, 2000; Bhattacharya and Kar, 2002; Sengupta et al., 2004; Bhattacharya, 2004                                                      |

*Table continues*

TABLE 2. *Continued*

| Age (Ma) | Location                                                                  | P (GPa) | T (°C)    | References                                                                                                    |
|----------|---------------------------------------------------------------------------|---------|-----------|---------------------------------------------------------------------------------------------------------------|
| 1200–940 | Mollendo-Camana Block, Arequipa-Antofalla Basement, central Andean Margin | 1.2     | 1000      | Martingole and Martelat, 2003; Loewy et al., 2004                                                             |
| ~770     | Niquelândia and Barro Alto complexes, Northern Brasília Belt, Brazil      | 0.8     | 980       | Ferreira Filho et al., 1994, 1998; Moraes and Fuck, 2000; Pimentel et al., 2004                               |
| 645–635  | Anápolis-Itaçu complex, Southern Brasília Belt, Brazil                    | 0.875   | 980       | Moraes et al., 2002; Piuzeana et al., 2003; Baldwin et al., 2005, unpubl. data                                |
| ~635     | Labwor Hills, Uganda                                                      | 0.95    | 1075      | Leggo et al., 1971; Sandiford et al., 1987                                                                    |
| 610–550  | Central Highland Complex, Sri Lanka                                       | 1.1–1.2 | 1000–1100 | Kriegsman and Schumacher, 1999; Osanai et al., 2003; Sajeev and Osanai, 2004                                  |
| 550–530  | Southern granulite terrain, India                                         | 1.1     | 980–1030  | Mohan et al., 1986; Brown and Raith, 1996; Raith et al., 1997                                                 |
| 530–520  | Kerala Khondalite Belt, India                                             | 1.0     | 950       | Chacko et al., 1996; Pattison et al., 2003; Morimoto et al., 2004; Santosh et al., 2005; Shabeer et al., 2005 |
| 517 ± 9  | Lutzow-Holm Bay, East Antarctica                                          | 1.1     | 1000      | Motoyoshi and Ishikawa, 1997; Frazier et al., 2000; Kawakami and Motoyoshi, 2004                              |
| 511 ± 4  | Prydz Bay localities, East Antarctica                                     | 1.1–1.2 | 1000–1050 | Harley, 1998; Kelsey et al., 2003a, 2003b                                                                     |
| ~510     | Madurai Block, Southern India                                             | 1.0     | 1000      | Mohan et al., 1986; Brown and Raith, 1996; Raith et al., 1997; Tamashiro et al., 2004                         |
| ~254     | Kontum Massif, Vietnam                                                    | 1.1     | 1000      | Nam et al., 2001; Nakano et al., 2004; Osanai et al., 2004                                                    |
| ~33      | Gruf Unit, central Alps                                                   | 1.1     | 980–1030  | Droop and Bucher-Nurminen, 1984; Liati and Gebauer, 2003                                                      |

European Variscides (see Table 3 and references therein). About one-third of recognized high-pressure granulite belts have evolved from medium-temperature eclogite-facies assemblages, and about one-quarter of the remainder evolve into UHT metamorphic conditions, both apparently without distinction between Proterozoic and Phanerozoic examples. Compositional zoning of minerals and reaction coronas, symplectites, and kelyphites are common as multivariant reactions are crossed during this evolution. Three periods of E-HPG metamorphism coincide with the first formation of supercratons (Superior/Scandia) and then the Nuna and Rodinia supercontinent cycles, but unlike UHT metamorphism *sensu stricto*, E-HPG metamorphism is characteristic not just of Gondwana assembly but also of the Lower Paleozoic Caledonides and

Variscides. E-HPG metamorphism is rare in the post-Carboniferous rock record.

#### *Blueschist (BS) and high-pressure to ultrahigh-pressure metamorphism*

BS belts appear for the first time in the Neoproterozoic Era, where two definite examples, the Aksu Group, western China, and inliers of the Anti-Atlas belt in central Morocco, both are overlain by sediments of Ediacaran age (Maruyama et al., 1996). Other possible Neoproterozoic blueschists are those in the suture between the Yangtze and Cathaysia blocks of south China, within the Delhi Fold Belt of Rajasthan, India, and in Anglesey, Wales. BS rocks may be associated with low-temperature eclogites (Maruyama et al., 1996).

TABLE 3. Peak P-T Conditions and Age of Peak Metamorphism for Medium-Temperature Eclogite-High-Pressure Granulite (E-HPG) Metamorphic Belts

| Age (Ma)     | Location                                                                                          | P (Gpa)   | T (°C)    | References                                                                                           |
|--------------|---------------------------------------------------------------------------------------------------|-----------|-----------|------------------------------------------------------------------------------------------------------|
| 2720 ± 8     | Gridino eclogite-bearing mélange zone, Belomorian mobile belt, Fennoscandian Shield               | 1.75      | 770–880   | Volodichev et al., 2004                                                                              |
| ~2487        | Jianping Complex, North China craton                                                              | 1.1       | 800       | Kröner et al., 1998a; Wei et al., 2001                                                               |
| 2700–2400    | S Badakhshan Block, Afghanistan                                                                   | 1.2–1.3   | 700–750   | Faryad, 1999                                                                                         |
| ~2400        | Sharyzhgai Complex, SW margin, Siberian craton                                                    | 2.3–3.0   | 920–1030  | Ota et al., 2004a                                                                                    |
| 1999.1 ± 1.1 | Usagaran orogen, Tanzania                                                                         | 1.8       | 750       | Möller et al., 1995; Collins et al., 2004                                                            |
| 1848 ± 6     | Ubendian, Tanzania                                                                                | 1.2–1.4   | 840–900   | Sklyarov et al., 1998; Boven et al., 1999; Muhongo et al., 2002                                      |
| 1904.0 ± 0.3 | Snowbird tectonic zone, Canadian Shield                                                           | 1.8–2.0   | 920–1,000 | Baldwin et al., 2003, 2004                                                                           |
| 1900–1,850   | Sanggan area, North China craton                                                                  | 1.1–1.5   | 750–870   | Guo et al., 2002                                                                                     |
| 1900–1800    | Hengshan, North China Craton                                                                      | 1.35–1.55 | 770–840   | Zhao et al., 2001; O'Brien et al., 2005                                                              |
| ~1790        | Rudall complex, central Western Australia                                                         | 1.2       | 800       | Smithies and Bagas, 1997; Bagas, 2004                                                                |
| 1134–1129    | Llano uplift, Texas                                                                               | 1.4–2.4   | 610–775   | Carlson et al., 2007                                                                                 |
| 1050–1020    | Manicouagan imbricate zone, and Molsen Lake, Gagnon and Lelekuau terranes, Grenville belt, Canada | 1.5–2.0   | 800–900   | Indares, 1993, 1995, 1997, 2003; Indares and Dunning, 2001                                           |
| 1100–960     | Zambezi belt, Zimbabwe                                                                            | 1.9       | 760       | Dirks and Sithole, 1999                                                                              |
| ~975         | Sveconorwegian belt, SW Sweden                                                                    | 1.0–1.2   | 705–795   | Möller, 1998, 1999                                                                                   |
| ~630         | Guaxupé                                                                                           | 1.4–1.5   | 1040      | Campos Neto and Caby, 1999, 2000; Del Lama et al., 2000; Moraes et al., 2003; Valeriano et al., 2004 |
| ~630         | Varghina nappe complex, Brasilia orogen, SE Brazil                                                | 1.75      | 950       | Campos Neto and Caby, 1999, 2000; Del Lama et al., 2000; Moraes et al., 2003; Valeriano et al., 2004 |
| 610 ± 2      | Dahomeyides, Ghana-Togo                                                                           | 1.4       | 800–900   | Attoh et al., 1997; Attoh, 1998a, 1998b;                                                             |
| ~605         | NE Ox Inlier, NW Ireland                                                                          | 1.4–1.5   | 850–900   | Sanders et al., 1987                                                                                 |
| 560–520      | Musgrave Block, Australia                                                                         | 1.2–1.3   | 700–750   | Scrimgeour and Close, 1999                                                                           |
| ~530         | Lufilian Arc-Zambesi Belt, Zambia                                                                 | 1.0–1.3   | 700–750   | John et al., 2004                                                                                    |
| ~500         | South Altyn Tagh, NW China                                                                        | 1.9–2.7   | 870–1050  | J. X. Zhang et al., 2005                                                                             |
| ~470         | Carolina terrane, Southern Appalachians                                                           | 1.2–1.5   | 700–800   | Shervais et al., 2003                                                                                |
| 470–460      | Moine, Scotland                                                                                   | 1.2       | 700       | Friend et al., 2000                                                                                  |
| 432 ± 6      | Roan, Western Gneiss Region, Norway                                                               | 1.45      | 870       | Johansson and Möller, 1986; Möller, 1988; Dallmeyer et al., 1992a                                    |
| 410–390      | Payer Land, NE Greenland                                                                          | 1.4–1.7   | 800–850   | Elvevold and Gilotti, 2000; Gilotti and Elvevold, 2002; Elvevold et al., 2003                        |
| 440–390      | Bragança-Morais, NE Portugal                                                                      | 1.3–1.4   | 800–850   | Marques et al., 1996                                                                                 |

*Table continues*

TABLE 3. *Continued*

| Age (Ma)    | Location                         | P (Gpa) | T (°C)    | References                                                                                                                                   |
|-------------|----------------------------------|---------|-----------|----------------------------------------------------------------------------------------------------------------------------------------------|
| 402 ± 1     | Góry Sowie, Sudetes, Poland      | 1.5–2.0 | 900–1000  | Kryza et al., 1996; O'Brien, 1997                                                                                                            |
| 395–380     | Munchberg Massif, Germany        | 1.8     | 925       | Gayk et al., 1995                                                                                                                            |
| 370–360     | Armorican Massif, France         | 2.0     | 950       | Ballèvre et al., 1994; Bosse et al., 2000; Le Hébel, 2002                                                                                    |
| 369–360     | Sniecznik, Sudetes, Poland       | 2.1–2.8 | 800–1,000 | Pouba et al., 1985; Bakun-Czubarow, 1992; Kryza et al., 1996; Klemm and Bröcker, 1999                                                        |
| 350–340     | Massif Central, France           | 1.0–1.2 | 700–750   | Pin and Vielzeuf, 1983, 1988; Dufour et al., 1985; Gardien et al., 1990; Costa et al., 1993                                                  |
| 341.5 ± 0.8 | Granulitgebirge, Saxony, Germany | 2.2     | 1010–1060 | Rötzler, 1992; von Quadt, 1993; Kröner et al., 1998b; Rötzler and Romer, 2001                                                                |
| ~341        | Erzgebirge, Germany              | 2.1     | 830       | Willner et al., 1997; Kröner and Willner, 1998; Nasdala and Massonne, 2000                                                                   |
| 342 ± 5     | N. Bohemia, Czech Republic       | 1.5–1.7 | 800–850   | Kotková et al., 1996                                                                                                                         |
| 339.8 ± 2.6 | S. Bohemia, Czech Republic       | 1.5     | 900       | Fiala et al., 1987; Vrána, 1989; O'Brien, 1999; Kröner et al., 2000                                                                          |
| ~341        | Moravia, Czech Republic          | 1.4–2.0 | 750–1050  | Medaris et al., 1995a, 1995b                                                                                                                 |
| ~340        | Lower Austria                    | 1.4–1.6 | 950–1050  | Scharbert and Carswell, 1983; Dallmeyer et al., 1992b; Carswell and O'Brien, 1993; Becker, 1997; Cooke et al., 2000; Cooke and O'Brien, 2001 |
| 340–335     | Schwarzwald, Germany             | 1.5     | 950–1000  | Kalt et al., 2000                                                                                                                            |
| ~330        | Uiten zone, Eastern Alps         | 1.45    | 700       | Godard et al., 1996; Tumati et al., 2003                                                                                                     |
| ~330        | Santa Lucia, Corsica             | 1.2–1.5 | 800       | Pin and Vielzeuf, 1983, 1988; Libourel, 1988                                                                                                 |
| 125–110     | Fjordland, New Zealand           | 1.4     | 750–800   | Ireland and Gibson, 1998; Clarke et al., 2000                                                                                                |
| ~100        | Jijal Complex, Pakistan          | 1.5–1.6 | 800–900   | Yamamoto, 1993; Jan and Karim, 1995; Yamamoto and Yoshino, 1998; Yoshino et al., 1998                                                        |

HP-UHP metamorphism has been recognized during multiple periods in Earth history (see Table 5 and references therein): the Edicaran (e.g., Minas Gerais (?), Brazil; Mali, western Africa); the Cambrian (e.g., Kokchetav Massif, northern Kazakhstan; southwest Altyn Tagh, west China); the Ordovician–Silurian (e.g., Tromsø Nappe, Norway; Dulan Belt, North Qaidam, west China); the Devonian (e.g., Western Gneiss Region, Norway, northeastern Greenland; Maksyutov Complex, southern Urals; Saxonian Erzgebirge, Germany; Bohemian Massif,

central Europe; and the French Massif Central); the Permo-Triassic (e.g., Dabie-Sulu belt, east-central China; western Tianshan (?), west China); the Early Cretaceous (e.g., central Indonesia, southwestern Sulawesi); through the Cenozoic Era (e.g., Dora Maira Massif, Italy; Zermatt-Saas Zone, Western Alps; Tso-Morari Complex, India; Kaghan Valley, Pakistan); and finally the Earth's youngest UHP rocks of Miocene–Pliocene age (Fergusson Island, eastern Papua New Guinea). P-T-t paths from UHP metamorphic terranes vary from hairpins, in which

TABLE 4. Peak P-T Conditions and Age of Peak Metamorphism for Lawsonite Blueschists and Lawsonite Eclogites

| Age (Ma) | Location                                            | P (Gpa)            | T (°C)             | References                                                                          |
|----------|-----------------------------------------------------|--------------------|--------------------|-------------------------------------------------------------------------------------|
| ~475     | Motalafjella, Spitsbergen                           | 1.5–1.6<br>1.8–2.4 | 380–400<br>580–640 | Agard et al., 2005<br>Hirajima et al., 1988; Dallmeyer et al., 1989                 |
| ~470     | Port Macquarie, New South Wales, Australia          | >1.8–<2.6          | 560                | Fukui et al., 1995; Och et al., 2003                                                |
| ~400     | Western Gneiss Region, Norway                       | 2.9                | 710                | Ravna, 1982; Carswell et al., 2003; Root et al., 2004                               |
| ~370     | Ile de Groix, Armorican Massif, France              | 1.8–2.0            | 450                | Ballèvre et al., 2003; Bosse et al., 2005                                           |
| ~350     | Western Tianshan, NW China                          | 1.5–1.9            | 540–580            | Klemd et al., 2002, 2005; Wei et al. 2003                                           |
| ~340     | Maksyutovo Complex, Southern Urals, Russia          | 1.8–2.1            | 520–540            | Schulte and Sindern, 2002                                                           |
| ~240     | Huangzhen-Zhujiachong, South Dabie terrane, China   | 3.3                | 670                | Castelli et al., 1998; Li et al., 2004                                              |
| ~230     | Qinglongshan, Sulu Terrane, China                   | 3.0–3.5            | 600–700            | Mattinson et al., 2004                                                              |
| ~218     | Pinchi Lake, British Columbia, Canada               | 1.3<br>2.4–2.7     | 500<br>450         | Ghent et al. (1993)<br>Ghent, pers. commun., October 10, 2006                       |
| ~145     | Cazadero, Northern California, USA                  | 1.0–1.3<br>1.8–2.2 | 360–400<br>430–460 | Shibakusa and Mackawa, 1997;<br>Anczkiewicz et al., 2004<br>Tsujimori et al., 2006b |
| ~145     | Jenner, Northern California, USA                    | 0.9<br>2.2         | 320<br>435         | Ravna et al. 1994; Anczkiewicz et al., 2004<br>Ravna and Terry, 2004                |
| ~132     | Motagua fault zone, Guatemala                       | 2.6                | 480                | Brueckner et al., 2005; Tsujimori et al., 2006a                                     |
| ~120     | Bantimala, Sulawesi, Indonesia                      | 2.1                | 520                | Miyazaki et al., 1996; Parkinson et al., 1998; Tsujimori et al., 2006b              |
| ~95      | Sistan suture zone, Iran                            | 2.25               | 600                | Fotoohi Rad et al., 2005                                                            |
| ~87      | Orhaneli, NW Turkey                                 | 2.1–2.7            | 430                | Okay et al., 1998; Okay, 2002                                                       |
| ~85      | Corsica, France                                     | 2.0                | 390                | Caron and Péquignot, 1986; Tsujimori et al., 2006b                                  |
| ~80      | Samaná Peninsula, Dominican Republic                | 1.6                | 360                | Goncalves et al., 2000; Zack et al., 2004                                           |
| ~80      | Garnet Ridge, Colorado Plateau, USA                 | 3.5                | 540                | Usui et al., 2003; Tsujimori et al., 2006b                                          |
| ~50      | Sifnos, Greece                                      | 2.0                | 550–600            | Schmädicke and Will, 2003; Forster and Lister, 2005                                 |
| ~44      | Diahot terrane, Pam Peninsula, New Caledonia        | 1.5                | 410                | Spandler et al., 2005; Potel et al., 2006                                           |
|          | (metamorphic peak)                                  | 2.1–2.4            | 475–580            | Fitzherbert et al., 2003, 2005                                                      |
| ~32      | Eclogite zone, Tauern Window, Eastern Alps, Austria | 2.2                | 600                | Hoschek, 2001, 2004; Kühn et al., 2005                                              |

TABLE 5. Peak P-T Conditions and Age of Peak Metamorphism for Granulite and Ultrahigh-Pressure (UHP) Metamorphic Belts

| Age (Ma)    | Location                                                         | P (Gpa)    | T (°C)    | References                                                                                     |
|-------------|------------------------------------------------------------------|------------|-----------|------------------------------------------------------------------------------------------------|
| ~630        | Pan-African nappes of Minas Gerais State, SE Brazil              | >2.7       | 800       | Parkinson et al., 2001                                                                         |
| ~620        | Pan-African nappes of Northern Mali, West Africa                 | >2.7       | 700–750   | Caby, 1994; Jahn et al. 2001                                                                   |
| 530–528     | Kumdy-Kol, Kokchetav Massif (diamond-grade), northern Kazakhstan | 4.0–6.0    | 920–1000  | Sobolev and Shatsky, 1990; Shatsky et al., 1995; Rubatto et al., 2005;                         |
| 508–500     | Kulet, Kokchetav Massif (coesite-grade), northern Kazakhstan     | 3.8–4.0    | 730       | Sobolev and Shatsky, 1990; Shatsky et al., 1995; Rubatto et al., 2005;                         |
| 503.9 ± 5.3 | Jianggalesayi, southwest Altyn Tagh, west China                  | 2.8–3.0    | 820–850   | L. Liu et al., 2001, 2003b; J. X. Zhang et al., 2001                                           |
| 498–490     | Lantermann Range, Northern Victoria Land, Antarctica             | >2.7       | 750       | Ghiribelli et al., 2002; Palmeri et al., 2003                                                  |
| 452 ± 1.7   | Tromsø Nappe, Norway                                             | 3.4        | 735       | Corfu et al., 2003; Ravna and Roux, 2006                                                       |
| 440–420     | Dulan Belt, North Qaidam                                         | 2.9–3.3    | 630–740   | Song et al., 2003; Mattinson et al., 2005a, 2000b                                              |
| 420–400     | Massif Central, France                                           | >2.8       | 740–780   | Lardeaux et al., 2001                                                                          |
| ~400        | Western Gneiss Region, Norwegian Caledonides                     | >2.8       | 790       | Smith, 1984, 1988; Wain, 1997; Cuthbert et al., 2000; Ravna and Terry, 2004; Root et al., 2004 |
| 389 ± 4     | Maksyutov Complex, southern Urals                                | 3.2        | 650       | Chesnokov and Popov, 1965; Leech et al., 2002; Bostick et al., 2003                            |
| 361–347     | Northeast Greenland, Greenland Caledonides                       | 3.6        | 970       | Gilotti and Ravna, 2002; McClelland et al., 2005, 2006                                         |
| 360 ± 7     | Erzgebirge, Saxony, Germany                                      | >4.2       | 900–1000  | Massonne, 1999, 2001; Nasdala and Massonne, 2000                                               |
| 360–340     | Leaota Massif, South Carpathians                                 | >2.7       | 700–750   | Săbău, 2000; Medaris et al., 2003                                                              |
| 355–340     | Sudetes Mountains, SW Poland                                     | 3.0        | 700–800   | Bakun-Czubarow, 1991, 1992                                                                     |
| 355–340     | Bohemian Massif, Czech Republic                                  | 4.5–4.9    | 1050–1150 | Nakamura et al., 2004                                                                          |
| ~345        | Makbal, Kyrgyzstan                                               | 2.6        | >610–680  | Tagiri et al., 1995                                                                            |
| ~345        | At-bashy, Kyrgyzstan (younger?)                                  | >2.5       | 660       | Tagiri et al., 1995                                                                            |
| ~345*       | Western Tianshan, China                                          | >4.95–5.07 | 560–610   | L. Zhang et al., 2002, 2003, 2005a, 2005b, 2005c; Gao and Klemd (2003)                         |
| ~240        | Dabie Shan, Central China                                        | 4.0        | 700       | Wang et al., 1989, 1995; Wang and Liou, 1991; Okay, 1993; Ravna and Terry, 2004                |
| ~230        | Su-Lu Terrane, Eastern China                                     | 3.5        | 660       | R. Y. Zhang et al., 1995; L. Liu et al., 2003a; F. Liu et al., 2004a, 2004b                    |
| *233–225    | Western Tianshan, China                                          | >4.95–5.07 | 560–610   | L. Zhang et al., 2002, 2003, 2005a, 2005b                                                      |
| 120–115     | Bantimala Complex, Central Sulawesi, Indonesia (eclogite)        | >2.7       | 720–750   | Parkinson et al. (1998)                                                                        |

*Table continues*

TABLE 5. *Continued*

| Age (Ma)   | Location                                   | P (Gpa) | T (°C)  | References                                                                                  |
|------------|--------------------------------------------|---------|---------|---------------------------------------------------------------------------------------------|
| ~90        | Pohorje Mountains, Eastern Alps, Slovenia  | 3.0–3.1 | 760–825 | Janák et al., 2004; Thöni, 2005                                                             |
| 73.5 ± 3.4 | Kimi, East Rhodope, northern Greece        | >2.5    | 700–780 | Mposkos and Kostopoulos, 2001; Liati et al., 2002                                           |
| 53.5 ± 7   | Tso-Morari Complex, Indian Himalaya        | >2.8    | >640    | Mukherjee and Sachan, 2001; Mukherjee et al., 2003; Sachan et al., 2004; Leech et al., 2005 |
| 52 ± 18    | Zermatt-Saas Zone, Western Alps            | 2.6–3.0 | 590–630 | Reinecke, 1991; van der Klauw et al., 1997                                                  |
| 46.3 ± 0.2 | Kaghan Valley, Pakistan Himalaya           | 2.7–3.2 | 700–770 | O'Brien et al., 2001; Kaneko et al., 2003; Parrish et al., 2003                             |
| 35.4 ± 1   | Dora-Maira Massif, Western Alps            | 4.35    | 740     | Chopin, 1984; Chopin et al., 1991; Hermann, 2003                                            |
| 8–3        | Fergusson Island, eastern Papua New Guinea | 2.75    | 700     | S. L. Baldwin et al., 2005                                                                  |

the exhumation path is close to the burial path in terms of  $dT/dP$ , and in the range 150–350°C/GPa (4–10°C/km), implying syn-subduction exhumation, to CW paths with isentropic decompression segments to Barrovian-type thermal regimes of 350–750°C/GPa (10–20°C/km), consistent with thermal relaxation after initial fast exhumation to Moho depths.

What is the depth limit after which continental crust is irretrievably “lost” to the mantle? The mineral assemblages of common continental crustal rocks at pressures around 10 GPa—based on experimental studies and assemblages calculated using internally consistent thermodynamic datasets (L. Liu et al., 2005; Massonne and Brandelik, 2005)—suggest that continental crust will become denser than ambient mantle at a depth of ~300 km, inhibiting its buoyant return to a more normal depth. Part of the Altyn Tagh in western China may have been returned from pressures that were this extreme—around the “point of no return” (L. Liu et al., 2005).

Is there any evidence of crustal recycling into the deep mantle in the Proterozoic or Archean Eons? Two studies show enigmatic chemistry of diamond that may record early UHP metamorphism. First, microdiamonds from the 1,800 Ma Akluilak minette dikes (Nunavut, Canada) are similar to diamonds formed in subducted metamorphic rocks (Cartigny et al., 2004)—they have high concentrations of

un-aggregated nitrogen and positive  $\delta^{15}\text{N}$ —suggesting that they formed within supracrustal rocks taken down to ultrahigh pressures before being sampled by the minette magma. The postulated UHP metamorphism may have been related to either the Thelon or Trans-Hudson orogenic system and, inasmuch as a cold environment is required, it may indicate that something akin to modern subduction was active during the Paleoproterozoic Era on some parts of the Earth. Second, populations of sulfide inclusions in Archean diamonds from the Orapa kimberlite pipe in the Kaapvaal-Zimbabwe craton, Botswana, preserve mass-independent sulfur isotope fractionations (Farquhar et al., 2002). The data indicate that material was transferred from the atmosphere to the mantle by sedimentation, lithification, metamorphism, and convective stirring, but imply that sulfur was not well mixed in the diamond source regions. Evidence for these connections is preserved because sulfide was encased in diamond and transported back to the surface, but it is unclear whether these connections require a process akin to modern subduction.

#### *Archean metamorphism*

The Eoarchean through Mesoarchean rock record, and particularly the greenstone belts, generally records P-T conditions characteristic of low-to-moderate-P–moderate-to-high-T metamorphic

facies [e.g., the ~3,800 Ma metamorphism recorded by the Isua Supracrustal Belt of southern West Greenland (Komiya et al., 2002) and the ~3,230 Ma metamorphism recorded in the Barberton and related greenstone belts of South Africa (Dziggel et al., 2002, Diener et al., 2005)]. Thermal gradients were around 750°C/GPa (~20°C/km)—consistent with generally thinner lithosphere during the Archean. Although ordinary granulites are common throughout the Archean Eon, extreme thermal conditions are not recorded before the Neoarchean. Furthermore, blueschist-facies rocks are not documented, and there is no metamorphic imprint of subduction of continental crust into the mantle. However, the chemistry of Mesoarchean to Neoarchean eclogite xenoliths and the chemistry and paragenesis of Neoarchean diamonds in kimberlites within cratons suggest some process associated with supercraton formation at convergent margins was operating to take basalt and other supracrustal material into the mantle by the Neoarchean Era (e.g. Jacob, 2004; Shirey et al., 2004).

An unusual feature of Archean granulite-amphibolite facies terranes is that the granulite-amphibolite facies isograd commonly is a retrograde feature (e.g., McGregor and Friend, 1997). This requires the widespread availability of an H<sub>2</sub>O-rich fluid phase (volatile phase and/or melt phase) during the waning stages of the metamorphic evolution, and together with the suggestion that low-temperature water-saturated melting may be common early in Earth history (Watson and Harrison, 2005), suggests a tectonic setting and/or process particular to this Eon.

### *Inferences and implications*

Overall, the restricted time span of different types of metamorphism through Earth history and the periods of metamorphic quiescence during the Proterozoic Eon (Fig. 8) suggest: (1) a link with major events in the mantle, supporting the postulate of discontinuous geodynamic evolution of the Earth; (2) a different style of tectonics in the Archean Eon in comparison with the Proterozoic Eon; and (3) a different style of tectonics in the Proterozoic Eon in comparison with the Phanerozoic Eon.

The metamorphic record indicates a dramatic change in the thermal environments of crustal metamorphism through the Neoarchean to the Archean-to-Proterozoic transition, which may reflect the switch to a “Proterozoic plate tectonics regime.”

Such a premise is consistent with the first occurrence of ophiolite-like complexes in Proterozoic suture zones (Moore, 2002) and with the increase in  $\delta^{18}\text{O}$  of magmas through the Paleoproterozoic, which Valley et al. (2005) argue may reflect maturation of the crust, the beginning of recycling of supracrustal rocks, and their increasing involvement in magma genesis via subduction. This is also consistent with assembly of the continental crust into progressively larger units to form supercratons (Bleeker, 2003), perhaps indicating a change in the pattern of mantle convection during the transition to the Proterozoic Eon, a period of Earth history that was characterized by the assembly, partial dispersal, and reconfiguration of much of the continental lithosphere into several supercontinents. Although the style of Proterozoic subduction remains cryptic, the change in tectonic regime whereby interactions between discrete lithospheric plates generated tectonic settings with contrasting thermal regimes was a landmark event in Earth history.

The concentration of UHT metamorphic belts in the interval Neoarchean to Cambrian, and their broad correlation in time with amalgamation of continental crust into supercratons and supercontinents, suggests that UHT metamorphism, which commonly records tight clockwise P-T paths, may record closure and thickening of backarcs with their characteristic high heat flow (cf., Currie and Hyndman, 2005; Hyndman et al., 2005b). This certainly appears to be true for the aggregation of the continental crust in Gondwana, where the geological evidence for tectonic setting is more complete. However, it is less clear whether this tectonic setting is appropriate for all UHT metamorphic belts, although it is an attractive solution to the heat problem.

The thermal environment of E-HPG metamorphism [350–750°C/GPa (~10–20°C/km)] and maximum pressures of metamorphism suggest that crustal rocks have been taken to mantle depths, although the pressures could simply reflect collision and doubling in thickness by crustal stacking following subduction of oceanic plates. However, given the contrasting metamorphic record of the Proterozoic versus the Phanerozoic, it is difficult to disagree with Stern (2005) that subduction as we know it on the modern Earth did not commence before the Neoproterozoic Era.

The transition to the “modern plate tectonics regime” through the Neoproterozoic is one in which

ocean crust became thinner whereas the lithosphere became thicker (e.g., Moores, 2002), and the anastomosing pattern of Brasiliano–Pan-African orogens that created Gondwana was replaced by semi-circumferential large-scale plate-margin orogens (e.g., the Terra Australis, Caledonide, and Tethyan orogens). These features may indicate a change to a smaller number of plates, perhaps reflecting a fundamental change in the pattern of mantle convection that led to a sharp decrease in the thermal gradient of subduction-zone metamorphism and facilitated deeper subduction of continental crust during the Phanerozoic (Fig. 7). Many authors have noted the confluence of other changes to the Earth system during this transition (e.g., Maruyama and Liou, 1998, 2005). The repeated occurrence of UHP metamorphism (with the inference of subducted continental crust) since the Ediacaran Period suggests a common process in convergent tectonic settings, and the P-T conditions coupled with high-precision ages indicate fast rates of subduction to generate the low thermal gradient (150–350°C/GPa) and rates of exhumation similar to modern plate velocities (1–2 cm yr<sup>-1</sup>). These features are characteristic of the modern plate tectonics regime.

### Secular Change in Tectonics

Although there are differences in opinion about how a hotter early Earth may have dissipated heat, there is general agreement that the mean mantle temperature must have been higher, crustal heat production must have been higher, the mean thermal boundary layer thickness must have been lower, and the mean residence time of oceanic lithosphere in the thermal boundary layer must have been much shorter (e.g., Pollack, 1997). There are multiple consequences for the organization of the TBL and for Archean–Proterozoic tectonics.

On modern Earth, plates and plate tectonics partly represent a surface conductive thermal boundary layer, which may return to the interior of Earth via subduction, but where continents and lithospheric mantle are buoyant, due to chemical composition and, therefore, mineralogy, and do not readily sink just because they are cold. The plate mosaic is likely a self-organized network of plates and forces readily reorganized by changes in stress (Anderson, 2002). Hargraves (1986) argued that younger lithosphere on the early Earth would subduct more slowly, and that if heat flow at about the Archean–Proterozoic transition was three times the

present heat flow, then ridge length in the ocean basins would have to have been as much as 27 times the present ridge length to dissipate the heat. In an alternative calculation, Abbott and Menke (1990) suggested that the length of plate boundaries at 2.4 Ga was about 2.2 times the present boundary lengths, with more than 60 individual plates moving with velocities and associated sea-floor spreading rates about 16% slower than today. More recently, Korenaga (2006) also has argued for more sluggish plate tectonics when the mantle was hotter to allow lithosphere with a thicker depleted mantle component to become negatively buoyant by thermal contraction, but he concluded that consequently the concept of many small plates in the Archean is physically implausible. The point of these examples is that whatever the configuration of ridges and plate boundaries, and whatever the number of plates, tectonics in the Archean was likely to have been different in many of the details in comparison with the modern plate tectonics regime.

The effect of increased radiogenic heat production (RHP) on temperatures generated in subduction-to-collisional orogens through time has been modeled in a preliminary manner by Chacko (2003). Because of higher abundances of heat-producing elements, older crust would have produced 1.2, 1.6, and 2.3 times more heat at 1.0, 2.0, and 3.0 Ga, respectively. The effect of increased RHP on the thermal evolution of crust instantaneously thickened from 35 to 70 km suggests that collisional orogens as old as 1.0 Ga would have experienced maximum mid-crustal (20–40 km) temperatures that were approximately 70°C hotter than those of modern orogens. Collisional orogens at 2.0 and 3.0 Ga would have produced mid-crustal temperatures that were approximately 180 and 370°C, respectively, hotter than modern orogens. Evidence of these significantly higher and progressively declining temperatures of metamorphism is lacking in the rock record, and in those circumstances where high-grade or UHT metamorphic conditions are attributed to radioactive heating alone, the U and Th contents are high and modeling the heat production at the age of metamorphism achieves the temperatures required (Andreoli et al., 2006; Kramers et al., 2001). Indeed, the Archean rock record is dominated by rather ordinary conditions of crustal metamorphism and there is accumulating evidence that thermal conditions in Earth's very early continental crust also were rather ordinary (e.g., Valley, 2005; Watson and Harrison, 2005, 2006; Watson et al.,

2006), although the evidence is by no means without dispute (e.g., Nemchin et al., 2006); one inference we may draw from this observation is that a model of instantaneous doubling of the continental crust is probably inappropriate as a model of Archean orogenesis (cf. Cagnard et al., 2006).

It is outside the scope of this paper to speculate on Archean tectonics. However, there are documented differences in structural styles between Archean, Proterozoic, and Proterozoic orogenic belts (e.g., Hamilton, 1998; Marshak, 1999; Cagnard et al., 2006) and dramatic changes in the metamorphic record through the Archean to Proterozoic and Proterozoic to Phanerozoic transitions, as documented in this paper. These differences and changes probably were linked to—and reflect—non-linear switches in mantle convection, although exactly what these might be is a matter of speculation and modeling. For example, whether the Archean to Proterozoic transition effected a change from convective overturn events to slab-induced overturn events in a layered mantle (Davies, 1995) or effected a change to episodic catastrophic subduction of ocean lithosphere and consequent plume events (O’Neal et al., 2005) demonstrates that whatever the mechanism might be, two-layer convecting mantle models generate episodic whole-mantle convection overturn events on timescales appropriate to major transitions in Earth history. Furthermore, it is possible that some form of oceanic “plate tectonics” operated in the Archean despite the high temperature of the mantle (e.g., Davies, 2006). The relationship between superplume events and heterogeneities in the deep mantle (Burke and Torsvik, 2004) and the strongly exothermic perovskite–post-perovskite phase transition within the lower mantle (e.g., Yuen and Monnereau, 2005) remains enigmatic, but demonstrate the importance of coupling in Earth-system behavior.

### Concluding Notes

1. I propose that one defining feature of a plate tectonics regime is the development of two contrasting thermal environments where plates converge—on modern Earth these are the subduction zone and the backarc, respectively—that are registered in the rock record as contemporaneous different types of metamorphism, viz. E-HPG and UHT metamorphism in Proterozoic–Cambrian orogens, and BS to UHP and moderate-P, high-T metamorphic belts in the Phanerozoic orogens (Figs. 7 and 8).

2. I suggest that we investigate further the implications of the crustal recycling implied by the geochemistry of Archean and Proterozoic diamonds to determine whether they represent (scant) evidence of UHP metamorphism before the Ediacaran (?). This will enable us to establish whether the Proterozoic plate tectonics regime involved deep subduction of continental crust—as might be implied by the diamonds—but somehow prevented its exhumation, or whether the warmer thermal environment—implied by the widespread occurrence of E-HPG metamorphism and the apparent absence of BS and UHP metamorphism—somehow prevented deep subduction of continental crust before the Neoproterozoic Era.

3. I argue the change from a Proterozoic plate tectonic regime to a modern plate tectonic regime during the Neoproterozoic–Cambrian reflects a change in global geodynamics and pattern of mantle convection. The change was responsible for the reconfiguration of Rodinia and its transformation into Gondwana—the last of the supercontinents—and the progressive destruction of a long-standing hemispheric-scale oceanic realm by closure of the Rheic Ocean (during the Paleozoic) and subduction successively along the Paleo-Tethyan and Tethyan trenches (during the Mesozoic; e.g., Collins, 2003).

4. Finally, UHT metamorphism clearly must be limited by the liquidus for crustal rocks, although for all practical purposes an upper limit of  $\sim 1,100^{\circ}\text{C}$  seems to operate in nature. However, the maximum depth in the mantle from which crustal protoliths may be exhumed back to Earth’s surface—the point of no return—is not well established, although it appears to be around 10 GPa. Clearly, we should determine the natural limits of terrane-scale crustal metamorphism at the extremes of temperature and pressure.

### Acknowledgments

I thank Louie for the invitation to take part in the celebration of his career achievements at Stanford in December 2005, Gary Ernst for facilitating the celebration and the publication of papers from the event, Barry Reno for help in drafting the figures, and Tim Johnson and Aaron Martin for rapid but constructive reviews at short notice! Research in the Laboratory for Crustal Petrology is supported by the University of Maryland and NSF grant EAR-0227553.

## REFERENCES

- Abbott, D., and Menke, W., 1990, Length of the global plate boundary at 2.4 Ga: *Geology*, v. 18, p. 58–61.
- Agard, P., Labrousse, L., Elvevold, S., and Lepvrier, C., 2005, Discovery of Paleozoic Fe-Mg carpholite on Motalafjella, Svalbard Caledonides: A milestone for subduction zone gradients: *Geology*, v. 33, p. 761–764.
- Anczkiewicz, R., Platt, J. P., Thirlwall, M. F., and Wakabayashi, J., 2004, Franciscan subduction off to a slow start: Evidence from high-precision Lu-Hf garnet ages on high-grade blocks: *Earth and Planetary Science Letters*, v. 225, p. 147–161.
- Anderson, D. L., 2002, How many plates? *Geology*, v. 30, p. 411–414.
- Andreoli, M. A. G., Hart, R. J., Ashwal, L. D., and Coetzee, H., 2006, Correlations between U, Th content, and metamorphic grade in the Western Namaqualand Belt, South Africa, with implications for radioactive heating of the crust: *Journal of Petrology* [doi:10.1093/petrology/egl004].
- Arima, M., and Barnett, R. L., 1984, Sapphirine bearing granulites from the Sipiwek Lake area of the late Archean Pikwitonei granulite terrain, Manitoba, Canada: *Contributions to Mineralogy and Petrology*, v. 88, p. 102–112.
- Arima, M., Kerrich, R., and Thomas, A., 1986, Sapphirine-bearing paragneiss from the northern Grenville province in Labrador, Canada: Protolith composition and metamorphic P-T conditions: *Geology*, v. 14, p. 844–847.
- Attoh, K., 1998a, High-pressure granulite facies metamorphism in the Pan-African Dahomeyide orogen, West Africa: *Journal of Geology*, v. 106, p. 236–246.
- Attoh, K., 1998b, Models for orthopyroxene-plagioclase and other corona reactions in metanorites, Dahomeyide orogen, West Africa: *Journal of Metamorphic Geology*, v. 16, p. 345–362.
- Attoh, K., Dallmeyer, R. D., and Affaton, P., 1997, Chronology of nappe assembly in the Pan-African Dahomeyide orogen, West Africa: Evidence from  $^{40}\text{Ar}/^{39}\text{Ar}$  mineral ages: *Precambrian Research*, v. 82, p. 153–171.
- Baba, S., 1998, Proterozoic anticlockwise P-T path of the Lewisian complex of South Harris, outer Hebrides, NW Scotland: *Journal of Metamorphic Geology*, v. 16, p. 819–841.
- Baba, S., 1999, Sapphirine-bearing orthopyroxene-kyanite/sillimanite granulites from South Harris, NW Scotland: Evidence for Proterozoic UHT metamorphism in the Lewisian: *Contributions to Mineralogy and Petrology*, v. 136, p. 33–47.
- Baba, S., 2003, Two stages of sapphirine formation during prograde and retrograde metamorphism in the Paleoproterozoic Lewisian complex in South Harris, NW Scotland: *Journal of Petrology*, v. 44, p. 329–354.
- Bagas, L., 2004, Proterozoic evolution and tectonic setting of the northwest Paterson Orogen, western Australia: *Precambrian Research*, v. 128, p. 475–496.
- Bakun-Czubarow, N., 1991, On the possibility of occurrence of quartz pseudomorphs after coesite in the eclogite-granulite rock series of the Złote Mountains in the Sudetes (SW Poland): *Archiwum Mineralogiczne*, v. 47, p. 5–16.
- Bakun-Czubarow, N., 1992, Quartz pseudomorphs after coesite and quartz exsolutions in eclogitic omphacites of the Złote Mountains in the Sudetes (SW Poland): *Archiwum Mineralogiczne*, v. 48, p. 3–25.
- Baldwin, J. A., Bowring, S. A., and Williams, M. L., 2003, Petrologic and geochronologic constraints on high-pressure, high-temperature metamorphism in the Snowbird tectonic zone, Canada: *Journal of Metamorphic Geology*, v. 21, p. 81–98.
- Baldwin, J. A., Bowring, S. A., Williams, M. L., and Mahan, K. H., 2006, Geochronological constraints on the evolution of high-pressure felsic granulites from an integrated electron microprobe and ID-TIMS geochemical study: *Lithos* [doi:10.1016/j.lithos.2005.08.009].
- Baldwin, J. A., Bowring, S. A., Williams, M. L., and Williams, I. S., 2004, Eclogites of the Snowbird Tectonic Zone: Petrological and U-Pb geochronological evidence for Paleoproterozoic high-pressure metamorphism in the western Canadian Shield: *Contributions to Mineralogy and Petrology*, v. 147, p. 528–548.
- Baldwin, J. A., Powell, R., Brown, M., Moraes, R., and Fuck, R. A., 2005, Mineral equilibria modeling of ultrahigh-temperature metamorphism: an example from the Anápolis-Itaúçu Complex, central Brazil: *Journal of Metamorphic Geology*, v. 23, p. 511–531.
- Baldwin, S. L., Webb, L. E., Monteleone, B. D., Little, T. A., Fitzgerald, P. G., and Chappell, J. L., 2005, Metamorphism and exhumation of the youngest known HP-UHP terrane on Earth, eastern Papua New Guinea [abs.]: *EOS (Transactions of the American Geophysical Union)*, v. 86, abstract V54 B-01.
- Ballèvre, M., Marchand, J., Godard, G., Goujou, J. C., Wyls, J.-C., and Wyls, R., 1994, Eo-Hercynian events in the Armorican Massif, in Keppie, J. D., ed., *Pre-Mesozoic geology in France and related areas*: Berlin, Germany, Springer, p. 183–194.
- Ballèvre, M., Pitra, P., and Bohn, M., 2003, Lawsonite growth in the epidote blueschist from the Ile de Groix (Armorican Massif, France): A potential geobarometer: *Journal of Metamorphic Geology*, v. 21, p. 723–735.
- Barbosa, J., Martin, H., and Peucat, J.-J., 2004, Paleoproterozoic dome-forming structures related to granulite-facies metamorphism, Jequié Block, Bahia, Brazil: Petrogenetic approaches: *Precambrian Research*, v. 135, p. 105–131.
- Becker, H., 1997, Sm-Nd garnet ages and cooling history of high-temperature garnet peridotite massifs and

- high-pressure granulites from lower Austria: Contributions to Mineralogy and Petrology, v. 127, p. 224–236.
- Bégin, N. J., and Pattison, D. R. M., 1994, Metamorphic evolution of granulites in the Minto block, northern Québec: Extraction of peak P–T conditions taking account of late Fe–Mg exchange: Journal of Metamorphic Geology, v. 12, p. 411–428.
- Berman, R. G., and Bostock, H. H., 1997, Metamorphism in the northern Taltson magmatic zone, Northwest Territories: Canadian Mineralogist, v. 35, p. 1069–1091.
- Bertrand, Ph., Ouzegane, Kh., and Kienast, J.-R., 1992, P–T–X relationships in the Precambrian Al–Mg-rich granulites from In Ouzal, Hoggar, Algeria: Journal of Metamorphic Geology, v. 10, p. 17–31.
- Bhattacharya, S., 2004, Reply to comments by Sengupta, Raith, and Dasgupta on S. Bhattacharya and R. Kar (2002): Contributions to Mineralogy and Petrology, v. 147, p. 126–127.
- Bhattacharya, S., and Kar, R., 2002, High-temperature dehydration melting and decompressive P–T path in a granulite complex from the Eastern Ghats, India: Contributions to Mineralogy and Petrology, v. 143, p. 175–191.
- Bhowmik, S. K., Basu Sarbadhikari, A., Spiering, B., and Raith, M. M., 2005, Mesoproterozoic reworking of Paleoproterozoic ultrahigh-temperature granulites in the Central Indian tectonic zone and its implications: Journal of Petrology, v. 46, p. 1085–1119.
- Bingen, B., Demaiffe, D., and Delhal, J., 1988, Aluminous granulites of the Archean craton of Kasai (Zaire): Petrology and P–T conditions: Journal of Petrology, v. 29, p. 899–919.
- Bleeker, W., 2003, The late Archean record: A puzzle in ca. 35 pieces, Lithos, v. 71, p. 99–134.
- Bohlen, S. R., and Essene, E. J., 1979, A critical evaluation of two-pyroxene thermometry in Adirondack granulites: Lithos, v. 12, p. 335–345.
- Bohlen, S. R., and Essene, E. J., 1980, Evaluation of coexisting garnet-biotite, garnet-clinopyroxene, and other Mg–Fe exchange thermometers in the Adirondack granulites: Geological Society of America Bulletin, v. 91, p. 685–719.
- Böhm, C.O., 2004, Neoarchean high-grade metamorphism of the Pikwitonei Granulite Domain NW Superior Craton [abs.]: Geological Association of Canada–Mineralogical Association of Canada Joint Annual Meeting, St. Catharines 2004, Abstracts Volume 29, p. 330.
- Bosse, V., Féraud, G., Ruffet, G., Ballèvre, M., Peucat, J. J., and De Jong, K., 2000, Late Devonian subduction and early-orogenic exhumation of eclogite-facies rocks from the Champtoceaux Complex (Variscan belt, France): Geological Journal, v. 35, p. 297–325.
- Bosse, V., Féraud, G., Ballèvre, M., Peucat, J.-J., and Corsini, M., 2005, Rb–Sr and  $^{40}\text{Ar}/^{39}\text{Ar}$  ages in blueschists from the Ile de Groix (Armorican Massif, France): Implications for closure mechanisms in isotopic systems: Chemical Geology, v. 220, p. 21–45.
- Bostock, B. C., Jones, R. E., Ernst, W. G., Chen, C., Leech, M. L., and Beane, R. J., 2003, Low-temperature microdiamond aggregates in the Maksyutov metamorphic complex, South Ural Mountains, Russia: American Mineralogist, v. 88, p. 1709–1717.
- Boven, A., Theunissen, K., Sklyarov, E., et al., 1999, Timing of exhumation of a high-pressure mafic granulite terrane of the Paleoproterozoic Ubende belt (West Tanzania): Precambrian Research, v. 93, p. 119–137.
- Brandt, S., Klemd, R., and Okrusch, M., 2003, Ultrahigh-temperature metamorphism and multistage evolution of garnet-orthopyroxene granulites from the Proterozoic Epupa Complex, NW Namibia: Journal of Petrology, v. 44, p. 1121–1144.
- Brown, M., 1998a, Unpairing metamorphic belts: P–T paths and a tectonic model for the Ryoke Belt, southwest Japan: Journal of Metamorphic Geology, v. 16, p. 3–22.
- Brown, M., 1998b, Ridge-trench interactions and high–low-P metamorphism, with particular reference to the Cretaceous evolution of the Japanese Islands, in Treloar, P. J., and O'Brien, P. J., eds. What drives metamorphism and metamorphic reactions? Geological Society Special Publication no. 138, p. 131–163.
- Brown, M., 2001, From microscope to mountain belt: 150 years of petrology and its contribution to understanding geodynamics, particularly the tectonics of orogens: Journal of Geodynamics, v. 32, p. 115–164.
- Brown, M., 2002a, Prograde and retrograde processes in migmatites revisited: Journal of Metamorphic Geology, v. 20, p. 25–40.
- Brown, M., 2002b, Plate margin processes and “paired” metamorphic belts in Japan. Comment on “Thermal effects of ridge subduction and its implication for the origin of granitic batholith and paired metamorphic belts” by H. Iwamori: Earth and Planetary Science Letters, v. 199, p. 483–492.
- Brown, M., and Raith, M., 1996, First evidence of ultrahigh-temperature decompression from the granulite province of Southern India: Journal of the Geological Society, London, v. 153, p. 819–822.
- Brueckner, H. K., Henning, S., Sorensen, S., and Harlow, G. E., 2005, Synchronous Sm–Nd mineral ages from HP terranes on both sides of the Motagua Fault of Guatemala: Convergent suture and strike slip fault? [abs.]: EOS (Transactions, American Geophysical Union), v. 86, Fall Meeting Supplement, Abstract T23D-04.
- Burke, K., Dewey, J. F., and Kidd, W. S. F., 1977, World distribution of sutures—sites of former oceans: Tectonophysics, v. 40, p. 69–99.
- Burke, K., and Torsvik, T. H., 2004, Derivation of large igneous provinces of the past 200 million years from long-term heterogeneities in the deep mantle: Earth and Planetary Science Letters, v. 227, p. 531–538.

- Caby, R., 1994, Precambrian coesite from northern Mali: First record and implications for plate tectonics in the trans-Saharan segment of the Pan-African belt: *European Journal of Mineralogy*, v. 6, p. 235–244.
- Cagnard, F., Durrieu, N., Gapais, D., Brun, J.-P., and Ehlers, C., 2006, Crustal thickening and lateral flow during compression of hot lithospheres, with particular reference to Precambrian times: *Terra Nova*, v. 18, p. 72–78.
- Camacho, A., Lee, J. K. W., Hensen, B. J., and Braun, J., 2005, Short-lived orogenic cycles and the eclogitization of cold crust by spasmodic hot fluids: *Nature*, v. 435, p. 1191–1196.
- Campos Neto, M. D., and Caby, R., 1999, Neoproterozoic high-pressure metamorphism and tectonic constraints from the nappe system south of the Sao Francisco Craton, southeast Brazil: *Precambrian Research*, v. 97, p. 3–26.
- Campos Neto, M. D., and Caby, R., 2000, Terrane accretion and upward extrusion of high-pressure granulites in the Neoproterozoic nappes of southeast Brazil: Petrological and structural constraints: *Tectonics*, v. 19, p. 669–687.
- Carlson, W. D., Anderson, S. D., Mosher, S., Davidow, J. S., Crawford, W. D., and Lane, E. D., 2007, High-pressure metamorphism in the Texas Grenville Orogen: Mesoproterozoic subduction of the southern Laurentian continental margin: *International Geology Review*, v. 49, in press.
- Caron, J.-M., and Péquignot, G., 1986, The transition between blueschist and lawsonite-bearing eclogites based on observations from Corsican metabasalts: *Lithos*, v. 19, p. 205–218.
- Carswell, D. A., and O'Brien, P. J., 1993, Thermobarometry and geotectonic significances of high-pressure granulites: Examples from the Moldanubian Zone of the Bohemian Massif in Lower Austria: *Journal of Petrology*, v. 34, p. 427–459.
- Carswell, D. A., Tucker, R. D., O'Brien, P. J., and Krogh, T. E., 2003, Coesite micro-inclusions and the U/Pb age of zircons from the Hareidland Eclogite in the Western Gneiss region of Norway: *Lithos*, v. 67, p. 181–190.
- Cartigny, P., Chin, L., Viljoens, K. S., and Robinson, D., 2004, Early Proterozoic ultrahigh pressure metamorphism: Evidence from microdiamonds: *Science*, v. 304, p. 853–855.
- Cartwright, I., and Barnicoat, A. C., 1989, Evolution of the Scourian complex, in Daly, J. S., Cliff, R. A., and Yardley, B. W. D., eds., *Evolution of metamorphic belts*: Geological Society Special Publication 43, p. 297–301.
- Castelli, D., Rolfo, F., Compagnoni, R., and Xu, S. T., 1998, Metamorphic veins with kyanite, zoisite, and quartz in the Zhu-Jia-Chong eclogites, Dabie Shan, China: *The Island Arc*, v. 7, p. 159–173.
- Chacko, T., 2003, Are there systematic differences between the metamorphic temperatures of Phanerozoic, Proterozoic, and Archean collisional orogens? [abs.]: Geological Association of Canada, Mineralogical Association of Canada, Society of Economic Geologists Joint Meeting, Vancouver, B.C., Canada, Abstracts Volume 28, p. 66.
- Chacko, T., Kumar, G. R. R., and Newton, R. C., 1987, Metamorphic P–T conditions of the Kerala (south India) Khondalite Belt, a granulite facies supracrustal terrain: *Journal of Geology*, v. 95, p. 343–358.
- Chacko, T., Lamb, M., and Farquhar, J., 1996, Ultrahigh temperature metamorphism in the Kerala Khondalite Belt: Gondwana Research Group Memoir 3, p. 157–165.
- Chesnokov, B. V., and Popov, V. A., 1965, Phenomenon of volume growth of quartz grains in eclogites of South Urals: *Doklady Akademii Nauk SSSR*, v. 162, p. 909.
- Chopin, C., 1984, Coesite and pure pyrope in high grade blueschists of the Western Alps: A first record and some consequences: *Contributions to Mineralogy and Petrology*, v. 86, p. 107–118.
- Chopin, C., 2003, Ultrahigh-pressure metamorphism: tracing continental crust into the mantle: *Earth and Planetary Science Letters*, v. 212, p. 1–14.
- Chopin, C., Henry, C., and Marchand, A., 1991, Geology and petrology of the coesite-bearing terrain, Dora Maira Massif, Western Alps: *European Journal of Mineralogy*, v. 3, p. 263–291.
- Clarke, G. L., Klepeis, K. A., and Daczko, N. R., 2000, Cretaceous high-P granulites at Milford Sound, New Zealand: metamorphic history and emplacement in a convergent margin setting: *Journal of Metamorphic Geology*, v. 18, p. 359–374.
- Cloos, M., 1993, Lithospheric buoyancy and collisional orogenesis: Subduction of oceanic plateaus, continental margins, island arcs, spreading ridges, and seamounts: *Geological Society of America Bulletin*, v. 105, p. 715–737.
- Collins, W. J., 2002, Nature of extensional accretionary orogens: *Tectonics*, v. 21 [doi:10.1029/2000TC001272].
- Collins, W. J., 2003, Slab pull, mantle convection, and Pangaea assembly and dispersal: *Earth and Planetary Science Letters*, v. 205, p. 225–237.
- Collins, A. S., Reddy, S. M., Buchan, C., and Mruma, A., 2004, Temporal constraints on Paleoproterozoic eclogite formation and exhumation (Usagaran Orogen, Tanzania): *Earth and Planetary Science Letters*, v. 224, p. 175–192.
- Coney, P. J., Jones, D. L., and Monger, J. W. H., 1980, Cordilleran suspect terranes: *Nature*, v. 28, p. 329–333.
- Cooke, R. A., and O'Brien, P. J., 2001, Resolving the relationship between high P-T rocks and gneisses in collisional terranes: An example from the Gföhl gneiss-granulite association in the Moldanubian Zone, Austria: *Lithos*, v. 58, p. 33–54.
- Cooke, R. A., O'Brien, P. J., and Carswell, D. A., 2000, Garnet zoning and the identification of equilibrium mineral compositions in high-pressure-temperature

- granulites from the Moldanubian Zone, Austria: *Journal of Metamorphic Geology*, v. 18, p. 551–569.
- Coolen, J., 1980, Chemical petrology of the Furua granulite complex, southern Tanzania: *GUA Papers of Geology, Series 1*, p. 13.
- Corfu, F., Ravna, E. J. K., and Kullerød, K., 2003, A Late Ordovician U-Pb age for the Tromsø Nappe eclogites, uppermost allochthon of the Scandinavian Caledonides: *Contributions to Mineralogy and Petrology*, v. 145, p. 502–513.
- Costa, S., Maluski, H., and Lardeaux, J.-M., 1993,  $^{40}\text{Ar}$ – $^{39}\text{Ar}$  chronology of Variscan tectono-metamorphic events in an exhumed crustal nappe: The Monts du Lyonnais complex (Massif Central, France): *Chemical Geology*, v. 105, p. 339–359.
- Cowan, D. S., Brandon, M. T., and Garver, J. I., 1997, Geologic tests of hypotheses for large coastwise displacements—a critique illustrated by the Baja British Columbia controversy: *American Journal of Science*, v. 297, p. 117–173.
- Currie, C. A., and Hyndman, R. D., 2005, The thermal structure of subduction zones: The case for a hot back-arc: *EOS (Transactions of the American Geophysical Union)*, v. 86, abstract T11B-0366.
- Cuthbert, S. J., Carswell, D. A., Krogh-Ravna, E. J., and Wain, A., 2000, Eclogites and eclogites in the Western Gneiss Region, Norwegian Caledonides: *Lithos*, v. 52, p. 165–195.
- Dahl, P. S., Hamilton, M. A., and Jercinovic, M. J., et al., 2005, Comparative isotopic and chemical geochronometry of monazite, with implications for U-Th-Pb dating by electron microprobe: An example from metamorphic rocks of the eastern Wyoming craton (USA): *American Mineralogist*, v. 90, p. 619–638.
- Dale, J., Powell, R., White, R. W., Elmer, F. L., and Holland, T. J. B., 2005, A thermodynamic model for Ca-Na clin amphiboles in  $\text{Na}_2\text{O}$ -CaO-FeO-MgO- $\text{Al}_2\text{O}_3$ - $\text{SiO}_2$ - $\text{H}_2\text{O}$ -O for petrological calculations: *Journal of Metamorphic Geology*, v. 23, p. 771–791.
- Dallmeyer, R. D., Johansson, L., and Möller, C., 1992a, Chronology of Caledonian high-pressure granulite-facies metamorphism, uplift, and deformation within northern parts of the Western Gneiss Region, Norway: *Geological Society of America Bulletin*, v. 104, p. 444–455.
- Dallmeyer, R. D., Neubauer, F., and Hock, V., 1992b, Chronology of Late Paleozoic tectonothermal activity in the southeastern Bohemian Massif, Austria (Moldanubian and Moravo-Silesian Zones)— $\text{Ar}^{40}$ - $\text{Ar}^{39}$  mineral age controls: *Tectonophysics*, v. 210, p. 135–153.
- Dallmeyer, R. D., Peucat, J. J., Hirajima, T., and Ohta, Y., 1989, Tectonothermal chronology within a blueschist-eclogite complex, west-central Spitsbergen, Svalbard: Evidence from  $^{40}\text{Ar}/^{39}\text{Ar}$  and Rb-Sr mineral ages: *Lithos*, v. 24, p. 291–304.
- Dallwitz, W. B., 1968, Co-existing sapphirine and quartz in granulite from Enderby Land, Antarctica: *Nature*, v. 219, p. 476.
- Dasgupta, S., Sanyal, S., Sengupta, P., and Fukuoka, M., 1994, Petrology of granulites from Anakapalle—evidence for Proterozoic decompression in the Eastern Ghats, India: *Journal of Petrology*, v. 35, p. 433–459.
- Dasgupta, S., Sengupta, P., Fukuoka, M., and Chakraborti, S., 1992, Dehydration melting, fluid buffering and decompressional P-T path in a granulite complex from the Eastern Ghats, India: *Journal of Metamorphic Geology*, v. 10, p. 777–788.
- Davies, G. F., 1995, Punctuated tectonic evolution of the Earth: *Earth and Planetary Science Letters*, v. 136, p. 363–379.
- Davies, G. F., 2006, Gravitational depletion of the early Earth's upper mantle and the viability of early plate tectonics: *Earth and Planetary Science Letters* [doi:10.1016/j.epsl.2006.01.053].
- Dawson, J. B., and Smith, J. V., 1987, Reduced sapphirine granulite xenoliths from the Lace Kimberlite, South Africa: Implications for the deep structure of the Kaapvaal craton: *Contributions to Mineralogy and Petrology*, v. 95, p. 376–383.
- Dawson, J. B., Harley, S. L., Rudnick, R. L., and Ireland, T., 1997, Equilibrium and reaction in Archaean sapphirine granulite xenoliths from the Lace kimberlite pipe, South Africa: *Journal of Metamorphic Geology*, v. 15, p. 253–266.
- Del Lama, E. A., Zanardo, A., Oliveira, M. A. F., and Morales, N., 2000, Exhumation of high-pressure granulites of the Guaxupé Complex, southeastern Brazil: *Geological Journal*, v. 35, p. 231–249.
- Dempster, T. J., Harrison, T. N., Brown, P. E., and Hutton, D. H. W., 1991, Low-pressure granulites from the Ketilidian mobile belt of southern Greenland: *Journal of Petrology*, v. 32, p. 979–1004.
- de Roever, W. P., 1956, Some differences between post-Paleozoic and older regional metamorphism: *Geologie en Mijnbouw*, v. 18, p. 123–127.
- de Roever, W. P., 1965, On the cause of the preferential distribution of certain metamorphic minerals in orogenic belts of different age: *Geologische Rundschau*, v. 54, p. 933–943.
- Dewey, J. F., and Kidd, W. S. F., 1977, Geometry of plate accretion: *Geological Society of America Bulletin*, v. 88, p. 960–968.
- Diener, J. F. A., Stevens, G., Kisters, A. F. M., and Poujol, M., 2005, Metamorphism and exhumation of the basal parts of the Barberton greenstone belt, South Africa: Constraining the rates of Mesoarchaean tectonism: *Precambrian Research*, v. 143, p. 87–112.
- Dirks, P. H. G. M., and Sithole, T. A., 1999, Eclogites in the Makuti gneisses of Zimbabwe: Implications for the tectonic evolution of the Zambezi Belt in southern Africa: *Journal of Metamorphic Geology*, v. 17, p. 593–612.

- Dobrzhinetskaya, L., and Green, H. W., 2005, Ultra-deep subduction of continental crust: Evidence from natural rocks and experimental investigations [abs.]: EOS (Transactions of the American Geophysical Union), v. 86, abstract V51E-01.
- Dobrzhinetskaya, L. F., Renfro, A. P., and Green, H. W., 2004, Synthesis of skeletal diamonds: Implications for microdiamond formation in orogenic belts: *Geology*, v. 32, p. 869–872.
- Dobrzhinetskaya, L. F., Wirth, R., Green, H. W., 2005, Direct observation and analysis of a trapped COH fluid growth medium in metamorphic diamond: *Terra Nova*, v. 17, p. 472–477.
- Domanik, K. J., and Holloway, J. R., 2000, Experimental synthesis and phase relations of phengitic muscovite from 6.5 to 11 GPa in a calcareous metapelite from the Dabie Mountains, China: *Lithos*, v. 52, p. 51–77.
- Droop, G. T. R., 1989, Reaction history of garnet-sapphirine granulites and conditions of Archaean high-pressure granulite facies metamorphism in the Central Limpopo Mobile Belt, Zimbabwe: *Journal of Metamorphic Geology*, v. 7, p. 383–403.
- Droop, G. T. R., and Bucher-Nurminen, K., 1984, Reaction textures and metamorphic evolution of sapphirine-bearing granulites from the Gruf Complex, Italian Central Alps: *Journal of Petrology*, v. 25, p. 766–803.
- Dufour, E., Lardeaux, J. M., and Coffrant, D., 1985, Eclogites and granulites in the Monts du Lyonnais area: An eo-Hercynian plurifacial metamorphic evolution: *Comptes Rendus de l'Academie des Sciences, Paris*, v. 300, p. 141–144.
- Dziggel, A., Stevens, G., Poujol, M., Anhaeusser, C. R., and Armstrong, R. A., 2002, Metamorphism of the granite–greenstone terrane south of the Barberton greenstone belt, South Africa: an insight into the tectono-thermal evolution of the ‘lower’ portions of the Onverwacht Group: *Precambrian Research*, v. 114, p. 221–247.
- Ellis, D. J., 1980, Osumilite-sapphirine-quartz granulites from Enderby Land, Antarctica—P-T conditions of metamorphism, implications for garnet-cordierite equilibria and the evolution of the deep crust: *Contributions to Mineralogy and Petrology*, v. 74, p. 201–210.
- Ellis, D. J., Sheraton, J. W., England, R. N., and Dallwitz, W. B., 1980, Osumilite-sapphirine-quartz granulites from Enderby Land, Antarctica—mineral assemblages and reactions: *Contributions to Mineralogy and Petrology*, v. 72, p. 123–143.
- Elvevold, S., and Gilotti, J. A., 2000, Pressure-temperature evolution of retrogressed kyanite eclogites, Weinschenk Island, North-East Greenland Caledonides: *Lithos*, v. 53, p. 127–147.
- Elvevold, S., Thrane, K., and Gilotti, J. A., 2003, Metamorphic history of high-pressure granulites in Payer Land, Greenland Caledonides: *Journal of Metamorphic Geology*, v. 21, p. 49–63.
- Ernst, W. G., 1972, Occurrence and mineralogic evolution of blueschist belts with time: *American Journal of Science*, v. 272, p. 657–668.
- Ernst, W. G., 2005, Alpine and Pacific styles of Phanerozoic mountain building: Subduction-zone petrogenesis of continental crust: *Terra Nova*, v. 17, p. 165–188.
- Farquhar, J., Chacko, T., and Ellis, D. J., 1996, Preservation of oxygen isotope compositions in granulites from northwestern Canada and Enderby Land, Antarctica: Implications for high-temperature isotopic thermometry: *Contributions to Mineralogy and Petrology*, v. 125, p. 213–224.
- Farquhar, J., Wing, B. A., McKeegan, K. D., Harris, J. W., Cartigny, P., and Thiemens, M. H., 2002, Mass-independent sulfur of inclusions in diamond and sulfur recycling on early Earth: *Science*, v. 298, p. 2369–2372.
- Faryad, S. W., 1999, Metamorphic evolution of the Precambrian South Badakhshan block, based on mineral reactions in metapelites and metabasites associated with whiteschists from Sare Sang (Western Hindu Kush, Afghanistan): *Precambrian Research*, v. 98, p. 223–241.
- Ferreira Filho, C. F., Kamo, S. L., Fuck, R. A., Krogh, T. E., and Naldrett, A. J., 1994, Zircon and rutile U-Pb geochronology of the Niquelândia layered mafic and ultramafic intrusion, Brazil: Constrains for the timing of magmatism and the high-grade metamorphism: *Precambrian Research*, v. 68, p. 241–255.
- Ferreira Filho, C. F., Moraes, R., Fawcett, J. J., and Naldrett, A. J., 1998, Amphibolite to granulite progressive metamorphism in the Niquelândia Complex, Central Brazil: Regional tectonic implication: *Journal of South American Earth Sciences*, v. 11, p. 35–50.
- Fiala, J., Matejovská, O., and Vaňková, V. B., 1987, Moldanubian granulites: Source material and petrogenetic considerations: *Neues Jahrbuch für Mineralogie, Abhandlungen*, v. 157, p. 133–165.
- Fitzherbert, J. A., Clarke, G. L., and Powell, R., 2003, Lawsonite-omphacite-bearing metabasites of the Pam Peninsula, NE New Caledonia: Evidence for disrupted blueschist- to eclogite-facies conditions: *Journal of Petrology*, v. 44, 1805–1831.
- Fitzherbert, J. A., Clarke, G. L., and Powell, R., 2005, Preferential retrogression of high-P sediments and the preservation of blueschist to eclogite-facies metabasites during exhumation, Diahot terrane, NE New Caledonia: *Lithos*, v. 83, p. 67–96.
- Fitzsimons, I. C. W., and Harley, S. L., 1994, The influence of retrograde cation exchange on granulite P-T estimates and a convergence technique for the recovery of peak metamorphic conditions: *Journal of Petrology*, v. 35, p. 543–576.
- Fonarev, V. I., Pilugin, S. M., Savko, K. A., and Novikova, M. A., 2006, Exsolution textures of ortho- and clinopyroxene in high-grade BIF of the Voronezh Crystalline Massif: Evidence of ultrahigh-temperature metamor-

- phism: *Journal of Metamorphic Geology*, v. 24, p. 135–151.
- Forster, M. A., and Lister, G. S., 2005, Several distinct tectono-metamorphic slices in the Cycladic eclogite-blueschist belt, Greece: *Contributions to Mineralogy and Petrology*, v. 150, p. 523–545.
- Fotoohi Rad, G. R., Droop, G. T. R., Amini, S., and Moazzen, M., 2005, Eclogites and blueschists of the Sistan Suture Zone, eastern Iran: A comparison of P-T histories from a subduction mélange: *Lithos*, v. 84, p. 1–24.
- Frazier, G., McDougall, I., Ellis, D. J., and Williams, I. S., 2000, Timing and rate of isothermal decompression in Pan-African granulites from Rundvågshetta, east Antarctica: *Journal of Metamorphic Geology*, v. 18, p. 441–454.
- Friend, C. R. L., Jones, K. A., and Burns, I. M., 2000, New high-pressure granulite event in the Moine Supergroup, northern Scotland: Implications for Taconic (early Caledonian) crustal evolution: *Geology*, v. 28, 543–546.
- Friend, C. R. L., and Kinny, P. D., 2001, A reappraisal of the Lewisian Gneiss Complex: Geochronological evidence for its tectonic assembly from disparate terranes in the Proterozoic: *Contributions to Mineralogy and Petrology*, v. 142, p. 198–218.
- Frost, B. R., Frost, C. D., Chamberlain, K. R., and Hughes, P. J., 2005, The 2.68–2.67 Teton–Wind River domain: Evidence for active-margin tectonics and continent-continent collision on the western margin of the Wyoming Province [abs.]: *Geological Society of America, Abstracts with Programs* 37, p. 505.
- Fukui, S., Watanabi, T., Itaya, T., and Leitch, E. C., 1995, Middle Ordovician high PT metamorphic rocks in eastern Australia: Evidence from K-Ar ages: *Tectonics*, v. 14, p. 1014–1020.
- Gao, J., and Klemd, R., 2003, Formation of HP-LT rocks and their tectonic implications in the western Tianshan Orogen, NW China: *Geochemical and age constraints: Lithos*, v. 66, p. 1–22.
- Gardien, V., Tegye, M., Lardeaux, J. M., Misseri, M., and Dufour, E., 1990, Crust-mantle relationships in the French Variscan chain: The example of the Southern Monts du Lyonnais unit (eastern French Massif Central): *Journal of Metamorphic Geology*, v. 8, p. 477–492.
- Gayk, T., Kleinschrodt, R., Langosch, A., and Seidel, E., 1995, Quartz exsolution in clinopyroxenes of high-pressure granulites from the Münchberg massif: *European Journal of Mineralogy*, v. 7, p. 1217–1220.
- Ghent, E. D., Stout, M. Z., and Erdmer, P., 1993, Pressure-temperature evolution of lawsonite-bearing eclogites, Pinchi Lake, British Columbia: *Journal of Metamorphic Geology*, v. 11, p. 279–290.
- Ghiribelli, B., Frezzotti, M. L., and Palmeri, R., 2002, Coesite in eclogites of the Lanterman Range (Antarctica): Evidence from textural and Raman studies: *European Journal of Mineralogy*, v. 14, p. 355–360.
- Gilotti, J. A., and Elvevold, S., 2002, Extensional exhumation of a high-pressure granulite terrane in Payer Land, Greenland Caledonides: Structural, petrologic, and geochronologic evidence from metapelites: *Canadian Journal of Earth Sciences*, v. 39, p. 1169–1187.
- Gilotti, J. A., and Ravna, E. J. K., 2002, First evidence for ultrahigh-pressure metamorphism in the North-East Greenland Caledonides: *Geology*, v. 30, p. 551–554.
- Godard, G., Martin, S., Prosser, G., Kienast, J. R., and Morten, L., 1996, Variscan migmatites, eclogites and garnet-peridotites of the Ulten zone, Eastern Austroalpine system: *Tectonophysics*, v. 259, p. 313–341.
- Goncalves, P., Guillot, S., Lardeaux, J.-M., Nicollet, C., and Mercier de Lepinay, B., 2000, Thrusting and sinistral wrenching in a pre-Eocene HP-LT Caribbean accretionary wedge (Samaná Peninsula, Dominican Republic): *Geodinamica Acta*, v. 13, p. 119–132.
- Goncalves, P., Nicollet, C., and Montel, J.-M., 2004, Petrology and *in situ* U-Th-Pb monazite geochronology of ultrahigh-temperature metamorphism from the Andriamena mafic unit, north-central Madagascar. Significance of a petrographical P-T path in a poly-metamorphic context: *Journal of Petrology*, v. 45, p. 1923–1957.
- Grambling, J. A., 1981, Pressures and temperatures in Precambrian metamorphic rocks: *Earth and Planetary Science Letters*, v. 53, p. 63–68.
- Green, H. W., 2005, Psychology of a changing paradigm: 40+ years of high-pressure metamorphism: *International Geology Review*, v. 47, p. 439–456.
- Green, H. W., Dobrzynetskiy, L., and Bozhilov, K. N., 2000, Mineralogical and experimental evidence for very deep exhumation from subduction zones: *Journal of Geodynamics*, v. 30, p. 61–76.
- Gregoire, M., Mattioli, N., Nicollet, C., Cottin, J. Y., Leyrit, H., Weis, D., Shimizu, N., and Giret, A., 1994, Oceanic mafic granulite xenoliths from the Kerguelen archipelago: *Nature*, v. 367, p. 360–363.
- Guo, J. H., O'Brien, P. J., and Zhai, M. G., 2002, High pressure granulites in the Sanggan area, North China craton: Metamorphic evolution, P-T paths and geotectonic significance: *Journal of Metamorphic Geology*, v. 20, p. 741–757.
- Hacker, B. R., 2006, Pressures and temperatures of ultrahigh-pressure metamorphism: Implications for UHP tectonics and H<sub>2</sub>O in subducting slabs: *International Geology Review*, v. 48, p. 1053–1066.
- Hamilton, W. B., 1998, Archean magmatism and deformation were not products of plate tectonics: *Precambrian Research*, v. 91, p. 143–179.
- Hansen, E. C., Janardhan, A. S., Newton, R. C., Prame, W. K. B. N., and Kumar, G. R., 1987, Arrested charnockite formation in southern India and Sri Lanka: *Contributions to Mineralogy and Petrology*, v. 96, p. 225–244.
- Hargraves, R. B., 1986, Faster spreading or greater ridge length in the Archean: *Geology*, v. 14, p. 750–752.

- Harley, S. L., 1985, Garnet-orthopyroxene bearing granulites from Enderby Land, Antarctica: Metamorphic pressure-temperature-time evolution of the Archaean Napier Complex: *Journal of Petrology*, v. 26, p. 819–856.
- Harley, S. L., 1989, The origins of granulites—a metamorphic perspective: *Geological Magazine*, v. 126, p. 215–247.
- Harley, S. L., 1998, Ultrahigh temperature granulite metamorphism (1050°C, 12 kbar) and decompression in garnet (Mg 70)–orthopyroxene–sillimanite gneisses from the Rauer Group, East Antarctica: *Journal of Metamorphic Geology*, v. 16, p. 541–562.
- Harley, S. L., 2004, Extending our understanding of ultrahigh temperature crustal metamorphism: *Journal of Mineralogical and Petrological Sciences*, v. 99, p. 140–562.
- Harley, S. L., and Green, D. H., 1982, Garnet ortho-pyroxene barometry for granulites and peridotites: *Nature*, v. 300, p. 697–701.
- Harley, S. L., Hensen, B. J., and Sheraton, J. W., 1990, Two-stage decompression in orthopyroxene–sillimanite granulites from Forefinger Point, Enderby Land, Antarctica: Implications for the evolution of the Archaean Napier Complex: *Journal of Metamorphic Geology*, v. 8, p. 591–613.
- Harley, S. L., and Motoyoshi, Y., 2000, Al zoning in orthopyroxene in a sapphirine quartzite: Evidence for >1120°C UHT metamorphism in the Napier complex, Antarctica, and implications for the entropy of sapphirine: *Contributions to Mineralogy and Petrology*, v. 138, p. 293–307.
- Harlov, D. E., and Milke, R., 2002, Stability of corundum + quartz relative to kyanite and sillimanite at high temperature and pressure: *American Mineralogist*, v. 87, p. 424–432.
- Harris, N. B. W., Holt, R. W., and Drury, S. A., 1982, Geobarometry, geothermometry and late Archean geotherms from the granulite facies terrain of south India: *Journal of Geology*, v. 90, p. 509–527.
- Harris, N. B. W., and Holland, T. J. B., 1984, The significance of cordierite-hypersthene assemblages from the Beitbridge region of the Central Limpopo Belt: evidence for rapid decompression in the Archaean: *American Mineralogist*, v. 69, p. 1036–1049.
- Hartz, E. H., Podladchikov, Y. Y., and Jettsetuen, E., 2005, UHP P-T-Time loops: A record of ultradeep subduction or tectonic overpressure? *Mitteilungen der Österreichischen Mineralogischen Gesellschaft*, v. 150.
- Hayob, J. L., Essene, E. J., Ruiz, J., Ortega-Gutiérrez, F., and Aranda-Gómez, J. J., 1989, Young high-temperature granulites from the base of the crust in central Mexico: *Nature*, v. 342, p. 265–268.
- Hensen, B. J., and Zhou, B., 1995, Retention of isotopic memory in garnets partially broken down during an overprinting granulite-facies metamorphism—implications for the Sm-Nd closure temperature: *Geology*, v. 23, p. 225–228.
- Hensen, B. J., Zhou, B., and Thost, D. E., 1995, Are reaction textures reliable guides to metamorphic histories? Timing constraints from garnet Sm-Nd chronology for “decompression” textures in granulites from Sostrene Island, Prydz Bay, Antarctica: *Geological Journal*, v. 30, p. 261–271.
- Hermann, J., 2003, Experimental evidence for diamond-facies metamorphism in the Dora-Maira Massif: *Lithos*, v. 70, p. 163–182.
- Hirajima, T., Banno, S., Hiroi, Y., and Ohta, Y., 1988, Phase petrology of eclogites and related rocks from the Motalafjella high-pressure metamorphic complex in Spitsbergen (Arctic Ocean) and its significance: *Lithos*, v. 22, p. 75–97.
- Hiroi, Y., Kishi, S., Nohara, T., Sato, K., and Goto, J., 1998, Cretaceous high-temperature rapid loading and unloading in the Abukuma metamorphic terrane, Japan: *Journal of Metamorphic Geology*, v. 16, p. 67–81.
- Hisada, K., and Miyano, T., 1996, Petrology and microthermometry of aluminous rocks in the Botswana Limpopo Central Zone: Evidence for isothermal decompression and isobaric cooling: *Journal of Metamorphic Geology*, v. 14, p. 183–197.
- Hodges, K. V., 2003, Geochronology and thermochronology in orogenic systems, in Rudnick, R. L., ed., *The crust: Treatise on geochemistry*, v. 3: Oxford, UK: Elsevier-Pergamon, p. 263–292.
- Hokada, T., 2001, Feldspar thermometry in ultrahigh-temperature metamorphic rocks: Evidence of crustal metamorphism attaining ~1100°C in the Archean Napier complex, east Antarctica: *American Mineralogist*, v. 86, p. 932–938.
- Hokada, T., and Harley, S. L., 2004, Zircon growth in UHT leucosome: Constraints from zircon-garnet rare earth elements (REE) relations in Napier complex, east Antarctica: *Journal of Mineralogical and Petrological Sciences*, v. 99, p. 180–190.
- Hokada, T., Misawa, K., Yokoyama, K., Shiraishi, K., and Yamaguchi, A., 2004, SHRIMP and electron microprobe chronology of UHT metamorphism in the Napier complex, east Antarctica: Implications for zircon growth at >1,000°C: *Contributions to Mineralogy and Petrology*, v. 147, p. 1–20.
- Holland, T., and Powell, R., 1996a, Thermodynamics of order-disorder in minerals. 1. Symmetric formalism applied to minerals of fixed composition: *American Mineralogist*, v. 81, p. 1413–1424.
- Holland, T., and Powell, R., 1996b, Thermodynamics of order-disorder in minerals. 2. Symmetric formalism applied to solid solutions: *American Mineralogist*, v. 81, p. 1425–1437.
- Holland, T. J. B., and Powell, R., 1998, An internally consistent thermodynamic data set for phases of petrolog-

- ical interest: *Journal of Metamorphic Geology*, v. 16, p. 309–343.
- Holland, T., and Powell, R., 2003, Activity-composition relations for phases in petrological calculations: An asymmetric multicomponent formulation: *Contributions to Mineralogy and Petrology*, v. 145, p. 492–501.
- Hollis, J. A., Harley, S. L., White, R. W., and Clarke, G. L., 2006, Preservation of evidence for prograde metamorphism in UHT HP granulites, South Harris, Scotland: *Journal of Metamorphic Geology*, v. 24, p. 263–279.
- Hoschek, G., 2001, Thermobarometry of metasediments in metabasites from the eclogite zone of the Hohe Tauern, Eastern Alps, Austria: *Lithos*, v. 59, p. 127–150.
- Hoschek, G., 2004, Comparison of calculated P-T pseudosections for a kyanite eclogite from the Tallon Window, Eastern Alps, Austria: *European Journal of Mineralogy*, v. 16, p. 59–72.
- Hughes, P. J., Frost, B. R., and Swapp, S. M., 2005, High-pressure granulite metamorphism and P-T path of the Archean layered gneiss unit, northern Teton, Wyoming [abs.]: *Geological Society of America, Abstracts with Programs*, v. 37, p. 505.
- Hyndman, R. D., Currie, C. A., and Mazzotti, S. P., 2005a, Subduction zone backarcs, mobile belts, and orogenic heat: *GSA Today*, v. 15, p. 4–10.
- Hyndman, R. D., Currie, C., and Mazzotti, S., 2005b, The origin of Global mountain belts: Hot subduction zone backarcs [abs.]: *EOS (Transactions of the American Geophysical Union)*, v. 86, abstract T11B-0367.
- Indares, A., 1993, Eclogitized gabbros from the eastern Grenville Province: Texture, metamorphic context, and implication: *Canadian Journal of Earth Sciences*, v. 30, p. 159–173.
- Indares, A., 1995, Metamorphic interpretation of high-pressure–temperature metapelites with preserved growth zoning in garnet, Eastern Grenville Province, Canadian Shield: *Journal of Metamorphic Geology*, v. 13, p. 475–486.
- Indares, A., 1997, Garnet-kyanite clinopyroxenites and garnet-kyanite restites from the Manicouagan Imbricate Zone: A case of high-P–high-T metamorphism in the Grenville Province: *Canadian Mineralogist*, v. 35, p. 1161–1171.
- Indares, A. D., 2003, Metamorphic textures and P-T evolution of high-P granulites from the Lelukuau terrane, NE Grenville Province: *Journal of Metamorphic Geology*, v. 21, p. 35–48.
- Indares, A., and Dunning, G., 2001, Partial melting of high-P-T metapelites from the Tshenukutish terrane (Grenville Province): *Petrography and U-Pb geochronology: Journal of Petrology*, v. 42, p. 1547–1565.
- Ireland, T. R., and Gibson, G. M., 1998, SHRIMP monazite and zircon geochronology of high grade metamorphism in New Zealand: *Journal of Metamorphic Geology*, v. 16, p. 149–167.
- Irfune, T., Ringwood, A. E., and Hibberson, W. O., 1994, Subduction of continental crust and terrigenous and pelagic sediments: an experimental study: *Earth and Planetary Science Letters*, v. 126, p. 351–368.
- Jacob, D. E., 2004, Nature and origin of eclogite xenoliths from kimberlites: *Lithos*, v. 77, p. 295–316.
- Jaekel, P., Kröner, A., Kamo, S. L., Brandl, G., and Wendt, J. L., 1997, Late Archean to Early Proterozoic granitoid magmatism and high-grade metamorphism in the central Limpopo belt, South Africa: *Journal of the Geological Society*, v. 154, 25–44.
- Jahn, B., Caby, R., and Monie, P., 2001, The oldest UHP eclogites of the world: Age of UHP metamorphism, nature of protoliths, and tectonic implications: *Chemical Geology*, v. 178, p. 143–158.
- James, D. T., Kamo, S., Krogh, T., and Nadeau, L., 2002, Preliminary report on U-Pb ages for intrusive rocks from the western Mealy Mountains and Wilson Lake terranes, Grenville Province, southern Labrador: Current Research: St. John's, Newfoundland, Department of Mines and Energy Geological Survey, Report 02-1, p. 67–77.
- Jan, M. Q., and Karim, A., 1995, Coronas and high-P veins in metagabbros of the Kohistan island arc, northern Pakistan: Evidence for crustal thickening during cooling: *Journal of Metamorphic Geology*, v. 13, p. 357–366.
- Janák, M., Froitzheim, F., Lupták, B., Brabec, M., and Krogh Ravna, E. J., 2004, First evidence for ultrahigh-pressure metamorphism of eclogites in Pohorje, Slovenia: Tracing deep continental subduction in the eastern Alps: *Tectonics*, v. 23, TC5014 [doi:10.1029/2004TC001641].
- Janardhan, A. S., Newton, R. C., and Hansen, E. C., 1982, The transformation of amphibolite facies gneiss to charnockite in southern Karnataka and northern Tamil Nadu, India: *Contributions to Mineralogy and Petrology*, v. 79, p. 130–149.
- Jarick, J., 2000, Die thermotektonometamorphe Entwicklung des Eastern Ghats Belt, Indien—ein Test der SWEAT-Hypothese: Unpubl. Ph.D. thesis, Johann Wolfgang Goethe-Universität, Frankfurt.
- Johansson, L., and Möller, C., 1986, Formation of sapphirine during retrogression of a basic high-pressure granulite, Roan, Western Gneiss Region, Norway: *Contributions to Mineralogy and Petrology*, v. 94, p. 29–41.
- Johnston, S. T., 2001, The great Alaskan terrane wreck: Reconciliation of paleomagnetic and geological data in the northern Cordillera: *Earth and Planetary Science Letters*, v. 193, p. 259–272.
- Johnson, T., and Brown, M., 2004, Quantitative constraints on metamorphism in the Variscides of southern Brittany—a complementary pseudosection approach: *Journal of Petrology*, v. 38, p. 1237–1259.
- John, T., Schenk, V., Mezger, K., and Tembo, F., 2004, Timing and P-T evolution of whiteschist metamorphism in the Lufillian Arc-Zambezi Belt Orogen

- (Zambia): Implications for the assembly of Gondwana: *Journal of Geology*, v. 112, p. 71–90.
- Kalt, A., Altherr, R., and Hanel, M., 2000, The Variscan basement of the Schwarzwald: *European Journal of Mineralogy*, v. 12, p. 1–43.
- Kaneko, Y., Katayama, I., Yamamoto, H., et al., 2003, Timing of Himalayan ultrahigh-pressure metamorphism: Sinking rate and subduction angle of the Indian continental crust beneath Asia: *Journal of Metamorphic Geology*, v. 21, p. 589–599.
- Kawakami, T., and Motoyoshi, Y., 2004, Timing of attainment of the spinel + quartz coexistence in garnet-sillimanite leucogneiss from Skallevikshalsen, Lützow-Holm Complex, east Antarctica: *Journal of Mineralogical and Petrological Sciences*, v. 99, p. 311–319.
- Kelly, N. M., and Harley, S. L., 2005, An integrated micro-textural and chemical approach to zircon geochronology: Refining the Archaean history of the Napier complex, east Antarctica: *Contributions to Mineralogy and Petrology*, v. 149, p. 57–84.
- Kelsey, D. E., Powell, R., Wilson, C. J. L., and Steel, D. A., 2003a, (Th + U)-Pb monazite ages from Al-Mg-rich metapelites, Rauer Group, east Antarctica: *Contributions to Mineralogy and Petrology*, v. 146, p. 326–340.
- Kelsey, D. E., White, R. W., Powell, R., Wilson, C. J. L., and Quinn, C. D., 2003b, New constraints on metamorphism in the Rauer Group, Prydz Bay, east Antarctica: *Journal of Metamorphic Geology*, v. 21, p. 739–759.
- Kelsey, D. E., White, R. W., Holland, T. J. B., and Powell, R., 2004, Calculated phase equilibria in  $K_2O$ -FeO-MgO-Al<sub>2</sub>O<sub>3</sub>-SiO<sub>2</sub>-H<sub>2</sub>O for sapphirine-quartz-bearing mineral assemblages: *Journal of Metamorphic Geology*, v. 22, p. 559–578.
- Kienast, J. R., and Ouzegane, K., 1987, Polymetamorphic Al-Mg rich parageneses in Archaean rocks from Hoggar, Algeria: *Geological Journal*, v. 22, p. 57–79.
- Klemd, R., and Bröcker, M., 1999, Fluid influence on mineral reactions in ultrahigh-pressure granulites: A case study in the Snieznik Mts. (West Sudetes, Poland): *Contributions to Mineralogy and Petrology*, v. 136, p. 358–373.
- Klemd, R., Bröcker, M., Hacker, B. R., Gao, J., Gans, P., and Wemmer, K., 2005, New age constraints on the metamorphic evolution of the high-pressure/low-temperature belt in the western Tianshan Mountains, NW China: *Journal of Geology*, v. 113, p. 157–168.
- Klemd, R., Schröter, F. C., Will, T. M., and Gao, J., 2002, P-T evolution of glaucophane-omphacite bearing HP-LT rocks in the western Tianshan Orogen, NW China: New evidence for “Alpine-type” tectonics: *Journal of Metamorphic Geology*, v. 20, p. 239–254.
- Komiya, T., Hayashi, M., Maruyama, S., and Yurimoto, H., 2002, Intermediate-P/T type Archean metamorphism of the Isua Supracrustal Belt: Implications for secular change of geothermal gradients at subduction zones and for Archean plate tectonics: *American Journal of Science*, v. 302, p. 806–826.
- Korenaga, J., 2006, Archean geodynamics and the thermal evolution of Earth, *in* Benn, K., Mareschal, J.-C., and Condie, K. C., eds., *Archean geodynamics and environments*: Washington, DC, American Geophysical Union, Geophysical Monograph 164, p. 7–32.
- Korhonen, F. J., and Stout, J. H., 2004, Low-variance sapphirine-bearing assemblages from Wilson Lake, Grenville Province of Labrador, *in* Tollo, R. P., Corriveau, L., McLelland, J., and Bartholomew, M. J., eds., *Proterozoic tectonic evolution of the Grenville Orogen in North America*: Geological Society of America Memoir 197, p. 81–103.
- Kotková, J., Kröner, A., Todt, W., and Fiala, J., 1996, Zircon dating of North Bohemian granulites, Czech Republic: Further evidence for the Lower Carboniferous high-pressure event in the Bohemian Massif: *Geologische Rundschau*, v. 85, p. 154–161.
- Kramers, J. D., Kreissig, K., and Jones, M. Q. W., 2001, Crustal heat production and style of metamorphism: A comparison between two Archean high-grade provinces in the Limpopo belt, Southern Africa: *Precambrian Research*, v. 112, p. 149–163.
- Kreissig, K., Holzer, L., Frei, R., Villa, I. M., Kramers, J. D., Kröner, A., Smit, C. A., and van Reenen, D. D., 2000, Geochronology of the Hout River shear zone and the metamorphism in the Southern Marginal Zone of the Limpopo Belt, southern Africa: *Precambrian Research*, v. 109, p. 145–173.
- Kriegsman, L. O., and Schumacher, J. C., 1999, Petrology of sapphirine-bearing and associated granulites from central Sri Lanka: *Journal of Petrology*, v. 40, p. 1211–1239.
- Kröner, A., Cui, W. Y., Wang, S. Q., Wang, C. Q., and Nemchin, A. A., 1998a, Single zircon ages from high-grade rocks of the Jianping Complex, Liaoning Province, NE China: *Journal of Asian Earth Sciences*, v. 16, p. 519–532.
- Kröner, A., Jaeckel, P., Brandl, G., Nemchin, A. A., and Pidgeon, R. T., 1999, Single zircon ages for granitoid gneisses in the central zone of the Limpopo belt, southern Africa and geodynamic significance: *Precambrian Research*, v. 93, p. 299–337.
- Kröner, A., Jaeckel, P., Reischmann, T., and Kroner, U., 1998b, Further evidence for an early Carboniferous (similar to 340 Ma) age of high-grade metamorphism in the Saxonian granulite complex: *Geologische Rundschau*, v. 86, p. 751–766.
- Kröner, A., O'Brien, P. J., Nemchin, A. A., and Pidgeon, R. T., 2000, Zircon ages for high pressure granulites from South Bohemia, Czech Republic, and their connection to Carboniferous high temperature processes: *Contributions to Mineralogy and Petrology*, v. 138, p. 127–142.
- Kröner, A., and Willner, A. P., 1998, Time of formation and peak of Variscan HP-HT metamorphism of quartz-

- feldspar rocks in the central Erzgebirge, Saxony, Germany: Contributions to Mineralogy and Petrology, v. 132, p. 1–20.
- Kryza, R., Pin, C., and Vielzeuf, D., 1996, High-pressure granulites from the Sudetes (south-west Poland): Evidence of crustal subduction and collisional thickening in the Variscan Belt: Journal of Metamorphic Geology, v. 14, p. 531–546.
- Kühn, A., Glodny, J., and Ring, U., 2005, Rapid Oligocene exhumation of the eclogite zone, Tauern Window, Eastern Alps: Mitteilungen der Österreichischen Mineralogischen Gesellschaft, v. 150, p. 80.
- Lardeaux, J. M., Ledru, P., Daniel, I., and Duchene, S., 2001, The Variscan French Massif Central—a new addition to the ultrahigh pressure metamorphic “club”: Exhumation processes and geodynamic consequences: Tectonophysics, v. 332, p. 143–167.
- Leech, M. L., Metzger, E. P., Wooden, J. L., Jones, R. E., Schwartz, C. L., and Beane, R. J., 2002, New Eclogitization and protolith ages for the Maksyutov Complex (south Ural Mountains) based on U-Pb zircon SHRIMP data [abs.]: EOS (Transactions of the American Geophysical Union), v. 83, abstract V51B-1263.
- Leech, M. L., Singh, S., Jain, A. K., Klemperer, S. L., and Manickavasagam, R. M., 2005, The onset of India-Asia continental collision: Early, steep subduction required by the timing of UHP metamorphism in the western Himalaya: Earth and Planetary Science Letters, v. 234, p. 83–97.
- Leggo, P. J., Aftalion, M., and Pidgeon, R. T., 1971, Discordant zircon U-Pb ages from Uganda basement: Nature-Physical Science, v. 231, p. 81–84.
- Le Hébel, F., Vidal, O., and Kienast, J. R., et al., 2002, Evidence for HP-LT Hercynian metamorphism within the “Porphyroides” of south Brittany, France: Comptes Rendus Geoscience, v. 334, p. 205–211.
- Leong, K. M., and Moore, J. M., 1972, Sapphirine-bearing rocks from Wilson Lake, Labrador: Canadian Mineralogist, v. 11, p. 777–790.
- Li, X.-P., Zheng, Y.-F., Wu, Y.-B., Chen, F., Gong, B., and Li, Y.-L., 2004, Low-T eclogite in the Dabie Terrane of China: Petrological and isotopic constraints on fluid activity and radiometric dating: Contributions to Mineralogy and Petrology, v. 148, p. 443–470.
- Liati, A., and Gebauer, D., 2003, Geochronological constraints of the time of metamorphism in the Gruf Complex (central Alps) and implications for the Adula-Cima Lunga Nappe system: Schweizerische Mineralogische und Petrographische Mitteilungen, v. 83, 159–172.
- Liati, A., Gebauer, D., and Wysoczanski, R., 2002, U-Pb SHRIMP-dating of zircon domains from UHP mafic rocks in the Rhodope zone (N. Greece): Evidence for early Cretaceous crystallization and late Cretaceous metamorphism: Chemical Geology, v. 184, p. 281–300.
- Libourel, G., 1988, Le Complexe de Santa-Lucia di Mercurio (Corse): Un nouveau jalon de la base de la croûte varisque en Méditerranée occidentale: Compte Rendus de l'Académie des Sciences, Paris, v. 307, p. 1067–1073.
- Liou, J. G., Tsujimori, T., Zhang, R. Y., Katayama, I., and Maruyama, S., 2004, Global UHP metamorphism and continental subduction/collision: The Himalayan model: International Geology Review, v. 46, p. 1–27.
- Liu, F., Xu, Z., Liou, J. G., Katayama, I., Masago, H., Maruyama, S., and Yang, J., 2002, Ultrahigh-pressure mineral inclusions in zircons from gneissic core samples of the Chinese continental scientific drilling site in eastern China: European Journal of Mineralogy, v. 14, p. 499–512.
- Liu, F., Zhang, Z., Katayama, I., Xu, Z., and Maruyama, S., 2003, Ultrahigh-pressure metamorphic records hidden in zircons from amphibolites in Sulu terrane, eastern China: The Island Arc, v. 12, p. 256–267.
- Liu, F. L., Xu, Z. Q., and Liou, J. G., 2004a, Tracing the boundary between UHP and UP metamorphic belts in the southwestern Sulu terrane, Eastern China: Evidence from mineral inclusions in zircons from metamorphic rocks: International Geology Review, v. 46, p. 409–425.
- Liu, F. L., Xu, Z. Q., and Xue, H. M., 2004b, Tracing the protolith, UHP metamorphism, and exhumation ages of orthogneiss from the SW Sulu terrane (eastern China): SHRIMP U-Pb dating of mineral inclusion-bearing zircons: Lithos, v. 78, p. 411–429.
- Liu, L., Sun, Y., Chen, D., Zhang, A., and Luo, J., 2003a, Discovery of relic majoritic garnet in felsic metamorphic rocks of Qinling complex, north Qinling orogenic belt, China: Alice Wain Memorial Western Norway Eclogite Field Symposium, Selje, Western Norway, Abstract Volume, p. 82.
- Liu, L., Sun, Y., Wang, Y., Chen, D. L., Luo, J. H., and Zhang, A. D., 2003b, Ultrahigh-P evidence for country rocks of the garnet-bearing lherzolite in Altyn Tagh—exsolution of clinopyroxene in the garnet of retrograde eclogite [abs.]: Alice Wain Memorial Western Norway Eclogite Field Symposium, Selje, Norway, Abstract Volume, p. 81.
- Liu, L., Wang, Y., Sun, Y., Xiao, P., Chen, D., Luo, J., and Che, Z., 2001, The discovery and significance of magnetite-bearing garnet lherzolite from Altun Eclogite Zone, Western China [abs.]: Sixth International Eclogite Conference, Sept. 1–7, Niihama, Japan, Abstract, p. 79–80.
- Liu, L., Zhang, J., Green, H.W., and Jin, Z., 2005, Evidence of former stishovite in metamorphosed sediments: Exhumation from >300 km: EOS (Transactions of the American Geophysical Union), v. 86, abstract V51E-08.
- Loewy, S. L., Connelly, J. N., and Dalziel, I. W. D., 2004, An orphaned basement block: The Aarequipa-Antofalla basement of the central Andean margin of South America: Geological Society of America Bulletin 116, p. 171–187.

- Love, G. J., Kinny, P. D., and Friend, C. R. L., 2004, Timing of magmatism and metamorphism in the Gruinard Bay area of the Lewisian gneiss complex: Comparisons with the Assynt terrane and implications for terrane accretion: *Contributions to Mineralogy and Petrology*, v. 146, p. 620–636.
- Mancktelow, N. S., 1995, Non-lithostatic pressure during sediment subduction and the development and exhumation of high pressure metamorphic rocks: *Journal of Geophysical Research*, v. 100, p. 571–583.
- Mann, P., and Taira, A., 2004, Global tectonic significance of the Solomon Islands and Ontong Java Plateau convergent zone: *Tectonics*, v. 389, p. 137–190.
- Marques, F. O., Ribeiro, A., and Munhá, J. M., 1996, Geodynamic evolution of the Continental Allochthonous Terrane (CAT) of the Bragança Nappe Complex, NE Portugal: *Tectonics*, v. 15, p. 747–762.
- Marshak, S., 1999, Deformation style way back when: Thoughts on the contrasts between Archean/Paleoproterozoic and contemporary orogens: *Journal of Structural Geology*, v. 21, p. 1175–1182.
- Martignole, J., and Martelat, J.-E., 2003, Regional-scale Grenvillian-age UHT metamorphism in the Mollendo-Camana block (basement of the Peruvian Andes): *Journal of Metamorphic Geology*, v. 21, 99–120.
- Maruyama, S., 1997, Pacific-type orogeny revisited: Miyashiro-type orogeny proposed: *Island Arc*, v. 6, p. 91–120.
- Maruyama, S., and Liou, J. G., 1998, Initiation of ultrahigh-pressure metamorphism and its significance on the Proterozoic–Phanerozoic boundary: *The Island Arc*, v. 7, p. 6–35.
- Maruyama, S., and Liou, J. G., 2005, From snowball to Phanerozoic Earth: *International Geology Review*, v. 47, p. 775–791.
- Maruyama, S., Liou, J. G., and Terabayashi, M., 1996, Blueschists and eclogites of the world and their exhumation: *International Geology Review*, v. 38, p. 490–596.
- Mason, A. J., Parrish, R. R., and Brewer, T. S., 2004, U-Pb geochronology of Lewisian orthogneiss in the outer Hebrides, Scotland: Implications for the tectonic setting and correlation of the South Harris complex: *Journal of the Geological Society*, v. 161, p. 45–54.
- Massonne, H.-J., 1999, A new occurrence of microdiamonds in quartzo-feldspathic rocks of the Saxonian Erzgebirge, Germany, and their metamorphic evolution, in Gurney, J. J., Gurney, L. G., Pascoe, M. D., and Richardson, S. H., eds., *Proceedings of the 7th International Kimberlite Conference*, Cape Town: Cape Town, South Africa, Redroof Publishing, v. 2 (The Nixon Volume), p. 533–539.
- Massonne, H.-J., 2001, First find of coesite in the ultrahigh-pressure metamorphic area of central Erzgebirge, Germany: *European Journal of Mineralogy*, v. 13, p. 565–570.
- Massonne, H.-J., and Brandelik, 2005, Calculated phase relations in the system KFMASH between 6 and 16 GPa [abs.]: EOS (Transactions of the American Geophysical Union), v. 86, abstract V51E-02.
- Matsuda, T., and Uyeda, S., 1971, On the Pacific-type orogeny and its model—extension of the paired belts concept and possible origin of marginal seas: *Tectonophysics*, v. 11, p. 5–27.
- Mattinson, C. G., Zhang, R. Y., Tsujimori, T., and Liou, J. G., 2004, Epidote-rich calc-kyanite-phengite eclogites, Sulu Terrane, eastern China: P-T- $f_{O_2}$  estimates and the significance of the epidote-talc assemblage in eclogite: *American Mineralogist*, v. 89, p. 1772–1783.
- Mattinson, C. G., Liou, J. G., Bird, D. K., Wooden, J. L., Wu, C., and Yang, J., 2005a, Zircon geochronology and REE geochemistry, north Qaidam UHP Terrane, north-west China: *Mitteilungen der Österreichischen Mineralogischen Gesellschaft*, v. 150, p. 106.
- Mattinson, C. G., Liou, J. G., Bird, D. K., Wooden, J. L., Wu, C., and Yang, J., 2005b, Geochronology of prograde and eclogite-facies zircon growth, North Qaidam UHP terrane, NW China [abs.]: EOS (Transactions of the American Geophysical Union), v. 86, abstract V43A-1556.
- McClelland, W. C., Gilotti, J. A., Power, S. E., and Mazdab, F. K., 2005, Dating of UHP metamorphism, NE Greenland Caledonides: *Mitteilungen der Österreichischen Mineralogischen Gesellschaft*, v. 150, p. 107.
- McClelland, W. C., Power, S. E., Gilotti, J. A., Mazdab, F. K., and Wopenka, B., 2006, U-Pb SHRIMP geochronology and trace-element geochemistry of coesite-bearing zircons, north-east Greenland Caledonides, in Hacker, B. R., McClelland, W. C., and Liou, J. G., eds., *Ultrahigh-pressure metamorphism: Deep continental subduction: Geological Society of America Special Paper*, 403, p. 23–43.
- McGregor, V. R., and Friend, C. R. L., 1997, Field recognition of rocks totally retrogressed from granulite facies: An example from Archean rocks in the Paamiut region, south-west Greenland: *Precambrian Research*, v. 86, p. 59–70.
- Medaris, L. G., Beard, B. L., Johnson, C. M., Valley, J. W., Spicuzza, M. J., Jelinek, E., and Misar, Z., 1995a, Garnet pyroxenite and eclogite in the Bohemian Massif: geochemical evidence for Variscan recycling of subducted lithosphere: *Geologische Rundschau*, v. 84, p. 489–505.
- Medaris, G., Ducea, M., Ghent, E., and Iancu, V., 2003, Conditions and timing of high-pressure Variscan metamorphism in the South Carpathians, Romania: *Lithos*, v. 70, p. 141–161.
- Medaris, L. G., Jelinek, E., and Misař, Z., 1995b, Czech eclogites: Terrane settings and implications for Variscan tectonic evolution of the Bohemian Massif: *European Journal of Mineralogy*, v. 7, p. 7–28.
- Miyashiro, A., 1961, Evolution of metamorphic belts: *Journal of Petrology*, v. 2, p. 277–311.

- Miyashiro, A., 1973, Paired and unpaired metamorphic belts: *Tectonophysics*, v. 17, p. 241–254.
- Miyazaki, K., Zulkarnain, I., Sopaheluwakan, J., and Wakita, K., 1996, Pressure-temperature conditions and retrograde paths of eclogites, garnet-glaucophane rocks and schists from south Sulawesi, Indonesia: *Journal of Metamorphic Geology*, v. 14, 549–563.
- Mohan, A., Ackermann, D., and Lal, R. K., 1986, Reaction textures and P-T-X trajectory in the sapphirine-spinel bearing granulites from Ganguvarpatti, Southern India: *Neues Jahrbuch für Mineralogie Abhandlungen*, v. 154, p. 1–19.
- Möller, A., Appel, P., Mezger, K., and Schenk, V., 1995, Evidence for a 2 Ga subduction zone: Eclogites in the Usagaran belt of Tanzania: *Geology*, v. 23, p. 1067–1070.
- Möller, C., 1988, Geology and metamorphic evolution of the Roan area, Vestranden, Western Gneiss Region, Central Norwegian Caledonides: *Norges Geologiske Undersøkelse*, v. 413, p. 1–31.
- Möller, C., 1998, Decompressed eclogites in the Sveconorwegian (-Grenvillian) orogen of SW Sweden: petrology and tectonic implications: *Journal of Metamorphic Geology*, v. 16, p. 641–656.
- Möller, C., 1999, Sapphirine in SW Sweden: A record of Sveconorwegian (-Grenvillian) late-orogenic tectonic exhumation: *Journal of Metamorphic Geology*, v. 17, p. 127–141.
- Monger, J. W. H., 1997, Plate tectonics and northern Cordilleran geology: An unfinished revolution: *Geoscience Canada*, v. 24, p. 189–198.
- Moore, E. M., 2002, Pre-1 Ga (pre-Rodinian) ophiolites: Their tectonic and environmental implications: *GSA Bulletin*, v. 114, p. 80–95.
- Moraes, R., Brown, M., Fuck, R. A., Camargo, M. A., and Lima, T. M., 2002, Characterization and P-T evolution of melt-bearing ultrahigh-temperature granulites: An example from the Anápolis-Itaçu Complex of the Brasília Fold Belt, Brazil: *Journal of Petrology*, v. 43, p. 1673–1705.
- Moraes, R., and Fuck, R. A., 2000, Ultra-high-temperature metamorphism in central Brazil: The Barro Alto complex: *Journal of Metamorphic Geology*, v. 18, p. 345–358.
- Moraes, R., Fuck, R. A., and Brown, M., 2003, Qual é a Fonte de Calor Para Os Grãulitos de Temperatura Ultra-Alta? O Exemplo do Complexo Anápolis Itaçu [abs.], in IX Simpósio Nacional de Estudos Tectônicos, Boletim de Resumos, p. 70–71.
- Morimoto, T., Santosh, M., Tsunogae, T., and Yoshimura, Y., 2004, Spinel + quartz association from the Kerala khondalites, southern India: Evidence for ultrahigh-temperature metamorphism: *Journal of Mineralogical and Petrological Sciences*, v. 99, p. 257–278.
- Morse, S. A., and Talley, J. H., 1971, Sapphirine reactions in deep seated granulites near Wilson Lake, Central Labrador, Canada: *Earth and Planetary Science Letters*, v. 10, p. 325–328.
- Motoyoshi, Y., and Ishikawa, M., 1997, Metamorphic and structural evolution of granulites from Rundvågshetta, Lützow-Holm Bay, east Antarctica, in Ricci, C. A., ed., *The Antarctic region: Geological evolution and processes: Proceedings of the VII International Symposium on the Antarctic Earth Sciences*, Siena, Terra Antarctica, p. 65–72.
- Mouri, H., Andreoli, M.G.G., Kienast, J.R., Guiraud, M., and De Waal, S.A., 2003, First occurrence of the rare 'corundum + quartz' assemblage in the high-grade zone from the Namaqualand Metamorphic Complex, South Africa: evidence for higher P, T metamorphism? *Mineralogical Magazine*, v. 67, p. 1015–1021.
- Mposkos, E. D., and Kostopoulos, D. K., 2001, Diamond, former coesite, and supersilicic garnet in metasedimentary rocks from the Greek Rhodope: A new ultra-high-pressure metamorphic province established: *Earth and Planetary Science Letters*, v. 192, p. 497–506.
- Muhongo, S., Tuisku, P., Mnali, S., Temu, E., Appel, P., and Stendal, H., 2002, High-pressure granulite-facies metagabbros in the Ubendian Belt of SW Tanzania: Preliminary petrography and P-T estimates: *Journal of African Earth Sciences*, v. 34, p. 279–285.
- Mukherjee, B. K., and Sachan, H. K., 2001, Discovery of coesite from Indian Himalaya: A record of ultra-high pressure metamorphism in Indian continental crust: *Current Science*, v. 81, p. 1358–1361.
- Mukherjee, B. K., Sachan, H. K., Ogasawara, Y., Muko, A., and Yoshioka, N., 2003, Carbonate-bearing UHPM rocks from the Tso-Morari region, Ladakh, India: Petrological implications: *International Geology Review*, v. 45, p. 49–69.
- Müller, W., 2003, Strengthening the link between geochronology, textures, and petrology: *Earth and Planetary Science Letters*, v. 206, p. 237–251.
- Murphy, J. B., and Nance, R. D., 1991, Supercontinent model for the contrasting character of Late Proterozoic orogenic belts: *Geology*, v. 19, p. 469–472.
- Nandakumar, V., and Harley, S.L., 2000, A reappraisal of the pressure-temperature path of granulites from the Kerala Khondalite Belt, southern India: *Journal of Geology*, v. 108, p. 687–703.
- Nakamura, D., Svojtka, M., and Naemura, K., 2004, Very high-pressure (> 4 GPa) eclogite associated with the Moldanubian Zone garnet peridotite (Nove Dvory, Czech Republic): *Journal of Metamorphic Geology*, v. 22, p. 593–603.
- Nakano, N., Osanai, Y., Owada, M., Nam, T. N., Tsunogae, T., Toyoshima, T., and Binh, P., 2004, Decompression process of mafic granulite from eclogite to granulite facies under ultrahigh-temperature condition in the Kontum Massif, central Viet Nam: *Journal of Mineralogical and Petrological Sciences*, v. 99, p. 242–256.

- Nam, T. N., Sano, Y., Terada, K., Toriumi, M., Quynh, P. V., and Dung, L. T., 2001, First SHRIMP U-Pb zircon dating of granulites from the Kontum Massif (Viet Nam) and tectonothermal implications: *Journal of Asian Earth Sciences*, v. 19, p. 77–84.
- Nasdala, L., and Massonne, H.-J., 2000, Microdiamonds from the Saxonian Erzgebirge, Germany: *In situ* micro-Raman characterisation: *European Journal of Mineralogy*, v. 12, p. 495–498.
- Nelson, K. D., Zhao, W. J., Brown, L. D., Kuo, J., Che, J. K., Liu, X. W., Klemperer, S. L., Makovsky, Y., Meissner, R., Mechie, J., Kind, R., Wenzel, F., Ni, J., Nabelek, J., Chen, L. S., Tan, H. D., Wei, W. B., Jones, A. G., Booker, J., Unsworth, M., Kidd, W. S. F., Hauck, M., Alsdorf, D., Ross, A., Cogan, M., Wu, C. D., Sandvol, E., and Edwards, M., 1996, Partially molten middle crust beneath southern Tibet: Synthesis of project INDEPTH results: *Science*, v. 274, p. 1684–1688.
- Nemchin, A. A., Pidgeon, R. T., and Whitehouse, M. J., 2006, Re-evaluation of the origin and evolution of >4.2 Ga zircons from the Jack Hills metasedimentary rocks: *Earth and Planetary Science Letters* [doi:10.1016/j.epsl.2006.01.054].
- Nur, A., and Ben-Avraham, Z., 1989, Oceanic plateaus and the Pacific Ocean margins, in Ben-Avraham, Z., ed., *The evolution of the Pacific Ocean margins*: Oxford, UK, Oxford University Press, p. 7–19.
- O'Brien, P. J., 1997, Garnet zoning and reaction textures in overprinted eclogites, Bohemian Massif, European Variscides: A record of their thermal history during exhumation: *Lithos*, v. 41, p. 119–133.
- O'Brien, P. J., 1999, Asymmetric zoning profiles in garnet from HP-HT granulite and implications for volume and grain boundary diffusion: *Mineralogical Magazine*, v. 63, p. 227–238.
- O'Brien, P. J., and Rötzler, J., 2003, High-pressure granulites: Formation, recovery of peak conditions, and implications for tectonics: *Journal of Metamorphic Geology*, v. 21, p. 3–20.
- O'Brien, P. J., Walte, N., and Li, Jianghai, 2005, The petrology of two distinct granulite types in the Hengshan Mts., China, and tectonic implications: *Journal of Asian Earth Sciences*, v. 24, p. 615–627.
- O'Brien, P. J., Zotov, N., Law, R. D., Khan, M. A., and Jan, M. Q., 2001, Coesite in Himalayan eclogite and implications for models of India-Asia collision: *Geology*, v. 29, p. 435–438.
- Och, D. J., Leitch, E. C., Caprarelli, G., and Watanabe, T., 2003, Blueschist and eclogite in tectonic melange, Port Macquarie, New South Wales, Australia: *Mineralogical Magazine*, v. 67, p. 609–624.
- Okay, A. I., 1993, Petrology of a diamond and coesite-bearing terrain: Dabie Shan, China: *European Journal of Mineralogy*, v. 5, p. 659–675.
- Okay, A. I., 2002, Jadeite-chloritoid-glaucophane-lawsonite blueschists in north-west Turkey: Unusually high P/T ratios in continental crust: *Journal of Metamorphic Geology*, v. 20, p. 757–768.
- Okay, A. I., Harris, N. B. W., and Kelley, S. P., 1998, Exhumation of blueschists along a Tethyan Suture in northwest Turkey: *Tectonophysics*, v. 285, p. 275–299.
- O'Neill, C., Lenardic, A., Moresi, L., Torsvik, T., and Lee, C. A., 2005, Episodic Precambrian subduction [abs.]: EOS (Transactions of the American Geophysical Union), v. 86, abstract U43B-0835.
- Osanai, Y., Nakano, N., Owada, M., Nam, T. N., Toyoshima, T., Tsunogae, T., and Binh, P., 2004, Permo-Triassic ultrahigh-temperature metamorphism in the Kontum Massif, central Viet Nam: *Journal of Mineralogical and Petrological Sciences*, v. 99, p. 225–241.
- Osanai, Y., Sajeev, K., Owada, M., Kehelpannala, K. V. W., Prame, W. K. B. N., and Nakano, N., 2003, Evolution of highest-grade metamorphic rocks from Central Highland Complex, Sri Lanka: Geological Survey and Mines Bureau, Sri Lanka, Centenary Publication, p. 25–31.
- Ota, T., Gladkochub, D. P., Sklyarov, E. V., Mazukabzov, A. M., and Watanabe, T., 2004a, P-T history of garnet-websterites in the Sharyzhgalskiy Complex, southwestern margin of the Siberian Craton: Evidence for Paleoproterozoic high-pressure metamorphism: *Precambrian Research*, v. 132, p. 327–348.
- Ota, T., Terabayashi, M., and Katayama, I., 2004b, Thermobaric structure and metamorphic evolution of the Iratsu eclogite body in the Sanbagawa belt, central Shikoku, Japan: *Lithos*, v. 73, p. 95–126.
- Ouzegane, K., and Boumaza, S., 1996, An example of ultrahigh-temperature metamorphism: Orthopyroxene-sillimanite-garnet, sapphirine-quartz and spinel-quartz parageneses in Al-Mg granulites from In Hihau, In Ouzal, Hoggar: *Journal of Metamorphic Geology*, v. 14, p. 693–708.
- Ouzegane, K., Guiraud, M., and Kienast, J. R., 2003, Prograde and retrograde evolution in high-temperature corundum granulites (FMAS and KFMASH systems) from In Ouzal Terrane (NW Hoggar, Algeria): *Journal of Petrology*, v. 44, p. 517–545.
- Palmeri, R., Ghiribelli, B., Talarico, F., and Ricci, C. A., 2003, Ultra-high-pressure metamorphism in felsic rocks: The garnet-phengite gneisses and quartzites from the Lanterman Range, Antarctica: *European Journal of Mineralogy*, v. 15, p. 513–525.
- Paquette, J.-L., Goncalves, P., Devouard, B., and Nicollet, C., 2004, Micro-drilling ID-TIMS U-Pb dating of single monazites: A new method to unravel complex poly-metamorphic evolution. Application to the UHT granulites of Andriamena (North-Central Madagascar): *Contributions to Mineralogy and Petrology*, v. 147, p. 110–122.
- Parkinson, C. D., Miyazaki, K., Wakita, K., Barber, A. J., and Carswell, D. A., 1998, An overview and tectonic synthesis of the pre-Tertiary very-high-pressure meta-

- morphic and associated rocks of Java, Sulawesi, and Kalimantan, Indonesia: *The Island Arc*, v. 7, p. 184–200.
- Parkinson, C. D., Motoki, A., Onishi, C. E., and Maruyama, S., 2001, Ultrahigh-pressure pyrope-kyanite granulites and associated eclogites in Neoproterozoic nappes of Southeast Brazil [abs.], in UHPM Workshop 2001, Fluid/slab/mantle interactions and ultrahigh-P minerals, Waseda University, Tokyo, Abstract Volume, p. 87–90.
- Parrish, R. R., Gough, S., Searle, M., and Waters, D., 2003, Exceptionally rapid exhumation of the Kaghan UHP terrane, Pakistan from U-Th-Pb measurements on accessory minerals [abs.]: *Geological Society of America, Abstracts with Programs*, v. 35, p. 556.
- Pattison, D. R. M., and Bégin, N. J., 1994, Hierarchy of closure temperatures in granulites and the importance of an intergranular exchange medium (melt?) in controlling maximum Fe–Mg exchange temperatures: *Mineralogical Magazine*, v. 58A, p. 694–695.
- Pattison, D. R. M., Chacko, T., Farquhar, J., and McFarlane, C. R. M., 2003, Temperatures of granulite-facies metamorphism: Constraints from experimental phase equilibria and thermobarometry corrected for retrograde exchange: *Journal of Petrology*, v. 44, p. 867–900.
- Perchuk, L. L., Aranovich, L. Ya., Podlesskii, K. K., Lavrent'eva, I. V., Gerasimov, V. V., Kitsul, V. I., Korsakov, L. P., and Berdnikov, N. V., 1985, Precambrian granulites of the Aldan Shield, eastern Siberia, the USSR: *Journal of Metamorphic Geology*, v. 3, p. 265–310.
- Percival, J. A., 1991, Granulite-facies metamorphism and crustal magmatism in the Ashuanipi Complex, Quebec-Labrador, Canada: *Journal of Petrology*, v. 32, p. 1261–1297.
- Percival, J. A., and McGrath, P. H., 1986, Deep crustal history and tectonic history of the northern Kapuskasing uplift of Ontario: An integrated petrological-geophysical study: *Tectonics*, v. 5, p. 553–572.
- Perkins, D., III, and Chipera, S. J., 1985, Garnet-orthopyroxene-plagioclase-quartz barometry: Refinement and application to the English River subprovince and the Minnesota River Valley: *Contributions to Mineralogy and Petrology*, v. 89, p. 69–80.
- Petrini, K., and Podladchikov, Y., 2000, Lithospheric pressure-depth relationship in compressive regions of thickened crust: *Journal of Metamorphic Geology*, v. 18, p. 67–77.
- Pimentel, M. M., Ferreira Filho, C. F., and Armstrong, R. A., 2004, SHRIMP U-Pb and Sm-Nd ages of the Niquelândia layered complex: Meso- (1.25 Ga) and Neoproterozoic (0.79 Ga) extensional events in Central Brazil: *Precambrian Research*, v. 132, p. 133–153.
- Pin, C., and Vielzeuf, D., 1983, Granulites and related rocks in Variscan median Europe: A dualistic interpretation: *Tectonophysics*, v. 93, p. 47–74.
- Pin, C., and Vielzeuf, D., 1988, Les granulites de haute pression d'Europe moyenne témoins d'une subduction éo-hercynienne. Implications sur l'origine des groupes leptyno-amphiboliques: *Bulletin de la Société Géologique de France*, v. 8, no. 4, p. 13–20.
- Piuzana, D., Pimentel, M. M., Fuck, R. A., and Armstrong, R., 2003, Neoproterozoic granulite facies metamorphism and coeval granitic magmatism in the Brasília belt, Central Brazil: Regional implications of new SHRIMP U-Pb and Sm-Nd data: *Precambrian Research*, v. 125, p. 245–274.
- Pollack, H. N., 1997, Thermal characteristics of the Archaean, in De Wit, M. J., and Ashwal, L. D., eds., *Oxford monographs on geology and geophysics*. Oxford, UK and New York, NY: Clarendon Press, v. 35, p. 223–232.
- Potel, S., Ferreiro Mähmann, R., Stern, W. B., Mullis, J., and Frey, M., 2006, Very low-grade metamorphic evolution of pelitic rocks under high-pressure/low-temperature conditions, NW New Caledonia, (SW Pacific): *Journal of Petrology* [doi:10.1093/petrology/egl001].
- Pouba, Z., Padera, K., and Fiala, J., 1985, Omphacite granulite from the NE marginal area of the Bohemian Massif, (Rychleby Mts.): *Neues Jahrbuch für Mineralogie, Abhandlungen*, v. 151, p. 29–52.
- Powell, R., Guiraud, M., and White, R. W., 2005, Truth and beauty in metamorphic phase equilibria: Conjugate variables and phase diagrams: *Canadian Mineralogist*, v. 43, p. 21–33.
- Powell, R., and Holland, T., 1994, Optimal geothermometry and geobarometry: *American Mineralogist*, v. 79, p. 120–133.
- Powell, R., and Holland, T., 1999, Relating formulations of the thermodynamics of mineral solid solutions: Activity modeling of pyroxenes, amphiboles, and micas: *American Mineralogist*, v. 84, p. 1–14.
- Powell, R., Holland, T., and Worley, B., 1998, Calculating phase diagrams involving solid solutions via non-linear equations, with examples using THERMOCALC: *Journal of Metamorphic Geology*, v. 16, p. 577–588.
- Raith, M., Karmakar, S., and Brown, M., 1997, Ultra-high temperature metamorphism and multistage decompressional evolution of sapphirine granulites from the Palni Hill Ranges, southern India: *Journal of Metamorphic Geology*, v. 15, 379–400.
- Raith, M., Srikantappa, C., Ashamanjari, K. G., and Spiering, B., 1990, The granulite terrane of the Nilgiri Hills (southern India): Characterization of high grade metamorphism, in Vielzeuf, D., and Vidal, Ph., eds., *Granulites and crustal evolution*: Dordrecht, The Netherlands: Kluwer Academic, p. 339–365.

- Ravna, E. J. Krogh, 1982, Metamorphic evolution of Norwegian country-rock eclogites, as deduced from mineral inclusions and compositional zoning in garnets: *Lithos*, v. 15, p. 305–321.
- Ravna, E. J. Krogh, Oh, C. W., and Liou, J. G., 1994, Polyphase and anticlockwise P-T evolution for Franciscan eclogites and blueschists from Jenner, California, USA: *Journal of Metamorphic Geology*, v. 12, 121–134.
- Ravna, E. J. Krogh, and Roux, M. R. M., 2006, Metamorphic evolution of the Tønsvika eclogite, Tromsø Nappe—evidence for a new UHP province in the Scandinavian Caledonides: *International Geology Review*, v. 48, p. 861–881.
- Ravna, E. J. Krogh, and Terry, M. P., 2004, Geothermobarometry of UHP and HP eclogites and schists—an evaluation of equilibria among garnet-clinopyroxene-kyanite-phengite-coesite/quartz: *Journal of Metamorphic Geology*, v. 22, p. 570–592.
- Reinecke, T., 1991, Very-high-pressure metamorphism and uplift of coesite bearing metasediments from the Zermatt-Saas zone, Western Alps: *European Journal of Mineralogy*, v. 3, p. 7–17.
- Root, D. B., Hacker, B. R., Mattinson, J. M., and Wooden, J. L., 2004, Zircon geochronology and ca. 400 Ma exhumation of Norwegian ultrahigh-pressure rocks: an ion microprobe and chemical abrasion study: *Earth and Planetary Science Letters*, v. 228, p. 325–341.
- Rötzler, J., 1992, Zur Petrogenese im Sächsischen Granulitgebirge: die pyroxenfreien Granulite und die Metapelite: *Geotektonische Forschungen*, v. 77, p. 1–100.
- Rötzler, J., and Romer, R. L., 2001, P-T-t evolution of ultrahigh-temperature granulites from the Saxon Granulite Massif, Germany. Part I: Petrology: *Journal of Petrology*, v. 42, p. 1995–2013.
- Rubatto, D., 2002, Zircon trace element geochemistry: Partitioning with garnet and the link between U-Pb ages and metamorphism: *Chemical Geology*, v. 184, p. 123–138.
- Rubatto, D., Korsakov, A., Hermann, J., and Dobretsov, N. L., 2005, Diachronous UHP metamorphism in the Kokchetav Massif: *Mitteilungen der Österreichischen Mineralogischen Gesellschaft*, v. 150, p. 135.
- Săbău, G., 2000, A possible UHP-eclogite in the Leaota Mts. (South Carpathians) and its history from high-pressure melting to retrograde inclusion in a subducting melange: *Lithos*, v. 52, p. 253–276.
- Sachan, H. K., Mukherjee, B. K., Ogasawara, Y., Maruyama, S., Ishida, H., Muko, A., and Yoshioka, N., 2004, Discovery of coesite from Indus Suture Zone (ISZ), Ladakh, India: Evidence for deep subduction: *European Journal of Mineralogy*, v. 16, p. 235–240.
- Sajeev, K., and Osanai, Y., 2004, Ultrahigh-temperature metamorphism (1150°C, 12 kbar) and multistage evolution of Mg-, Al-rich granulites from the Central Highland Complex, Sri Lanka: *Journal of Petrology*, v. 45, p. 1821–1844.
- Sal'nikova, E. B., Glebovitsky, V. A., Kotov, A. B., Larin, A. N., Yakovleva, S. Z., Berezhnaya, N. G., Kovach, V. P., Drugova, G. N., and Anisimova, I. V., 2004, Metamorphic evolution of granulites from the Kurul'ta Block, Aldan Shield: U-Pb single zircon study: *Doklady, Russian Academy of Sciences, Earth Sciences Section*, v. 398, p. 968–972.
- Sanders, I. S., Daly, J. S., and Davies, G. R., 1987, Late Proterozoic high-pressure granulite facies metamorphism in the north-east Ox inlier, north-west Ireland: *Journal of Metamorphic Geology*, v. 5, p. 69–85.
- Sandiford, M., Neall, F. B., and Powell, R., 1987, Metamorphic evolution of aluminous granulites from Labwor Hills, Uganda: *Contributions to Mineralogy and Petrology*, v. 95, 217–225.
- Santosh, M., 1987, Cordierite gneisses of southern Kerala, India: Petrology, fluid inclusions, and implications for crustal uplift history: *Contributions to Mineralogy and Petrology*, v. 96, p. 343–356.
- Santosh, M., Collins, A. S., Morimoto, T., and Yokoyama, K., 2005, Depositional constraints and age of metamorphism in southern India: U-Pb chemical (EMPA) and isotopic (SIMS) ages from the Trivandrum Block: *Geological Magazine*, v. 142, p. 255–268.
- Santosh, M., Harris, N. B. W., Jackson, D. H., and Matthey, D. P., 1990, Dehydration and incipient charnockite formation: A phase equilibria and fluid inclusion study from South India: *Journal of Geology*, v. 98, p. 915–926.
- Scharbert, H. G., and Carswell, D. A., 1983, Petrology of garnet-clinopyroxene rocks in a granulite facies environment, Bohemian Massif of Lower Austria: *Bulletin Mineralogique*, v. 106, p. 761–774.
- Schmädicke, E., and Will, T. M., 2003, Pressure-temperature evolution of blueschist facies rocks from Sifnos, Greece, and implications for the exhumation of high-pressure rocks in the Central Aegean: *Journal of Metamorphic Geology*, v. 21, p. 799–811.
- Schmitz, M. D., and Bowring, S. A., 2003, Ultrahigh-temperature metamorphism in the lower crust during Neoproterozoic Ventersdorp rifting and magmatism, Kaapvaal Craton, southern Africa: *Geological Society of America Bulletin*, v. 115, p. 533–548.
- Schulte, B., and Sindern, S., 2002, K-rich fluid metasomatism at high-pressure metamorphic conditions: Lawsonite decomposition in rodingitized ultramafite of the Maksyutovo Complex, southern Urals (Russia): *Journal of Metamorphic Geology*, v. 20, p. 529–541.
- Schumacher, R., Schenk, V., Raase, P., and Vitanage, P. W., 1990, Granulite facies metamorphism of metabasic and intermediate rocks in the Highland Series of Sri Lanka, in Ashworth, J. R., and Brown, M., eds., *High-temperature metamorphism and crustal anatexis*: London, UK, Unwin Hyman, p. 235–271.

- Scrimgeour, I., and Close, D., 1999, Regional high-pressure metamorphism during intracratonic deformation: The Petermann Orogeny, central Australia: *Journal of Metamorphic Geology*, v. 17, p. 557–572.
- Searle, M., Hacker, B. R., and Bilham, R., 2001, The Hindu Kush seismic zone as a paradigm for the creation of ultrahigh-pressure diamond- and coesite-bearing continental rocks: *The Journal of Geology*, v. 109, p. 143–153.
- Sen, S. K., and Bhattacharya, A., 1984, An orthopyroxene–garnet thermometer and its application to the Madras charnockites: *Contributions to Mineralogy and Petrology*, v. 88, p. 64–71.
- Sengupta, P., Raith, M. N., and Dasgupta, S., 2004, Comment on: “High-temperature dehydration melting and decompressive P-T path in a granulite complex from the Eastern Ghats India”: *Contributions to Mineralogy and Petrology*, v. 147, p. 123–125.
- Seth, B., Armstrong, R. A., Brandt, S., Villa, I. M., and Kramers, J. D., 2003, Mesoproterozoic U-Pb and Pb-Pb ages of granulites in NW Namibia: Reconstructing a complete orogenic cycle: *Precambrian Research*, v. 126, p. 147–168.
- Shabeer, K. P., Satish-Kumar, M., Armstrong, R., and Buick, I. S., 2005, Constraints on the timing of Pan-African granulite-facies metamorphism in the Kerala Khondalite Belt of Southern India: SHRIMP mineral ages and Nd isotope systematics: *The Journal of Geology*, v. 113, p. 95–106.
- Shafer, J. T., Neal, C. R., and Mahoney, J. J., 2005, Crustal xenoliths from Malaita, Solomon Islands: A window to the lower crust of the Ontong Java Plateau [abs.]: EOS (Transactions of the American Geophysical Union), v. 86, abstract T11C-0397.
- Shatsky, V. S., Sobolev, N. V., and Vavilov, M. A., 1995, Diamond-bearing metamorphic rocks of the Kokchetav Massif (Northern Kazakhstan), in Coleman, R. G., and Wang, X., eds., *Ultrahigh pressure metamorphism*: Cambridge, UK, Cambridge University Press, p. 427–455.
- Shervais, J. W., Dennis, A. J., McGee, J. J., and Secor, D., 2003, Deep in the heart of Dixie: Pre-Alleghanian eclogite and HP granulite metamorphism in the Carolina terrane, South Carolina, USA: *Journal of Metamorphic Geology*, v. 21, p. 65–80.
- Shibakusa, H., and Maekawa, H., 1997, Lawsonite-bearing eclogitic metabasites in the Cazadero area, northern California: *Mineralogy and Petrology*, v. 61, p. 163–180.
- Shirey, S. B., Richardson, S. H., and Harris, J. W., 2004, Integrated models of diamond formation and craton evolution: *Lithos*, v. 77, p. 923–944.
- Sklyarov, E. V., Theunissen, K., Melnikov, A. I., Klerkx, J., Gladkochub, D. P., and Mruma, A., 1998, Paleoproterozoic eclogites and garnet pyroxenites of the Ubende belt (Tanzania): *Schweizerische Mineralogische und Petrographische Mitteilungen*, v. 78, p. 257–271.
- Smith, D. C., 1984, Coesite in clinopyroxene in the Caledonides and its implications for geodynamics: *Nature*, v. 310, p. 641–644.
- Smith, D. C., 1988, A review of the peculiar mineralogy of the “Norwegian Coesite Eclogite Province,” with crystal-chemical, petrological, geochemical, and geodynamic notes and an extensive bibliography, in Smith, D. C., ed., *Eclogites and eclogite-facies rocks*: Amsterdam, The Netherlands, Elsevier, p. 1–206.
- Smithies, R. H., and Bagas, L., 1997, High pressure amphibolite-granulite facies metamorphism in the Paleoproterozoic Rudall Complex, central Western Australia: *Precambrian Research*, v. 83, p. 243–265.
- Sobolev, N. J., and Shatsky, V. S., 1990, Diamond inclusions in garnet from metamorphic rocks: A new environment for diamond formation: *Nature*, v. 343, p. 742–746.
- Song, S. G., Yang, J. S., Liou, J. G., and Shi, R. D., 2003, Metamorphic evolution of the coesite-bearing ultrahigh-pressure terrane in the North Qaidam, northern Tibet, NW China: *Journal of Metamorphic Geology*, v. 21, p. 613–644.
- Spaggiari, C. V., Gray, D. R., and Foster, D. A., 2002b, Blueschist metamorphism during accretion in the Lachlan orogen, south-eastern Australia: *Journal of Metamorphic Geology*, v. 20, p. 711–726.
- Spaggiari, C. V., Gray, D. R., Foster, D. A., and Fanning, C. M., 2002a, Occurrence and significance of blueschist in the southern Lachlan orogen: *Australian Journal of Earth Sciences*, v. 49, p. 255–269.
- Spandler, C., Rubatto, D., and Hermann, J., 2005, Late Cretaceous–Tertiary tectonics of the southwest Pacific: Insights from U-Pb sensitive, high-resolution ion microprobe (SHRIMP) dating of eclogite facies rocks from New Caledonia: *Tectonics*, v. 24, p. TC3003 [doi:10.1029/2004TC001709].
- Srikantappa, C., Raith, M., and Spiering, B., 1985, Progressive charnockitization of a leptynite–khondalite suite in southern Kerala, India—evidence for formation of charnockites through a decrease in fluid pressure?: *Journal of the Geological Society of India*, v. 26, p. 859–872.
- Stern, R. J., 2005, Evidence from ophiolites, blueschists, and ultrahigh-pressure metamorphic terranes that the modern episode of subduction tectonics began in Neoproterozoic time: *Geology*, v. 33, p. 557–560.
- Suzuki, S., Arima, M., Williams, I. S., Shiraishi, K., and Kagami, H., 2006, Thermal history of UHT metamorphism in the Napier Complex, east Antarctica: Insights from zircon monazite and garnet ages: *Journal of Geology*, v. 114, p. 65–84.
- Tagiri, M., Yano, T., Bakirov, A., Yakajima, T., and Uchiumi, S., 1995, Mineral parageneses and metamorphic P-T paths of ultrahigh-pressure eclogites from Kyrgyzstan Tien-Shan: *Island Arc*, v. 4, p. 280–292.

- Tamashiro, I., Santosh, M., Sajeev, K., Morimoto, T., and Tsunogae, T., 2004, Multistage orthopyroxene formation in ultrahigh-temperature granulites of Ganguvappatti, southern India: Implications for complex metamorphic evolution during Gondwana assembly: *Journal of Mineralogical and Petrological Sciences*, v. 99, p. 279–297.
- Thöni, M., 2005, Dating eclogites in the eastern Alps: Approaches, results, interpretations: *Mitteilungen der Österreichischen Mineralogischen Gesellschaft*, v. 150, p. 153.
- Tsujimori, T., Sisson, V. B., Liou, J. G., Harlow, G. E., and Sorensen, S. S., 2006a, Petrologic characterization of Guatemalan lawsonite eclogite: Eclogitization of subducted oceanic crust in a cold subduction zone, *in* Hacker, B. H., McClelland, W. C., and Liou, J. G., eds., *Ultrahigh-pressure metamorphism: Deep continental subduction*: Geological Society of America Special Paper 403, p. 147–168.
- Tsujimori, T., Sisson, V. B., Liou, J. G., Harlow, G. E., and Sorensen, S. S., 2006b, Very-low-temperature record of the subduction process: A review of worldwide lawsonite eclogites: *Lithos* [doi:10.1016/j.lithos.2006.03.054].
- Tsunogae, T., Miyano, T., van Reenen, D. D., and Smit, C. A., 2004, Ultrahigh-temperature metamorphism of the Southern Marginal Zone of the Archean Limpopo Belt, South Africa: *Journal of Mineralogical and Petrological Sciences*, v. 99, p. 213–224.
- Tumati, S., Thoni, M., Nimis, P., Martin, S., and Mair, V., 2003, Mantle-crust interactions during Variscan subduction in the Eastern Alps (Nonsberg-Ulten zone): *Geochronology and new petrological constraints: Earth and Planetary Science Letters*, v. 210, p. 509–526.
- Turner, F. J., 1968, *Metamorphic petrology*: New York, NY, McGraw-Hill, 403 p.
- Turner, F. J., 1970, Uniqueness vs. conformity to pattern in petrogenesis: *American Mineralogist*, v. 55, p. 339–348.
- Unsworth, M. J., Jones, A. G., Wei, W., Marquis, G., Gokarn, S. G., and Spratt, J. E., 2005, Crustal rheology of the Himalaya and Southern Tibet inferred from magnetotelluric data: *Nature*, v. 438, p. 78–81.
- Usui, T., Nakamuri, E., Kobayashi, K., Maruyama, S., and Helmstaedt, H., 2003, Fate of the subducted Farallon plate inferred from eclogite xenoliths in the Colorado Plateau: *Geology*, v. 31, p. 589–592.
- Valeriano, C. M., Machado, N., Simonetti, A., et al., 2004, U-Pb geochronology of the southern Brasília belt (SE-Brazil): Sedimentary provenance, Neoproterozoic orogeny, and assembly of West Gondwana: *Precambrian Research*, v. 130, p. 27–55.
- Valley, J. W., 2005, A cool early earth? *Scientific American*, v. 293, p. 58–65.
- Valley, J. W., Lackey, J. S., Cavoisie, A. J., Clechenko, C. C., Spicuzza, M. J., Basei, M. A. S., Bindeman, I. N., Ferreira, V. P., Sial, A. N., King, E. M., Peck, W. H., Sinha, A. K., and Wei, C. S., 2005, 4.4 billion years of crustal maturation: Oxygen isotope ratios of magmatic zircon: *Contributions to Mineralogy and Petrology*, v. 150, p. 561–580.
- van der Klaww, S. N. G. C., Reinecke, T., and Stöckhert, B., 1997, Exhumation of ultrahigh-pressure metamorphic oceanic crust from Lago di Cignana, Piedmontese zone, western Alps: A structural record in metabasites: *Lithos*, v. 41, p. 79–102.
- Volodichev, O. I., Slabunov, A. I., Bibikova, E. V., Konilov, A. N., and Kuzenko, T. I., 2004, Archean eclogites in the Belomorian mobile belt, Baltic Shield: *Petrology*, v. 12, p. 540–560.
- von Quadt, A., 1993, The Saxonian Granulite Massif: New aspects from geochronological studies: *Geologische Rundschau*, v. 82, p. 516–530.
- Vrána, S., 1989, Perpotassic granulites from southern Bohemia. A new rock-type derived from partial melting of crustal rocks under upper mantle conditions: *Contributions to Mineralogy and Petrology*, v. 103, p. 510–522.
- Wain, A., 1997, New evidence for coesite in eclogite and gneisses: Defining an ultrahigh-pressure province in the Western Gneiss Region of Norway: *Geology*, v. 25, p. 927–930.
- Wang, X., and Liou, J. G., 1991, Regional ultrahigh-pressure coesite-bearing eclogitic terrane in central China: Evidence from country rocks, gneiss, marble and metapelite: *Geology*, v. 19, p. 933–936.
- Wang, X., Liou, J. G., and Mao, H. K., 1989, Coesite-bearing eclogites from the Dabie mountains, central China: *Geology*, v. 17, p. 1085–1088.
- Wang, X., Zhang, R., and Liou, J. G., 1995, UHPM terrane in East Central China, *in* Coleman, R. G., and Wang, X., eds., *Ultrahigh pressure metamorphism*: Cambridge, UK, Cambridge University Press, p. 356–390.
- Watson, E. B., and Harrison, T. M., 2005, Zircon thermometer reveals minimum melting conditions on earliest Earth: *Science*, v. 308, p. 841–844.
- Watson, E. B., and Harrison, T. M., 2006, Response to comments on “Zircon thermometer reveals minimum melting conditions on earliest Earth”: *Science*, v. 311 [doi:10.1126/science.112.1080].
- Watson, E. B., Wark, D. A., and Thomas, J. B., 2006, Crystallization thermometers for zircon and rutile: *Contributions to Mineralogy and Petrology*, v. 151, no. 4, p. 413–433 [doi: 10.1007/s00410-006-0068-5].
- Wei, C., and Powell, R., 2003, Phase relations in high-pressure metapelites in the system KFMASH ( $K_2O$ – $FeO$ – $MgO$ – $Al_2O_3$ – $SiO_2$ – $H_2O$ ) with application to natural rocks: *Contributions to Mineralogy and Petrology*, v. 145, p. 301–315.
- Wei, C., and Powell, R., 2004, Calculated phase relations in high-pressure metapelites in the system NKFMAH ( $Na_2O$ – $K_2O$ – $FeO$ – $MgO$ – $Al_2O_3$ – $SiO_2$ – $H_2O$ ): *Journal of Petrology*, v. 45, p. 183–202.

- Wei, C., and Powell, R., 2006, Calculated phase relations in the system NCKFMASH ( $\text{Na}_2\text{O}-\text{CaO}-\text{K}_2\text{O}-\text{FeO}-\text{MgO}-\text{Al}_2\text{O}_3-\text{SiO}_2-\text{H}_2\text{O}$ ) for high-pressure metapelites: *Journal of Petrology*, v. 47, p. 385–408.
- Wei, C. J., Powell, R., and Zhang, L. F., 2003, Eclogites from the south Tianshan, NW China: Petrological characteristic and calculated mineral equilibria in the  $\text{Na}_2\text{O}-\text{CaO}-\text{FeO}-\text{MgO}-\text{Al}_2\text{O}_3-\text{SiO}_2-\text{H}_2\text{O}$  system: *Journal of Metamorphic Geology*, v. 21, p. 163–179.
- Wei, C. J., Zhang, C. G., Zhang, A. L., Wu, T. H., and Li, J. H., 2001, Metamorphic P-T conditions and geological significance of high-pressure granulite from the Jianping complex, western Liaoning Province: *Acta Petrologica Sinica*, v. 17, 269–282.
- White, R. W., Powell, R., and Clarke, G. L., 2002, The interpretation of reaction textures in Fe-rich metapelitic granulites of the Musgrave Block, central Australia: Constraints from mineral equilibria calculations in the system  $\text{K}_2\text{O}-\text{FeO}-\text{MgO}-\text{Al}_2\text{O}_3-\text{SiO}_2-\text{H}_2\text{O}-\text{TiO}_2-\text{Fe}_2\text{O}_3$ : *Journal of Metamorphic Geology*, v. 20, p. 41–55.
- Whitney, D. L., Miller, R. B., and Paterson, S. R., 1999, P-T-t evidence for mechanisms of vertical tectonic motion in a contractional orogen: north-western US and Canadian Cordillera: *Journal of Metamorphic Geology*, v. 17, p. 75–90.
- Williams, M. L., and Jercinovic, M. J., 2002, Microprobe monazite geochronology: Putting absolute time into microstructural analysis: *Journal of Structural Geology*, v. 24, p. 1013–1028.
- Willner, A. P., Rotzler, K., and Maresch, W. V., 1997, Pressure-temperature and fluid evolution of quartzofeldspathic metamorphic rocks with a relic high-pressure, granulite-facies history from the Central Erzgebirge (Saxony, Germany): *Journal of Petrology*, v. 38, p. 307–336.
- Wilson, J. T., and Burke, K., 1972, 2 types of mountain building: *Nature*, v. 239, p. 448–449.
- Worley, B., and Powell, R., 2000, High-precision relative thermobarometry: Theory and a worked example: *Journal of Metamorphic Geology*, v. 18, p. 91–101.
- Yamamoto, H., 1993, Contrasting metamorphic P-T-time paths of the Kohistan granulites and tectonics of the western Himalayas: *Journal of the Geological Society*, v. 150, p. 843–856.
- Yamamoto, H., and Yoshino, T., 1998, Superposition of replacements in the mafic granulites of the Jijal complex of the Kohistan arc, northern Pakistan: Dehydration and rehydration within deep arc crust: *Lithos*, v. 43, p. 219–234.
- Yoshino, T., Yamamoto, H., Okudaira, T., and Toriumi, M., 1998, Crustal thickening of the lower crust in the Kohistan arc (N Pakistan) deduced from Al zoning in clinopyroxene and plagioclase: *Journal of Metamorphic Geology*, v. 16, p. 729–748.
- Yuen, D. A., and Monnereau, M., 2005, Superadiabaticity of the lowermost part of the mantle thermally controlled by the post-perovskite phase transition [abs.]: EOS (Transactions of the American Geophysical Union), v. 86, abstract MR22A-03.
- Zack, T., Rivers, T., Brunn, R., and Kronz, A., 2004, Cold subduction of oceanic crust: Implications from a lawsonite eclogite from the Dominican Republic: *European Journal of Mineralogy*, v. 16, p. 909–916.
- Zhang, J., Zhang, Z., Xu, Z., Yang, J., and Cui, J., 2001, Petrology and geochronology of eclogite from the western segment of the Altyn Tagh, northwestern China: *Lithos*, v. 56, p. 187–206.
- Zhang, J. X., Mattinson, C. G., Meng, F. C., and Wan, Y. S., 2005, An early Paleozoic HP/HT granulite-garnet peridotite association in the south Altyn Tagh, NW China: P-T history and U-Pb geochronology: *Journal of Metamorphic Geology*, v. 23, p. 491–510.
- Zhang, L. F., Ai, Y. L., Li, Q., Li, X. P., Song, S. G., and Wei, C. J., 2005a, The formation and tectonic evolution of UHP metamorphic belt in southwestern Tianshan, Xinjiang: *Acta Petrologica Sinica*, v. 21, p. 1029–1038.
- Zhang, L., Ai, Y., Rubatto, D., Song, B., Williams, S., Ellis, D., Li, X., Song, S., and Liou, J. G., 2005b, Triassic collision of western Tianshan orogenic belt, China: Evidences from SHRIMP U-Pb dating of zircon from UHP eclogitic rocks: *Mitteilungen der Österreichischen Mineralogischen Gesellschaft*, v. 150, p. 173.
- Zhang, L., Ellis, D. J., Arculus, R. J., Jiang, W., and Wei, C., 2003, “Forbidden zone” subduction of sediments to 150 km depth—the reaction of dolomite to magnesite plus aragonite in the UHPM metapelites from western Tianshan, China: *Journal of Metamorphic Geology*, v. 21, p. 523–529.
- Zhang, L., Ellis, D. J., and Jiang, W., 2002, Ultrahigh-pressure metamorphism in western Tianshan: Part I. Evidence from inclusions of coesite pseudomorphs in garnet and from quartz exsolution lamellae in omphacite in eclogites: *American Mineralogist*, v. 87, p. 853–860.
- Zhang, L. F., Song, S. G., Liou, J. G., Ai, Y. L., and Li, X. P., 2005c, Relict coesite exsolution in omphacite from Western Tianshan eclogites, China: *American Mineralogist*, v. 90, p. 181–186.
- Zhang, R. Y., Hirajima, T., Banno, S., Cong, B., and Liou, J. G., 1995, Petrology of ultrahigh-pressure rocks from the southern Su-Lu region, eastern China: *Journal of Metamorphic Geology*, v. 13, p. 659–675.
- Zhao, G. C., Cawood, P. A., Wilde, S. A., and Lu, L. Z., 2001, High-pressure granulites (retrograded eclogites) from the Hengshan Complex, North China Craton: Petrology and tectonic implications: *Journal of Petrology*, v. 42, p. 1141–1170.
- Zwart, H. J., 1967, The duality of orogenic belts: *Geologie en Mijnbouw*, v. 46, p. 283–309.