

Introduction to special section: Active Fault-Related Folding: Structural Evolution, Geomorphologic Expression, Paleoseismology, and Seismic Hazards

James F. Dolan¹ and Jean-Philippe Avouac²

Received 24 January 2007; revised 26 January 2007; accepted 31 January 2007; published 28 March 2007.

Citation: Dolan, J. F., and J.-P. Avouac (2007), Introduction to special section: Active Fault-Related Folding: Structural Evolution, Geomorphologic Expression, Paleoseismology, and Seismic Hazards, *J. Geophys. Res.*, 112, B03S01, doi:10.1029/2007JB004952.

[1] Folds are one of the obvious and common manifestations of deformation of the continental crust, and attempts to understand the origins of these structures extend back to the very beginnings of geology as a discipline within the Earth Sciences. In particular, the relationship between faults and folds, and the manner in which fault-related folds grow and evolve through time, has been the subject of intense interest within the structural geology and tectonics communities for more than a century. Indeed, it did not take long after geologists understood the basic principles of stratigraphy and geological mapping [Lyell, 1830] to realize that subsurface structures could be delineated with some accuracy from surface observations, through the use of simple geometric rules [e.g., Rogers, 1856]. The fact that folding is associated with thrust faulting and mountain building was also recognized in the early days of structural geology, together with the necessity for large tangential displacements at the Earth's surface across fold-thrust belts [Rogers, 1856; Moesch, 1867; Heim, 1878; Muller, 1878; Willis, 1891; Chamberlin, 1910; Argand, 1924; Rich, 1934]. Moreover, geologists quickly realized how to relate quantitatively folding and horizontal shortening by assuming conservation of mass [Chamberlin, 1910], and soon began investigating the mechanics of folding and the relationships between fold growth and detachment faulting and thrusting through the use of analogue experiments [e.g., Willis, 1891] (Figure 1). The necessity of low friction along the basal detachments of fold-thrust systems was also noticed early on [Hubbert and Rubey, 1959].

[2] Over the past several decades, the ever increasing availability of observations from natural exposures and geophysical exploration (much of the latter conducted by the petroleum industry) has led to tremendous progress in our understanding of the kinematics and mechanics of fault-related folding, and the relationship between fault slip and fold growth, at scales ranging from individual structures to entire orogens [e.g., Price, 1981; Boyer and Elliott, 1982; Suppe, 1983; Davis et al., 1983; Dahlstrom, 1990; Suppe and Medwedeff, 1990; Erslev, 1991; Allmendinger, 1998;

Allmendinger and Shaw, 2000]. Collectively, these advances have facilitated modeling and quantitative analysis of the origin and evolution of fault-related folds and have fostered an understanding of fold-thrust belts as integrated mechanical systems. That understanding how fold-and-thrust systems evolve is not only of academic and economic significance has been amply demonstrated by the recent occurrence of a number of highly destructive thrust earthquakes (e.g., 1980 M_s 7.3 El Asnam, M_w 6.5 Coalinga, 1994 M_w 6.7 Northridge, 1999 M_w 7.6 Chi-Chi, and 2005 M_w 7.6 Kashmir) [King and Vita-Finzi, 1981; Yielding et al., 1981; Philip and Meghraoui, 1983; Stein and King, 1984; Stein and Ekstrom, 1992; Scientists of the USGS and SCEC, 1994; Yu et al., 2001; Avouac et al., 2006]. These events have led to the growing recognition of the hazards posed by such structures to numerous urban centers around the world and highlight the need to understand the relationship between seismic slip on thrust faults and the resulting fold growth. These issues have also been the focus of much recent research by structural geologists, particularly those interested in understanding the mechanisms and rates of fold growth at timescales shorter than those provided by most exhumed examples of fold-thrust belts.

[3] Motivated by all of these issues, there has been a dramatic increase in recent research focused on active fold growth and associated fault slip. Many of these recent studies take advantage of technical advances in a number of disciplines, including (1) development of new methods to unravel the history of folding from analysis of growth strata (i.e., strata deposited above and adjacent to growing folds) [e.g., Suppe et al., 1992, 1997; Shaw and Suppe, 1994, 1996; Hardy et al., 1995; Storti and Poblet, 1997; Novoa et al., 2000; Shaw et al., 2005], (2) recognition that the kinematics of active folding and thrusting is recorded by alluvial, fluvial, and marine geomorphologic markers [e.g., Rockwell et al., 1984, 1988; Stein and King, 1984; Atwater et al., 1990; DeCelles et al., 1991; Dolan and Sieh, 1992; Avouac et al., 1993; Bullard and Lettis, 1993; Molnar et al., 1994; Burbank et al., 1996; Jackson et al., 1996; Benedetti et al., 2000; Van der Woerd et al., 2001; Ishiyama et al., 2004; Bennett et al., 2005], (3) advances in our general understanding of how tectonic uplift and climatic factors influence geomorphologic processes such as alluvial deposition and river entrenchment [e.g., Weldon, 1986; Bull, 1991; Schumm et al., 2000; Burbank and Anderson, 2001; Poisson and Avouac, 2004], and development of new

¹Department of Earth Sciences, University of Southern California, Los Angeles, California, USA.

²Tectonics Observatory, California Institute of Technology, Pasadena, California, USA.




Figure 1. Sketch of the apparatus constructed by Willis [1891] to investigate folding mechanisms. The layers, representing geological strata, are made out of wax, and the model is overlain with lead sheets to qualitatively take into account the effect of gravity.

methods to extract quantitative tectonic signals from the geomorphologic record of folding and thrusting [Lavé and Avouac, 2000; Kirby and Whipple, 2001; Thompson et al., 2002; Lague and Davy, 2003], (4) the use of both well-established and emerging geochronologic techniques (e.g., radiocarbon, cosmogenic radionuclide, optically stimulated luminescence dating, and magnetostratigraphy) to date growth strata and geomorphologic surfaces [Aitken, 1985; 1998; Noller et al., 2000; Gosse and Phillips, 2001], (5) analysis of high-resolution seismic reflection data collected across zones of active folding above thrust faults [e.g., Pratt et al., 1997, 2002; Shaw et al., 2002; Ishiyama et al., 2004], (6) development of new techniques to determine the paleoearthquake history of thrust faults through the recognition of ancient fold scarps [e.g., Dolan et al., 2003; Ishiyama et al., 2004], and (7) use of geodetic data, satellite imagery, seismic waveform modeling and precisely relocated seismicity to delineate coseismic and postseismic fault slip and associated fold growth, as well as the geometry of the causative fault [e.g., Shaw and Shearer, 1999; Ji et al., 2001; Johnson et al., 2001; Hsu et al., 2002; Dominguez et al., 2003; Avouac et al., 2006].

[4] This special section comprises 15 articles that address these issues using active examples of fault-related folding. These studies span a wide spectrum of study, from analyses of fold growth during individual earthquakes to kinematic models of the long-term development of entire fault fold systems to mechanical studies of deformation mechanisms. The papers here are all highly multidisciplinary in nature. In particular, their focus on the behavior and evolution of active structures allows the use of geodetic, seismologic, and geomorphologic data, which are not available for ancient examples. Such data facilitate the analysis of the

detailed rates, geometries, and styles of folding, which are typically much more difficult to discern in exhumed fold-thrust belts. Several of the papers focus on blind thrust faults, which are a common component of many fold-thrust systems but have generally proved to be difficult to investigate. Because these faults do not extend to the surface, analysis of their seismic behavior, and the degree to which folding above them is accommodated within discrete structural domains, as opposed to distributed deformation, has proved both problematic and controversial.

[5] Some of the specific questions that are addressed by the recent research described in this special section include the following:

[6] 1. How does the near-surface expression of folding relate to the deeper fold structure and the history of thrusting and folding?

[7] 2. Can we use data from active structures to test the validity of existing models, or are new models of fold growth required?

[8] 3. How can mechanical and analog models improve our understanding of the detailed mechanisms of fold growth?

[9] 4. How do fault-related folds grow in relation to the earthquake cycles on underlying faults? Do fault-related folds grow primarily during earthquakes, or is fold growth typically a more gradual process?

[10] 5. How do fold growth and fault slip rates vary over different timescales, ranging from seismic cycles to millions of years? How fast do folds (and by extension their causative faults) grow laterally?

[11] Many of the papers in this special section take advantage of the recent advances described above to determine the kinematics of folding and thrusting through

multidisciplinary approaches that combine subsurface imaging from geophysical techniques or well logs, geomorphologic investigations, dating of geomorphologic markers and growth strata, and structural modeling [Amos *et al.*, 2007; Y.-G. Chen *et al.*, 2007; J. Chen *et al.*, 2007; Daëron *et al.*, 2007; Guzowski *et al.*, 2007; Hubert-Ferrari *et al.*, 2007; Okamura *et al.*, 2007; Simoes *et al.*, 2007a, 2007b; Vergés *et al.*, 2007]. A number of these studies document the important result that near-surface folding associated with changes in thrust fault dip at depth (fault bend folds in the terminology of Suppe [1983]) is typically confined to narrow, structurally discrete zones, rather than distributed over the width of the fold, consistent with the idea that folding occurs within relatively narrow hinge zones [Y.-G. Chen *et al.*, 2007; Guzowski *et al.*, 2007; Hubert-Ferrari *et al.*, 2007; Ishiyama *et al.*, 2007; Leon *et al.*, 2007]. In particular, the geometries of growth and pregrowth strata in a number of these folds (e.g., Hubert-Ferrari *et al.*, Leon *et al.*) are consistent with kinematic models of kink band migration, yet show consistent steepening of limb dips in growth strata. Perhaps most spectacularly, Y.-G. Chen *et al.*, in their study of the thrust faults and folds that were active during the 1999 M_w 7.6 Chi-Chi, Taiwan, earthquake, document coseismic development of a discrete fold scarp formed due to deformation within a kink band associated with a deeply buried change in fault dip. This observation demonstrates that the history of coseismic deformation might be deciphered through geomorphologic observations of surface scarps and deformed fluvial and alluvial landforms, as well as through excavations of young growth strata that record the detailed history of recent folding, as described in papers by Leon *et al.*, Ishiyama *et al.*, and Streig *et al.* [2007]. These studies document stratigraphically and geomorphologically discrete uplift, as would be expected in temporally discrete, and likely coseismic, folding events.

[12] Interpreting the geometry and deformation history of folded strata, whether at the hundred-meter scale of paleoseismic investigations, or the multiple-kilometer scale of entire folds, is not an easy task, and generally requires some sort of kinematic modeling. For example, a number of studies included in this special section describe the successful application of the trishear kinematic model [Erslev, 1991; Allmendinger, 1998] to the analysis of coseismic fold scarps (Y.-G. Chen *et al.*, Streig *et al.*). When it comes to using such kinematic models to understand the long-term development of large-scale fault tip folds (i.e., folds formed at the tip of detachments or thrust faults that do not reach the surface) [Dahlstrom, 1990; Poblet and McClay, 1996; Mitra, 2003], the applicability of both the trishear fold model and more conventional flexural slip and kink band migration models, which has proved to be so effective in modeling fault bend folds, is limited. This is because fault tip folds generally grow as a result of heterogeneously distributed pure shear, rather than by bed-parallel simple shear, in a way that cannot be easily reproduced from the trishear model. In this regard, parameterization of the analogue model of Bernard *et al.* [2007] to allow adjustment in response to surface structural observations, geomorphologic measurements, and subsurface data (Daëron *et al.*, Simoes *et al.*(b)) represents a significant step forward. The manner in which these folds grow in detail, however,

and most especially the degree to which fold growth is coseismic or interseismic, remains poorly resolved. Moreover, these techniques work best in relatively young, structurally simple folds. Analysis of the kinematic and seismic behavior of more complicated examples, however, is extremely challenging. For example, Guzowski *et al.*, as part of their analysis of the blind thrusts faults and associated folds of the western San Joaquin fold-thrust belt, show that despite the simplicity of the fold structure near the surface, the deep structure observed on seismic reflection profiles is a relatively complex wedge system. They additionally show that multiplying the uplift signature of the 1983 M_w 6.5 Coalinga earthquake by thousands of such events does not yield fold geometries that are consistent with the actual geometry of the Coalinga anticlinorium. Similarly, complex wedge geometries are also observed below the Cerro Salinas anticline, which produced $M7$ earthquakes along the Andean front in Argentina (Vergés *et al.*), and below the Quilitag anticline near Kuqa (western China) (Hubert-Ferrari *et al.*).

[13] Despite the past success of kinematic fold models in interpreting both geomorphologic and subsurface data, this area of research would benefit from the further development of mechanical fold models (i.e., those that take into account how the kinematics of deformation relates to stress and rock rheology). This is an important issue because a model that might seem admissible from a kinematic point of view might be unrealistic from a mechanical point of view. In practice, this is an extremely difficult issue to address from both a theoretical and a numerical standpoint [e.g., Mandl, 1988; Maillot and Leroy, 2003]. New methods, however, such as the discrete element technique described here by Benesh *et al.* [2007], offer significant promise. In their study, Benesh *et al.* show that the record of near-surface folding recorded by growth strata depends strongly on rock rheology, in a way that kinematic modeling might not capture correctly. This is an area where much can also be learned from the detailed analysis of analogue experiments, such as those described in the special section by Bernard *et al.*, and earlier studies by Davis *et al.* [1983], Malavieille [1984], Adam *et al.* [2005], and Maillot and Koyi [2006]. The origin of the low basal friction suggested by the low taper angle of many thrust wedges is another issue that might benefit from insights provided by the analysis of the mechanics of active fold-thrust belts. Evidence of extremely high pore pressure within basal shear zones of subaerial thrust systems is, in fact, scarce, suggesting that the theory of Hubbert and Rubey [1959] may not be generally applicable in such settings. Given that the basal detachment mainly slips during recurring large earthquakes, one possibility would be that the apparent low friction would reflect a low dynamic friction, rather than a low effective friction [e.g., Avouac, 2003]. Low dynamic friction is a problem that has received significant recent attention, but that is not covered in this special section. It might be a specific property of thrust faults of geometric origin [e.g., Brune *et al.*, 1993; Oglesby *et al.*, 2000], or a more general property due to mechanisms that may not necessarily involve fluids [e.g., Kanamori and Brodsky, 2004].

[14] Collectively, the studies gathered in this special section highlight the benefits of the multidisciplinary study of active, fault-related folding, as well as the tremendous

advances made during the past several years in our understanding of the origins, evolution, mechanics, and seismic potential of fault-related folds. These studies also point to limitations of our current understanding of the mechanics of fault-related folding, providing direction for future research.

References

- Adam, J., J. L. Urai, B. Wieneke, O. Oncken, K. Pfeiffer, N. Kukowski, J. Lohrmann, S. Hoth, W. Van der Zee, and J. Schmatz (2005), Shear localisation and strain distribution during tectonic faulting: New insights from granular-flow experiments and high-resolution optical image correlation techniques, *J. Struct. Geol.*, **27**, 283–301.
- Aitken, M. J. (1985), *Thermo-luminescence Dating*, 359 pp., Elsevier, New York.
- Aitken, M. J. (1998), *An Introduction to Optical Dating: The Dating of Quaternary Sediments by the Use of Photon-Stimulated Luminescence*, 267 pp., Oxford Univ. Press, New York.
- Allmendinger, R. W. (1998), Inverse and forward numerical modeling of trishear fault propagation folds, *Tectonics*, **17**, 640–656.
- Allmendinger, R. W., and J. H. Shaw (2000), Estimation of fault propagation distance from fold shape: Implications for earthquake hazard assessment, *Geology*, **28**, 1099–1102.
- Amos, C. B., D. W. Burbank, D. C. Nobes, and S. A. L. Read (2007), Geomorphic constraints on listric thrust faulting: Implications for active deformation in the Mackenzie Basin, South Island, New Zealand, *J. Geophys. Res.*, **112**, B03S11, doi:10.1029/2006JB004291.
- Argand, E. (1924), La tectonique de l'Asie, paper presented at 13th International Geological Congress, Brussels.
- Atwater, B. F., D. A. Trumm, J. C. Tinsley III, R. S. Stein, A. B. Tucker, D. J. Donahue, A. J. T. Tull, and L. A. Payen (1990), Alluvial plains and earthquake recurrence at the Coalinga Anticline, *U.S. Geol. Surv. Prof. Pap.*, **1487**, 273–198.
- Avouac, J. P. (2003), Mountain building, erosion and the seismic cycle in the Nepal Himalaya, *Adv. Geophys.*, **46**, 1–79, doi:10.1016/S0065-2687(03)46001-9.
- Avouac, J. P., P. Tapponnier, M. Bai, H. You, and G. Wang (1993), active thrusting and folding along the northern Tien-Shan and Late Cenozoic rotation of the Tarim relative to Dzungaria and Kazakhstan, *J. Geophys. Res.*, **98**, 6755–6804.
- Avouac, J. P., F. Ayoub, S. Leprince, O. Konca, and D. V. Helmberger (2006), The 2005, Mw 7.6 Kashmir earthquake: Sub-pixel correlation of ASTER images and seismic waveforms analysis, *Earth Planet. Sci. Lett.*, **249**, 514–528.
- Benedetti, L., P. Tapponnier, G. C. P. King, B. Meyer, and I. Manighetti (2000), Growth folding and active thrusting in the Montello region, Veneto, northern Italy, *J. Geophys. Res.*, **105**, 8227–8228.
- Benesh, N. P., A. Plesch, J. H. Shaw, and E. K. Frost (2007), Investigation of growth fault-bend folding using discrete element modeling: Implications for signatures of active folding above blind-thrust faults, *J. Geophys. Res.*, doi:10.1029/2006JB004466, in press.
- Bennett, E. R., J. H. Youngson, J. A. Jackson, R. J. Norris, G. M. Raisbeck, F. You, and E. Fielding (2005), Growth of South Rough Ridge, Central Otago, New Zealand: Using in situ cosmogenic isotopes and geomorphology to study an active, blind reverse fault, *J. Geophys. Res.*, **110**, B02404, doi:10.1029/2004JB003184.
- Bernard, S., J.-P. Avouac, S. Dominguez, and M. Simoes (2007), Kinematics of fault-related folding derived from a sandbox experiment, *J. Geophys. Res.*, **112**, B03S12, doi:10.1029/2005JB004149.
- Boyer, S. E., and D. Elliott (1982), Thrust systems, *AAPG Bull.*, **66**, 1196–1230.
- Brune, J., S. Brown, and P. Johnson (1993), Rupture mechanism and interface separation in foam rubber models of earthquakes: A possible solution to the heat flow paradox and the paradox of large overthrusts, *Tectonophysics*, **218**, 59–67.
- Bull, W. B. (1991), *Geomorphic Responses to Climatic Change*, 326 pp., Oxford Univ. Press, New York.
- Bullard, T. F., and W. R. Lettis (1993), Quaternary fold deformation associated with blind thrust faulting, Los Angeles basin, California, *J. Geophys. Res.*, **98**, 8349–8369.
- Burbank, D. W., and R. S. Anderson (2001), *Tectonic Geomorphology*, 274 pp., Blackwell Sci., Malden, Mass.
- Burbank, D., A. Meigs, and N. Brozovic (1996), Interactions of growing folds and coeval depositional systems, *Basin Res.*, **8**, 199–223.
- Chamberlin, R. T. (1910), The Appalachian folds of central Pennsylvania, *J. Geol.*, **18**, 228–251.
- Chen, J., R. Heermance, D. W. Burbank, K. M. Scharer, J. Miao, and C. Wang (2007), Quantification of growth and lateral propagation of the Kashi anticline, southwest Chinese Tian Shan, *J. Geophys. Res.*, doi:10.1029/2006JB004345, in press.
- Chen, Y.-G., K.-Y. Lai, Y.-H. Lee, J. Suppe, W.-S. Chen, Y.-N. N. Lin, Y. Wang, J.-H. Hung, and Y.-T. Kuo (2007), Coseismic fold scarps and their kinematic behavior in the 1999 Chi-Chi earthquake Taiwan, *J. Geophys. Res.*, **112**, B03S02, doi:10.1029/2006JB004388.
- Daëron, M., J.-P. Avouac, and J. Charreau (2007), Modeling the shortening history of a fault tip fold using structural and geomorphic records of deformation, *J. Geophys. Res.*, **112**, B03S13, doi:10.1029/2006JB004460.
- Dahlstrom, C. D. A. (1990), Geometric constraints derived from the law of conservation of volume and applied to evolutionary models for detachment folding, *AAPG Bull.*, **74**, 336–344.
- Davis, D., J. Suppe, and F. A. Dahlen (1983), Mechanics of fold-and-thrust belts and accretionary wedges, *J. Geophys. Res.*, **88**, 1153–1172.
- DeCelles, P. G., M. B. Gray, K. D. Ridgway, R. B. Cole, P. Srivastava, N. Pequera, and D. A. Pivnik (1991), Kinematic history of foreland uplift from Paleocene synorogenic conglomerate, Beartooth Range, Wyoming and Montana, *Geol. Soc. Am. Bull.*, **103**, 1458–1475.
- Dolan, J. F., and K. Sieh (1992), Tectonic geomorphology of the northern Los Angeles basin: Seismic hazards and kinematics of young fault movement, in *Engineering Geology Field Trips: Orange County, Santa Monica Mountains, and Malibu, Guidebook and Volume*, edited by P. L. Ehlig and E. A. Steiner, pp. B-20 to B-26, Assoc. of Eng. Geol., South. Calif. Sect., Long Beach, 2–9 Oct.
- Dolan, J. F., S. A. Christofferson, and J. H. Shaw (2003), Recognition of paleoearthquakes on the Puente Hills blind thrust fault, California, *Science*, **300**, 115–118.
- Dominguez, S., J. Avouac, and R. Michel (2003), Horizontal coseismic deformation of the 1999 Chi-Chi earthquake measured from SPOT satellite images: Implications for the seismic cycle along the western foothills of central Taiwan, *J. Geophys. Res.*, **108**(B2), 2083, doi:10.1029/2001JB000951.
- Erslev, E. A. (1991), Trishear fault-propagation folding, *Geology*, **19**, 617–620.
- Gosse, J. C., and F. M. Phillips (2001), Terrestrial in situ cosmogenic nuclides: Theory and application, *Quat. Sci. Rev.*, **20**, 1475–1560.
- Guzofski, C. A., J. H. Shaw, G. Lin, and P. M. Shearer (2007), Seismically active wedge structure beneath the Coalinga anticline, San Joaquin basin, California, *J. Geophys. Res.*, doi:10.1029/2006JB004465, in press.
- Hardy, S., J. Poblet, K. McClay, and D. Waltham (1995), Mathematical modelling of growth strata associated with fault-related fold structures, in *Modern Developments in Structural Interpretation, Validation and Modelling*, edited by P. G. Buchanan and D. A. Nieuwland, *Geol. Soc. Spec. Publ.*, **99**, 265–282.
- Heim, A. (1878), *Der Mechanismus des Gebirgsbildung*, 233:207 pp.
- Hsu, Y., N. Bechor, P. Segall, S. Yu, L. Kuo, and K. Ma (2002), Rapid afterslip following the 1999 Chi-Chi, Taiwan earthquake, *Geophys. Res. Lett.*, **29**(16), 1754, doi:10.1029/2002GL014967.
- Hubbert, M. K., and W. W. Rubey (1959), Role of fluid pressure in mechanics of overthrust faulting. I. Mechanics of fluid-filled porous solids and its application to overthrust faulting, *Geol. Soc. Am. Bull.*, **79**, 115–166.
- Hubert-Ferrari, A., J. Suppe, R. Gonzalez-Mieres, and X. Wang (2007), Mechanisms of active folding of the landscape (southern Tian Shan, China), *J. Geophys. Res.*, doi:10.1029/2006JB004362, in press.
- Ishiyama, T., K. Mueller, M. Togo, A. Okada, and K. Takemura (2004), Geomorphology, kinematic history, and earthquake behavior of the active Kuwana wedge thrust anticline, central Japan, *J. Geophys. Res.*, **109**, B12408, doi:10.1029/2003JB002547.
- Ishiyama, T., K. Mueller, H. Sato, and M. Togo (2007), Coseismic fault-related fold model, growth structure, and the historic multisegment blind thrust earthquake on the basement-involved Yoro thrust, central Japan, *J. Geophys. Res.*, doi:10.1029/2006JB004377, in press.
- Jackson, J., R. Norris, and J. Youngson (1996), The structural evolution of active fault and fold systems in central Otago, New Zealand: Evidence revealed by drainage patterns, *J. Struct. Geol.*, **18**, 217–234.
- Ji, C., D. V. Helmberger, T.-R. A. Song, K.-F. Ma, and D. J. Wald (2001), Slip distribution and tectonic implications of the 1999 Chi-Chi, Taiwan earthquake, *Geophys. Res. Lett.*, **28**, 4379–4382.
- Johnson, K. J., Y.-J. Hsu, P. Segall, and S.-B. Yu (2001), Fault geometry and slip distribution of the 1999 Chi-Chi, Taiwan earthquake imaged from inversion of GPS data, *Geophys. Res. Lett.*, **28**, 2285–2288.
- Kanamori, H., and E. E. Brodsky (2004), The physics of earthquakes, *Rep. Progress Phys.*, **67**, 1429–1496.
- King, G. C. P., and C. Vita-Finzi (1981), Active folding in the Algerian earthquake of 10 October 1980, *Nature*, **292**, 22–26.
- Kirby, E., and K. Whipple (2001), Quantifying differential rock-uplift rates via stream profile analysis, *Geology*, **29**, 415–418.
- Lague, D., and P. Davy (2003), Constraints on the long-term colluvial erosion law by analyzing slope-area relationships at various tectonic

- uplift rates in the Siwaliks Hills (Nepal), *J. Geophys. Res.*, 108(B2), 2129, doi:10.1029/2002JB001893.
- Lavé, J., and J. P. Avouac (2000), Active folding of fluvial terraces across the Siwaliks Hills, Himalayas of central Nepal, *J. Geophys. Res.*, 105, 5735–5770.
- Leon, L. A., S. A. Christofferson, J. F. Dolan, J. H. Shaw, and T. L. Pratt (2007), Earthquake-by-earthquake fold growth above the Puente Hills blind thrust fault, Los Angeles, California: Implications for fold kinematics and seismic hazard, *J. Geophys. Res.*, doi:10.1029/2006JB004461, in press.
- Lyell, C. (1830), *Principles of Geology: The Surface of the Earth Is Shaped by Erosion, Sedimentation, Earthquakes and Volcanic Processes*, Murray, J., London.
- Maillot, B., and H. Koyi (2006), Thrust dip and thrust refraction in fault-bend folds: Analogue models and theoretical predictions, *J. Struct. Geol.*, 28, 36–49.
- Maillot, B., and Y. M. Leroy (2003), Optimal dip based on dissipation of back thrusts and hinges in fold-and-thrust belts, *J. Geophys. Res.*, 108(B6), 2320, doi:10.1029/2002JB002199.
- Malavieille, J. (1984), Modelisation experimentale des chevauchements imbriques: Application aux chaines de montagnes, *Bull. Soc. Geol. Fr.*, 26(1), 129–138.
- Mandl, G. (1988), *Mechanics of Tectonic Faulting*, *Dev. Struct. Geol.*, vol. X, edited by H. J. Zwart, Elsevier, New York.
- Mitra, S. (2003), A unified kinematic model for the evolution of detachment folds, *J. Struct. Geol.*, 25, 1659–1673.
- Moesch, C. (1867), Der Aargaw Jura und die nordlichen Gebiet des Cantons Zurich, *Beitr. Geol. Karte Schweiz*, IV.
- Molnar, P., et al. (1994), Quaternary climate-change and the formation of river terraces across growing anticlines on the north flank of the Tien-Shan, China, *J. Geol.*, 102, 583–602.
- Muller, A. (1878), Ueber di anomalen Lagerungsverhältnisse im westlichen Balser Jura, *Verhandl. Naturforsch. Gesellsch. Basel*, 428–462.
- Noller, J. S., J. M. Sowers, and W. R. Lettis (2000), *Quaternary Geochronology: Methods and Applications*, 582 pp., AGU, Washington, D. C.
- Novoa, E., J. Suppe, and J. H. Shaw (2000), Inclined-shear restoration of growth folds, *AAPG Bull.*, 84, 787–804.
- Oglesby, D., R. Archuleta, and S. Nielsen (2000), Dynamics of dip-slip faulting: Explorations in two dimensions, *J. Geophys. Res.*, 105, 13,643–13,653.
- Okamura, Y., T. Ishiyama, and Y. Yanagisawa (2007), Fault-related folds above the source fault of the 2004 mid-Niigata Prefecture earthquake, in a fold-and-thrust belt caused by basin inversion along the eastern margin of the Japan Sea, *J. Geophys. Res.*, doi:10.1029/2006JB004320, in press.
- Philip, H., and M. Meghraoui (1983), Structural analysis and interpretation of the surface deformations of the El-Asnam earthquake of October 10, 1980, *Tectonics*, 2, 17–49.
- Poblet, J., and K. McClay (1996), Geometry and kinematics of single-layer detachment folds, *AAPG Bull.*, 80, 1085–1109.
- Poisson, B., and J. P. Avouac (2004), Holocene hydrological changes inferred from alluvial stream entrenchment in north Tian Shan (north-western China), *J. Geol.*, 112, 231–249.
- Pratt, T. L., S. Y. Johnson, C. J. Potter, W. J. Stephenson, and C. Finn (1997), Seismic reflection images beneath Puget Sound, western Washington state: The Puget Sound Lowland thrust sheet hypothesis, *J. Geophys. Res.*, 102, 27,469–27,489.
- Pratt, T. L., J. H. Shaw, J. F. Dolan, S. A. Christofferson, R. A. Williams, J. K. Odum, and A. Plesch (2002), Shallow seismic imaging of folds above the Puente Hills blind-thrust fault, Los Angeles, California, *Geophys. Res. Lett.*, 29(9), 1304, doi:10.1029/2001GL014313.
- Price, R. A. (1981), The Cordilleran fold and thrust belt in the southern Canadian Rockies, in *Thrust and Nappe Tectonics*, edited by K. R. McClay and N. J. Price, *Geol. Soc. London Spec. Publ.*, 9, 427–448.
- Rich, J. L. (1934), Mechanics of low-angle overthrust faulting as illustrated by the Cumberland thrust block, Virginia, Kentucky and Tennessee, *Am. Assoc. Pet. Geol. Bull.*, 18, 1584–1596.
- Rockwell, T. K., E. A. Keller, M. N. Clark, and D. L. Johnson (1984), Chronology and rates of faulting of Ventura River terraces, California, *Geol. Soc. Am. Bull.*, 95, 1466–1474.
- Rockwell, T. K., E. A. Keller, and G. R. Dembroff (1988), Quaternary rate of folding of the Ventura Avenue anticline, western Transverse Ranges, southern California, *Geol. Soc. Am. Bull.*, 100, 850–858.
- Rogers, H. D. (1856), On the laws of structure of the more disturbed zones of the Earth's crust, *Trans. R. Soc. Edinburgh*, XXI, 442.
- Schumm, S. A., J. F. Dumont, and J. M. Holbrook (2000), *Active Tectonics and Alluvial Rivers*, 276 pp., Cambridge Univ. Press, New York.
- Scientists of the USGS and SCEC (1994), The magnitude 6.7 Northridge, California, earthquake of 17 January 1994, *Science*, 266, 389–397.
- Shaw, J. H., and J. Suppe (1994), Active faulting and growth folding in the eastern Santa Barbara Channel, California, *Geol. Soc. Am. Bull.*, 106, 607–626.
- Shaw, J. H., and P. M. Shearer (1999), An elusive blind-thrust fault beneath metropolitan Los Angeles, *Science*, 283, 1516–1518.
- Shaw, J. H., and J. Suppe (1996), Earthquake hazards of active blind-thrust faults under the central Los Angeles Basin, California, *J. Geophys. Res.*, 101, 8623–8642.
- Shaw, J. H., A. Plesch, J. F. Dolan, T. Pratt, and P. Fiore (2002), Puente Hills blind-thrust system, Los Angeles Basin, California, *Bull. Seismol. Soc. Am.*, 92, 2946–2960.
- Shaw, J. H., C. Connors, and J. Suppe (Eds.) (2005), *Seismic Interpretation of Contractual Fault-Related Folds*, *AAPG Stud. Geol.*, vol. 53, Am. Assoc. of Pet. Geol., Tulsa, Okla.
- Simoës, M., J. P. Avouac, and Y.-G. Chen (2007a), Slip rates on the Chelungpu and Chushiang thrust faults inferred from a deformed strath terrace along the Dungpuna river, west central Taiwan, *J. Geophys. Res.*, 112, B03S10, doi:10.1029/2005JB004200.
- Simoës, M., J.-P. Avouac, Y.-G. Chen, A. K. Singhvi, C.-Y. Wang, M. Jaiswal, Y.-C. Chan, and S. Bernard (2007b), Kinematic analysis of the Pakuashan fault tip fold, west central Taiwan: Shortening rate and age of folding inception, *J. Geophys. Res.*, 112, B03S14, doi:10.1029/2005JB004198.
- Stein, R. S., and G. Ekstrom (1992), Seismicity and geometry of a 110-km-long blind thrust fault. 2. Synthesis of the 1982–1985 California earthquake sequence, *J. Geophys. Res.*, 97, 4865–4883.
- Stein, R. S., and G. C. P. King (1984), Seismic potential revealed by surface folding: 1983 Coalinga, California earthquake, *Science*, 224, 869–872.
- Storti, F., and J. Poblet (1997), Growth stratal architectures associated to decollement folds and fault-propagation folds. Inferences on fold kinematics, *Tectonophysics*, 282, 353–373.
- Streig, A. R., C. M. Rubin, W.-S. Chen, Y.-G. Chen, L.-S. Lee, S. C. Thompson, C. Madden, and S.-T. Lu (2007), Evidence for prehistoric coseismic folding along the Tsaotun segment of the Chelungpu fault near Nan-Tou, Taiwan, *J. Geophys. Res.*, doi:10.1029/2006JB004493, in press.
- Suppe, J. (1983), Geometry and kinematics of fault-bend folding, *Am. J. Sci.*, 283, 684–721.
- Suppe, J., and D. A. Medwedeff (1990), Geometry and kinematics of fault-propagation folding, *Eclogae Geol. Helv.*, 83, 409–454.
- Suppe, J., G. T. Chou, and S. C. Hook (1992), Rates of folding and faulting determined from growth strata, in *Thrust Tectonics*, edited by K. R. McClay, 105–121, CRC Press, Boca Raton, Fla.
- Suppe, J., F. Sábát, J. A. Muñoz, J. Poblet, E. Roca, and J. Vergés (1997), Bed-by-bed fold growth by kink-band migration, Sant Llorenç de Morunys, eastern Pyrenees, *J. Struct. Geol.*, 19, 443–461.
- Thompson, S. C., R. J. Weldon, C. M. Rubin, K. Abdрахmatov, P. Molnar, and G. W. Berger (2002), Late Quaternary slip rates across the central Tien Shan, Kyrgyzstan, central Asia, *J. Geophys. Res.*, 107(B9), 2203, doi:10.1029/2001JB000596.
- Van der Woerd, J., X. Xiwei, L. Haibing, P. Tapponnier, B. Meyer, F. J. Ryerson, A.-S. Meriaux, and X. Zhiqin (2001), Rapid active thrusting along the northwestern range front of the Tanghe Nan Shan (western Gansu, China), *J. Geophys. Res.*, 106(B12), 30,475–30,504.
- Vergés, J., V. A. Ramos, A. Meigs, E. Cristallini, F. H. Bettini, and J. M. Cortés (2007), Crustal wedging triggering recent deformation in the Andean thrust front between 31°S and 33°S: Sierras Pampeanas-Precordillera interaction, *J. Geophys. Res.*, 112, B03S15, doi:10.1029/2006JB004287.
- Weldon, R. (1986), The Late Cenozoic geology of Cajon Pass: Implications for tectonics and sedimentation along the San Andreas fault, Ph.D. thesis, Calif. Inst. of Technol., Pasadena.
- Willis, B. (1891), The mechanics of Appalachian structure, *U.S. Geol. Surv. 13th Annual Report 1891–1892, part 2*, 281 pp.
- Yielding, G., J. A. Jackson, G. C. P. King, H. Sinval, C. Vitafinzi, and R. M. Wood (1981), Relations between surface deformation, fault geometry, seismicity, and rupture characteristics during the El-Asnam (Algeria) earthquake of 10 October 1980, *Earth Planet. Sci. Lett.*, 56, 287–304.
- Yu, S. B., et al. (2001), Preseismic deformation and coseismic displacements associated with the 1999 Chi-Chi, Taiwan, earthquake, *Bull. Seismol. Soc. Am.*, 91, 995–1012.

J.-P. Avouac, Tectonics Observatory, California Institute of Technology, MC 100-23, 1200 E. California Blvd., Pasadena, CA 91125, USA.

J. F. Dolan, Department of Earth Sciences, University of Southern California, Los Angeles, CA 90089-0740, USA. (dolan@earth.usc.edu)