

**Origin of volatiles emitted by Plinian mafic eruptions of the Chikurachki volcano, Kurile arc, Russia: trace element, boron and sulphur isotope constraints**

Andrey A. Gurenko<sup>a,b\*</sup>, Alexander B. Belousov<sup>c</sup>, Vadim S. Kamenetsky<sup>d,e</sup>, Michael E. Zelenski<sup>e</sup>

<sup>a</sup> *Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA*

<sup>b</sup> *Centre de Recherches Pétrographiques et Géochimiques, Université de Lorraine, 54501 Vandoeuvre-lès-Nancy, France*

<sup>c</sup> *Institute of Volcanology and Seismology, Petropavlovsk-Kamchatsky, 683006, Russia*

<sup>d</sup> *School of Physical Sciences, University of Tasmania, Hobart, TAS 7001, Australia*

<sup>e</sup> *Institute of Experimental Mineralogy RAS, Chernogolovka, 142432, Russia*

Revised manuscript

for submission to *Chemical Geology*

12 September, 2017

**Components:** abstract (398 words; 2,628 characters with spaces),  
main text (9,672 words; 61,177 characters with spaces),  
11 figures, 2 tables,  
the list of references includes 178 citations.

\* **Corresponding author and present address:** Andrey A. Gurenko, Centre de Recherches Pétrographiques et Géochimiques, 15, rue Notre-Dame des Pauvres, BP 20, 54501 Vandoeuvre-lès-Nancy, France. Phone: +33 (0)3 83 59 48 75, Fax: +33 (0)3 83 51 17 98, E-mail: [agurenko@crpg.cnrs-nancy.fr](mailto:agurenko@crpg.cnrs-nancy.fr)

**Abstract** – Chikurachki is a 1816-m high stratovolcano on Paramushir Island, Kurile arc, Russia, which has repeatedly produced highly explosive eruptions of mafic composition. The present work is aimed at constraining the origin of volatile components (CO<sub>2</sub>, H<sub>2</sub>O, F, S, and Cl), along with B and S isotopic compositions in a series of phenocryst-hosted melt inclusions and groundmass glasses from basaltic andesite pyroclasts of the 1853, 1986, and prehistoric Plinian eruptions of the volcano. The ranges of volatile concentrations in melt inclusions (47–1580 µg/g CO<sub>2</sub>, 0.4–4.2 wt.% H<sub>2</sub>O, 399–633 µg/g F, 619–3402 µg/g S and 805–1240 µg/g Cl) imply a sudden pressure release from ~460 through ~35 MPa that corresponds to ~1.2–16-km-depth range of magma ascent upon decompression. We conclude that rapid ascent of the volatile-rich basaltic magmas from ~16-km initial depth accompanied by near-surface bubble nucleation and growth, and subsequent magma fragmentation appear to be a primary reason for the Plinian character of the Chikurachki eruptions. Significant negative correlations of S with K, Zr, Nb, Ba, La, Ce, Pr ( $R = -0.8$  to  $-0.9$ ), no clear relationships of S with H<sub>2</sub>O, CO<sub>2</sub> and Cl, but strong positive correlations of S/K<sub>2</sub>O with H<sub>2</sub>O/K<sub>2</sub>O, Cl/K<sub>2</sub>O and F/K<sub>2</sub>O preclude magma degassing to be the only process affecting volatile concentrations dissolved in the melt. The  $\delta^{34}\text{S}$  values of the studied inclusion and groundmass glasses range from  $-1.6$  to  $+12.3\%$ , decrease with decreasing S, show significant positive correlations with H<sub>2</sub>O/K<sub>2</sub>O, Cl/K<sub>2</sub>O and F/Zr, and negative correlations with a number of incompatible trace elements. Neither open- nor close-system magma degassing can account for the observed range of  $\delta^{34}\text{S}$ . The  $\delta^{11}\text{B}$  values of the melt inclusions range from  $-7.0$  to  $+2.4\%$  with 13–23 µg/g B. The relationships of  $\delta^{11}\text{B}$  with B/K<sub>2</sub>O and B/Nb are inconsistent with magma contamination at shallow crustal depths. Linear character of 1/S vs.  $\delta^{34}\text{S}$  relationship suggests two-component mixing. The possible mixing end-members could be the magmas having similar major and trace element compositions, but strongly contrasting volatile contents and S isotopes. Based on the behaviour of fluid-mobile vs. fluid-immobile incompatible trace elements, we conclude that the subduction component likely

represents a mixture of subduction sediment-derived melt with up to 60% of slab-derived fluid. Admixture of ~1–8% of the inferred subduction component to the depleted mantle wedge source is required to account for the compositional range of the Chikurachki melt inclusions, and ~0.4–10% to account for the composition of Kurile arc mafic magmas.

### Keywords

Plinian mafic eruptions, Chikurachki volcano, Melt inclusions, Magmatic volatiles, Stable isotopes, Ion microprobe

### Research Highlights

- Origin of volatiles that caused Plinian basaltic eruptions of Chikurachki was studied.
- Degassing cannot account for the observed ~14‰ range of  $\delta^{34}\text{S}$  in melt inclusions.
- Behaviour of volatiles and  $\delta^{11}\text{B}$  in inclusion glasses precludes crustal contamination.
- Volatiles and  $\delta^{34}\text{S}$  of the glasses were mainly controlled by two-component mixing.
- The subduction component was a 4:6-mixture of slab-derived melt and fluid.

## 1. Introduction

Strongly explosive (Plinian) eruptions are characterised by high eruption rates, high eruption columns and widely dispersed fallout deposits, representing an efficient mechanism to transport volatiles from the Earth's interior to the upper atmosphere (Walker, 1973, 1980; Fischer and Schmincke, 1984; Wilson and Walker, 1987; Francis, 1993). Injection of water, halogens and sulphur into the atmosphere leads to the formation of solid and liquid aerosols, which may significantly affect global climate and the stability of the ozone layer (Brasseur and Granier, 1992). Plinian eruptions are mainly associated with water-rich intermediate and silicic magmas (andesite through dacite), whereas highly explosive eruptions of mafic composition (i.e., resembling basalt or basaltic andesite) are less common. Examples of Plinian mafic eruptions occur at Klyuchevskoy, Tolbachik, Chikurachki and Tyatya in the Kamchatka and Kurile arcs, Russia (Fedotov and Markhinin, 1983; Ovsyannikov and Muravyev, 1992; Belousov et al., 2003, 2015, 2017; Hasegawa et al., 2011), Masaya and Cerro Negro, Nicaragua (Williams, 1983; Roggensack et al., 1997), Arenal, Costa Rica (Soto and Alvarado, 2006; Szramek et al., 2006), Llaima, Chile (Ruth et al., 2016), Sumisu Rift, Japan (Gill et al., 1990), Ambrym, Vanuatu (Robin et al., 1993), Tofua, Tonga (Caulfield et al., 2011), Taupo and Tarawera, New Zealand (Walker, 1980; Walker et al., 1994), Shishaldin, Alaska (Szramek et al., 2010) and Etna, Italy (Houghton et al., 2004; Sable et al., 2006). The lack of clear relationships between eruption rate and style often associating with rapid change of eruptive behaviour of a volcano suggest that many different parameters such as conduit geometry, magma viscosity, changing conditions of magma ascent and fragmentation, and even the dynamics of syn-eruptive microlite growth may strongly affect the eruption figure (e.g., Fischer and Schmincke, 1984; Carey and Sparks, 1986; Houghton et al., 2004; Szramek et al., 2006, 2010; Houghton and Gonnermann, 2008).

The present work is a follow-up study to Gurenko et al. (2005a) and focuses on the detailed investigation of the behaviour of volatile components (CO<sub>2</sub>, H<sub>2</sub>O, F, S, and Cl) along with boron and sulphur isotopic compositions of the previously analysed olivine-

orthopyroxene- and plagioclase-hosted melt inclusions and groundmass glasses erupted by the Chikurachki volcano. Volatile-rich slab-derived melts and fluids play a central role in the origin of arc magmatism by lowering the solidus of a peridotite mantle wedge and triggering partial melting (e.g., Gill, 1981; McCulloch and Gamble, 1991; Hawkesworth et al., 1993; Pearce and Peate, 1995). Being a major inventory and a transport media of volatile components from descending plate to the overlying mantle, these melts and fluids (along with the factors listed above) may also have a link to the eruption style of volcanoes occurring in the subduction zone (SZ) setting. Here we explore the role of two concurrent processes, namely, subduction-related volatile flux versus possible gain of volatile components due to shallow crustal contamination of magma, also in relation of their possible contribution to the enrichment/depletion of the magma by sulphur and boron isotopes. Our new data provide additional constraints on the origin of volatile components in the mafic magmas erupted by the Chikurachki volcano during Plinian prehistoric and recent, 1853 and 1986 eruptions.

Another key problem in understanding the origin of arc magmas is the uncertainty in the provenance of volatile components. For example, Cl and H<sub>2</sub>O are thought to be primarily (>95%) from the subducting slab (e.g., Straub and Layne, 2003), whereas S and F can originate from metasomatically enriched subarc mantle or come directly from the slab (e.g., Alt et al., 1993; Metrich et al., 1999; Straub and Layne, 2003; Churikova et al., 2007; Selvaraja et al., 2017). Moreover, with increasing of subduction depth, the slab becomes progressively depleted in H<sub>2</sub>O and Cl, and to a lesser extent in S and F, implying efficient fractionation of H<sub>2</sub>O and Cl from S and F (e.g., Chaussidon et al., 1987; Metrich et al., 1999; Straub and Layne, 2003). Understanding the origin of volatiles that triggers the Plinian eruption of Chikurachki may also contribute to our understanding of fluid dynamics and mass transport in SZ settings, as well as help in elucidating the relative contributions of different subduction components to the generation of arc magmas.

## 2. Geological setting and samples

Chikurachki (50°19'24"N 155°27'39"E, 1816 m.a.s.l) is an active stratovolcano on Paramushir Island, the northern Kuriles (**Fig. 1**). The volcanism of Chikurachki results from westward subduction of the Pacific plate under the Okhotsk plate. The Late Holocene stratocone of the volcano is built on a thick sequence of Pleistocene lava flows (Gorshkov, 1967). The recorded activity on Chikurachki started around 7500 BC. According to the Kamchatka Volcanic Eruption Response Team (KVERT), the most recent eruptions of 1958, 1961, 1964, 1973, 2002, 2003, 2004, 2005, 2007, 2008 were predominately of strombolian (mildly explosive), but recent Plinian (highly explosive) eruptions occurred in 1853, 1986 (**Fig. 2**), and the last one in 2015.

### 2.2. Samples studied

This study is based on the analyses of glassy lapilli with porphyritic texture, which came from the 1853 (sample CHK3/4), 1986 (sample CHK16) and two prehistoric (samples CHK13/1 and 13/2) tephra fallouts, resulting from documented, highly explosive eruptions of Chikurachki (Ovsyannikov and Muravyev, 1992; Belousov et al., 2003). We studied primary melt inclusions hosted by olivine, orthopyroxene and plagioclase phenocrysts, most of which were naturally quenched to glass. Chemical compositions of the studied inclusions and groundmass glasses (major and trace element, and H<sub>2</sub>O, S and Cl concentrations) were reported by Gurenko et al. (2005a). Briefly, the erupted melts range from low- to medium-K tholeiitic basalts and andesites, which are typical for Kurile lavas (**Fig. S2.1 – Supporting online material**). Variation diagrams of FeO, CaO and K<sub>2</sub>O vs. SiO<sub>2</sub> show that glass compositions follow simple trends of magma fractionation (Fig. 5 in Gurenko et al., 2005a). The phenocrysts are dominated by euhedral, elongated, twinned individual crystals of plagioclase (Pl, ~0.2–3.0 mm, 70–90 vol.%, An<sub>74–95</sub>) that may also form glomerocrysts. Phenocrysts of subhedral to rounded olivine (Ol, Fo<sub>72–78</sub>) and orthopyroxene (Opx, *mg*-number = 72–75), both of 0.2–0.5 mm size, and euhedral to resorbed

clinopyroxene (Cpx, 0.3–1.0 mm, *mg*-number = 71–77) are present in subordinate amounts. The groundmass is strongly vesiculated (20–30 vol.% vesicles) and consists of variable proportions of rounded olivine, prismatic augite and isometric Fe-Ti oxide crystals mixed with needle-like plagioclase microlites (up to 200  $\mu\text{m}$ ).

The studied inclusion and groundmass glasses are slightly enriched in light rare earth elements (LREE) relative to heavy rare earth elements (HREE); e.g.,  $[\text{La}/\text{Sm}]_n = 1.8\text{--}2.4$ , where the subscript  $n$  denotes the respective element concentrations normalised to those of N-MORB taken from Hofmann (1988) (**Fig. 3, Table S2.1** – *Supporting online material*). The glasses also show significant depletions in high-field-strength elements (HFSE) relative to LREE but nearly equal concentrations of HFSE relative to HREE normalised to N-MORB (e.g.,  $[\text{Nb}/\text{La}]_n = 0.19\text{--}0.29$ ,  $[\text{Zr}/\text{Sm}]_n = 0.52\text{--}0.76$ ,  $[\text{Y}/\text{Yb}]_n = 0.86\text{--}1.24$ ), and enrichments in Th and large-ion lithophile elements (LILE) relative to Nb (e.g.,  $[\text{Ba}/\text{Nb}]_n = 41\text{--}69$ ,  $[\text{Th}/\text{Nb}]_n = 7\text{--}30$ ,  $[\text{K}/\text{Nb}]_n = 15\text{--}25$  and  $[\text{Sr}/\text{Sm}]_n = 2.9\text{--}6.9$ ). The inclusions, and to lesser extent the groundmass glasses are characterised by relatively high contents of volatile components (the concentrations of B,  $\text{H}_2\text{O}$ , Cl and F shown in **Fig. 3** are discussed in section 4.1), displaying positive anomalies in the multi-element diagram relative to respective trace elements of similar incompatibility (e.g.,  $[\text{Cl}/\text{Nb}]_n = 21\text{--}54$ ,  $[\text{B}/\text{K}]_n = 2.3\text{--}5.4$ ,  $[\text{H}_2\text{O}/\text{Ce}]_n = 4.9\text{--}23$ ,  $[\text{F}/\text{Nd}]_n = 2.2\text{--}3.2$ ,  $[\text{F}/\text{Zr}]_n = 3.9\text{--}7.8$ ). These values are generally in, or slightly exceeding, the range of volatile concentrations reported for the Kurile island arc basalts and basaltic andesites ( $\text{MgO} > 2.5$  wt.%) by Bailey et al. (1987a,b), Ishikawa and Tera (1997), Ikeda (1998), Ikeda et al. (2000), Takagi et al. (1999), Ishikawa et al. (2001) (**Fig. 3**).

### 3. Analytical methods

Ion microprobe analyses were obtained with a CAMECA IMS 1280 ion microprobe at the Northeast National Ion Microprobe Facility (Woods Hole Oceanographic Institution, Woods

Hole, USA) (for detailed description of the employed analytical protocols, see *Supporting online material*).

## 4. Results

### 4.1. Volatile concentrations

The concentrations of CO<sub>2</sub>, H<sub>2</sub>O, F, S, and Cl in naturally quenched, glassy inclusions hosted in Ol, Opx and Pl are listed in **Table S2.1** (*Supporting online material*), along with H<sub>2</sub>O, S and Cl reported by Gurenko et al. (2005a). The inclusions have a wide range of volatile concentrations (39–653 µg/g CO<sub>2</sub>, 0.4–4.2 wt.% H<sub>2</sub>O, 428–638 µg/g F, 676–3450 µg/g S, and 805–1243 µg/g Cl) that corresponds to the volatile contents previously reported by Gurenko et al. (2005a). Correction for post-entrapment crystallisation (PEC) of host mineral (up to 7.8 wt.% Ol, 0.8 wt.% Opx, and no correction for Pl) results in the following ranges: 39–626 µg/g CO<sub>2</sub>, 0.4–4.2 wt.% H<sub>2</sub>O, 412–634 µg/g F, 639–3408 µg/g S, 805–1241 µg/g Cl, which appeared to be nearly the same (given the maintained analytical uncertainty) as the uncorrected volatile concentrations. This observation implies that the eruption and quenching of magma occurred shortly after the inclusions have been entrapped by growing phenocrysts. Thus, the concentrations of H<sub>2</sub>O and CO<sub>2</sub> measured in the studied melt inclusions reflect the last vapour-melt equilibrium in the magmatic system upon the eruption, returning a minimum pressure of magma fractionation. According to the *VolatileCalc* solution model (Newman and Lowenstern, 2002), they correspond to H<sub>2</sub>O-CO<sub>2</sub> gas pressure of 33 to 276 MPa or to the range of depth from ~1 to ~9.5 km, assuming average and constant density of ~2900 kg/m<sup>3</sup> of the upper crust (**Fig. 4A**).

Several recent studies (Esposito et al., 2011; Steele-Macinnis et al., 2011; Hartley et al., 2014; Moore et al., 2015; Wallace et al., 2015; Mironov et al., 2015; Aster et al., 2016) have demonstrated that 40 to 90% of the original CO<sub>2</sub> dissolved in the melt at the time of inclusion

entrapment can be lost to a shrinkage bubble during post-entrapment cooling. A correction algorithm described by Wallace et al. (2015) was therefore used to reconstruct the original CO<sub>2</sub> and H<sub>2</sub>O concentrations in the entrapped melts (see *Supporting online material*). These calculations revealed that ~5 to 69% of the original CO<sub>2</sub> and 0.1 to 3.2% of the original H<sub>2</sub>O could have been lost to the shrinkage bubble, in agreement with the results of the above studies. The following ranges of volatile concentrations were obtained after the correction: 47–1580 µg/g CO<sub>2</sub>, 0.4–4.2 wt.% H<sub>2</sub>O, 399–633 µg/g F, 619–3402 µg/g S and 805–1240 µg/g Cl (**Table S2.1** – *Supporting online material*). According to the *VolatileCalc* solution model (Newman and Lowenstern, 2002), the corrected results correspond to the H<sub>2</sub>O-CO<sub>2</sub> gas pressures between 35 and 456 MPa or to ~ 1.2–15.7-km-range of crustal depth (**Fig. 4B**).

The concentrations of S in melt inclusions are shown in comparison with mid-ocean ridge basalts (MORB) and ocean island basalts (OIB) (**Fig. 5A**). Many of the Chirurachki glass inclusions have similar or higher S concentrations than those of MORB or OIB, whereas all groundmass glasses are depleted in S (<350 µg/g), consistent with sulphur loss presumably by degassing. However, sulphur in the inclusion glasses shows significant negative correlations with a number of trace and rare earth incompatible elements, such as K (–0.81), Ti (–0.59), Y (–0.61), Zr (–0.82), Nb(–0.71), Ba (–0.83), La (–0.79), Ce (–0.88), Pr (–0.77), Nd (–0.71), Sm (–0.68), Gd (–0.56), Dy (–0.73), Ho (–0.53), Er (–0.73), Yb (–0.66), Lu (–0.71), Th (–0.66), where the respective correlation coefficient (*R*) values are given in brackets and *N* = 21, if not specified (**Fig. 5B** and **Table S2.1** – *Supporting online material*). All these correlations are significant at 95% confidence level (for *N* = 18 and 21, a number of points in the correlations, a critical value of the Pearson correlation coefficient at the 0.05-level of significance for a two-tailed test is 0.47 and 0.43, respectively). There are no significant correlations of S with H<sub>2</sub>O, Cl and F. We observe, however, strong positive correlation between these elements, if their concentrations are normalised to K<sub>2</sub>O (or any other melt- and fluid-immobile incompatible element) that minimises

possible effects of partial melting and/or magma fractionation (e.g.,  $R = 0.83$  for S/K<sub>2</sub>O vs. H<sub>2</sub>O/K<sub>2</sub>O correlation, **Fig. 5C**,  $R = 0.80$  for S/K<sub>2</sub>O vs. Cl/K<sub>2</sub>O, **Fig. S2.1A** – *Supporting online material*,  $R = 0.85$  for S/K<sub>2</sub>O vs. F/K<sub>2</sub>O, **Fig. 5D**, and  $R = 0.88$  for Cl/K<sub>2</sub>O vs. H<sub>2</sub>O/K<sub>2</sub>O, **Fig. 5E**). Chlorine does not correlate with incompatible trace elements, except for Sr ( $R = -0.56$ ; **Fig. S2.1B** – *Supporting online material*), shows a significant correlation with fluorine ( $R = 0.70$ ; **Fig. S2.1C** – *Supporting online material*), which becomes much stronger, if we consider F/Zr vs. Cl/Nb ratios ( $R = 0.88$ ; **Fig. 5F**).

#### 4.2. Sulphur isotopes

Sulphur isotopic compositions of the studied melt inclusions and one groundmass glass are given in **Table S2.1** (*Supporting online material*) and presented in **Fig. 6**. The  $\delta^{34}\text{S}$  values are given in permil units (‰) relative V-CDT (the Vienna Canyon Diabolo Troilite,  $^{34}\text{S}/^{32}\text{S} = 0.0441626 \pm 0.0000039$ ; Ding et al., 2001). The  $\delta^{34}\text{S}$  values of the inclusion glasses range from  $-1.6$  to  $+12.3\%$ , and decrease with decreasing S concentrations ( $R = 0.85$ ; **Fig. 6A**). The S vs.  $\delta^{34}\text{S}$  relationship has a hyperbolic shape and is transformable into a strait line in the  $1/\text{S}-\delta^{34}\text{S}$  space ( $R = -0.84$ ; **Fig. 6B**), pointing towards two-component mixing scenario (Langmuir et al., 1978), though the effect of magma degassing cannot be completely excluded (see further discussion). The sample 13/1 matrix glass has  $\delta^{34}\text{S} = +0.39 \pm 0.43\%$ , 1 SE, contains very low S concentration (107  $\mu\text{g/g}$  S) and strongly deviates from the linear array defined by melt inclusions (**Fig. 6B**), being in contrast strongly degassed. Also, the  $\delta^{34}\text{S}$  values show significant positive correlations with H<sub>2</sub>O (0.54), CO<sub>2</sub> (0.55) and Sr (0.5), as well as negative correlations with light lithophile elements Li ( $-0.53$ ,  $N = 10$ ), Be ( $-0.73$ ,  $N = 10$ ) and B ( $-0.6$ ), and other incompatible elements, such as K ( $-0.8$ ), Zr ( $-0.71$ ), Nb ( $-0.64$ ), Ba ( $-0.75$ ), La ( $-0.69$ ), Ce ( $-0.72$ ), Pr ( $-0.57$ ), Sm ( $-0.63$ ), Gd ( $-0.76$ ), Dy ( $-0.71$ ), Tb ( $-0.49$ ), Er ( $-0.8$ ), Lu ( $-0.79$ ), Th ( $-0.53$ ), where the respective correlation coefficient ( $R$ ) values are given in brackets and  $N = 17$ , if not

specified, and the critical value of the Pearson correlation coefficient is 0.48. The significance of the correlations of  $\delta^{34}\text{S}$  with volatile components becomes higher, if volatile concentrations are normalised to incompatible elements as, for instance,  $\text{H}_2\text{O}/\text{K}_2\text{O}$  (0.79),  $\text{Cl}/\text{K}_2\text{O}$  (0.83) (**Fig. 6B,C**),  $\text{F}/\text{Zr}$  (0.84, **Fig. S2.1D** – *Supporting online material*).

#### 4.3. Boron concentrations and B isotopic composition

The concentrations of boron (12.5–22.9  $\mu\text{g/g}$  measured, and 12.5–21.7  $\mu\text{g/g}$  corrected for PEC) are similar to those previously reported by Gurenko et al. (2005a) (16.1–18.7  $\mu\text{g/g}$  B) (**Table S2.1** – *Supporting online material*). The B/K ratios ranging from 0.0025 to 0.0057 are systematically higher than those of primitive N-MORB (B/K of  $\sim 0.001$ ) and OIB (B/K of  $\sim 0.0003$ ) (Ryan and Langmuir, 1993; Chaussidon and Jambon, 1994; Gurenko and Chaussidon, 1997) but are in the midrange of B concentrations reported for Kurile arc magmas with  $\text{SiO}_2 < 60$  wt.% (4.5–36.6  $\mu\text{g/g}$  B; Ryan et al., 1995; Ishikawa and Tera, 1997). Very similar concentrations of B in the Chikurachki lavas (14.4–15.5  $\mu\text{g/g}$ ) were found, in particular, by Ishikawa and Tera (1997), who also noted a clear across-arc variation of B/Nb ratios (and  $\delta^{11}\text{B}$ , as well). At the slab depth of  $\sim 140$  km corresponding to that of the Chikurachki volcano (e.g., Tatsumi et al., 1994; Ishikawa and Tera, 1997; Ishikawa et al., 2001 and references therein), the B/Nb ratios in the Kurile lavas vary from  $\sim 6$  to 18 (Ishikawa and Tera, 1997), and the present B/Nb range obtained for the studied melt inclusions (from  $\sim 9$  to  $\sim 22$ ) is in agreement with these data.

The  $\delta^{11}\text{B}$  values of the studied naturally quenched olivine-hosted melt inclusions range from  $-7.0$  to  $+2.4\text{‰}$ , calculated relative to NBS 951 standard with  $^{11}\text{B}/^{10}\text{B} = 4.04558 \pm 0.00033$  (Spivack and Edmond, 1986) (**Table S2.1** – *Supporting online material*). The range is similar but extends to more negative values than those of the Kurile arc lavas (from  $-3.8$  to  $+5.9\text{‰}$ ; Ishikawa and Tera, 1997), being however within a broad interval of  $\delta^{11}\text{B}$ -values of arc magmas worldwide, from  $-21.3\text{‰}$  (e.g., Mt. Shasta; Rose et al., 2001) to  $+12\text{‰}$  (e.g., Izu arc; Straub and

Layne, 2002). In contrast to S isotopes,  $\delta^{11}\text{B}$  values show neither a correlation with the concentrations of B or 1/B values nor with major and incompatible trace elements. Similarly, a wide  $\delta^{11}\text{B}$  range of  $\sim 20\%$  (from  $-21.3$  to  $-1.1\%$  at relatively narrow range of boron concentrations ( $0.67\text{--}1.64$   $\mu\text{g/g}$  B) showing no clear correlation between each other was described in the Mt. Shasta basaltic andesite rocks (Rose et al., 2001).

## 5. Discussion

### 5.1. $\text{H}_2\text{O}\text{-Cl}\text{-B}$ and $\delta^{11}\text{B}$ relationships and the magma contamination scenario

Assimilation of crustal rocks is a common process that can significantly modify pristine chemical and isotopic compositions of parental arc magmas (e.g., Davidson, 1987; Thirlwall et al., 1996; Hickey et al., 1986; Turner et al., 1997; Macpherson and Matthey, 1998; Vroon et al., 2001; Bindeman et al., 2004; Walowski et al., 2016). Excess of halogens in mafic magmas, and especially coherent variation of  $\text{H}_2\text{O}$  and Cl was shown to be a good proxy to assess the role of assimilation of oceanic crust and/or addition of NaCl-rich brines or fluids during magma fractionation (e.g., Anderson, 1974; Michael and Schilling, 1989; Jambon et al., 1995; Michael and Cornell, 1998; Kent et al., 1999; Lassiter et al., 2002; Simons et al., 2002; le Roux et al., 2006; Shimizu et al., 2009; Gurenko and Kamenetsky, 2011; Kendrick et al., 2013; Cabral et al., 2014). However, as noted by le Roux et al. (2006), this approach is based on the assumption that the concentration of  $\text{K}_2\text{O}$  in the magma remains unaffected by later processes. These authors recommend using Cl/Nb and  $\text{H}_2\text{O}/\text{Ce}$  ratios because Nb and Ce have similar incompatible behaviour to Cl and  $\text{H}_2\text{O}$ , respectively, but are less affected by alteration, as compared to K.

In our case, a positive correlation of  $\text{H}_2\text{O}/\text{K}_2\text{O}$  with  $\text{Cl}/\text{K}_2\text{O}$  in the Chikurachki melt inclusions (**Fig. 5E**) can be interpreted in favour of magma contamination. However, strong positive correlations between  $\text{H}_2\text{O}/\text{K}_2\text{O}$  and  $\text{H}_2\text{O}/\text{Ce}$  ( $R = 0.97$ , not shown) and between  $\text{Cl}/\text{K}_2\text{O}$  and  $\text{Cl}/\text{Nb}$  ( $R = 0.88$ , not shown) suggest that K, Nb and Ce appear to behave coherently during

magma fractionation. Furthermore, the concentrations of K are unlikely to have been compromised and could be equally used for normalisation. The results of concurrent modelling of magma contamination using different compositions of probable crustal contaminants are shown in **Fig. 7** and **8**. In the modelling, we use the same element concentrations and  $\delta^{11}\text{B}$  values of seawater (SW), 15%- and 50%-NaCl brine (15%-NaCl and 50%-NaCl, respectively), and altered oceanic crust (AOC) mixing end-members, as in Gurenko and Kamenetsky (2011) (**Table 1**). Given the narrow range of incompatible element concentrations in the studied melt inclusions (**Fig. 3**), the concentrations of  $\text{K}_2\text{O}$  (0.804 wt.%), Nb (1.28  $\mu\text{g/g}$ ) and Ce (15.37  $\mu\text{g/g}$ ) of a presumably “uncontaminated” arc magma (UAM) were defined as average of all melt inclusions from **Table S2.1 – Supporting online material** (using PEC-corrected concentrations). The boron content and B isotopic composition of UAM (0.9  $\mu\text{g/g}$  B,  $\delta^{11}\text{B} = -3.7\text{‰}$ ) corresponding to back-arc basin basalts presumably unaffected by later shallow-crustal contamination were taken from Chaussidon and Marty (1995). The concentrations of  $\text{H}_2\text{O}$  and Cl in UAM were calculated using those of the neighbouring elements of similar incompatibility but showing no anomalies in the multi-element diagram (**Fig. 3**) to be in the lower left corner of the respective panels in **Fig. 7**, thereby matching the lower end of the trends defined by the inclusion compositions:  $[\text{H}_2\text{O}]_n = ([\text{Ce}]_n \times [\text{Pr}]_n)^{0.5}$ , equal to 0.256 wt.%  $\text{H}_2\text{O}$ , and  $[\text{Cl}]_n = ([\text{Nb}]_n \times [\text{K}]_n)^{0.5}$ , equal to 128.5  $\mu\text{g/g}$  Cl. The chemical and isotopic compositions of seawater (SW) is relatively uniform and well constrained, whereas the composition of seawater-derived NaCl-brines can be calculated from that of SW (**Table 1**). In contrast, the composition of altered oceanic crust (AOC) is variable. In particular, AOC from Gurenko and Kamenetsky (2011) is significantly more  $\text{H}_2\text{O}$ -, Cl- and  $\text{K}_2\text{O}$ -rich than that used for similar calculations by le Roux et al. (2006). In order to account for possibly wider ranges of  $\text{H}_2\text{O}$ , Cl, Nb and Ce, we considered two compositions of AOC: AOC1 from Gurenko and Kamenetsky (2011) and AOC2 from le Roux et al. (2006) (**Table 1**). Finally, we assigned a composition of siliceous marine sediment (SED),

which is considered as one possible, if not a single contamination component to account for the negative  $\delta^{11}\text{B}$  melt inclusion values, with 10.1 wt.%  $\text{H}_2\text{O}$ , 1.7 wt.%  $\text{K}_2\text{O}$ , 7.67  $\mu\text{g/g}$  Nb, 50.11  $\mu\text{g/g}$  Ce. The concentrations of  $\text{H}_2\text{O}$ ,  $\text{K}_2\text{O}$ , Nb and Ca used in calculations are those of the average sediment column of the Kurile trench from Plank and Langmuir (1998), and 1000  $\mu\text{g/g}$  Cl, 120  $\mu\text{g/g}$  B, and  $\delta^{11}\text{B} = -10\text{‰}$  were taken from Gurenko and Kamenetsky (2011), given that siliceous terrigenous sediments may be as low as  $-15\text{‰}$  of  $\delta^{11}\text{B}$  (Leeman and Sisson, 1996).

As noted by Gurenko and Kamenetsky (2011), it is not possible to unequivocally prove magma contamination only on the basis of  $\text{H}_2\text{O}/\text{K}_2\text{O}$  vs.  $\text{Cl}/\text{K}_2\text{O}$  relationships (or using Cl- and  $\text{H}_2\text{O}$ -to-incompatible-element normalised ratios). Similarly, if we consider solely the  $\text{H}_2\text{O}/\text{K}_2\text{O}$  vs.  $\text{Cl}/\text{K}_2\text{O}$  or  $\text{H}_2\text{O}/\text{K}_2\text{O}$ -B/ $\text{K}_2\text{O}$  variations in melt inclusions (equally,  $\text{H}_2\text{O}/\text{Ce}$  vs. Cl/Nb or  $\text{H}_2\text{O}/\text{Ce}$  vs. B/Nb), they can be explained by contamination of magma by seawater or NaCl-rich brines (**Fig. 7**). On the other hand, if in addition to  $\text{H}_2\text{O}$  and Cl we consider boron content and B-isotope data then the relationships between  $\text{Cl}/\text{K}_2\text{O}$  and B/ $\text{K}_2\text{O}$  (equally, Cl/Nb vs. B/Nb; **Fig. S2.2C – Supplementary online material**), or B/ $\text{K}_2\text{O}$  (or B/Nb) vs.  $\delta^{11}\text{B}$  (**Fig. 8**) confirm neither significant contribution of these components, nor the contribution of AOC1, AOC2 and SED components. Indeed, to account for the compositions of melt inclusions with strongly and coherently varying  $\text{H}_2\text{O}/\text{K}_2\text{O} = 0.3\text{--}8.9$  and  $\text{Cl}/\text{K}_2\text{O} = 0.06\text{--}0.24$ , it would require contamination either by SW (up to 7%) or more concentrated NaCl-brine, or assimilation of AOC1 (up to 60%) or both AOC1 + SW components (**Fig. 7A**). No significant effect of 15%-NaCl and 50%-NaCl brine, as well as the role of AOC2 and SED can be recognised in this diagram. The same conclusions can be derived from the  $\text{H}_2\text{O}/\text{Ce}$  vs. Cl/Nb variations (**Fig. 7B**), with only a difference that contamination by AOC1 (up to 70%) is required. However, if we consider mixing of the same components in coordinates  $\text{H}_2\text{O}/\text{K}_2\text{O}$  vs. B/ $\text{K}_2\text{O}$  or  $\text{H}_2\text{O}/\text{Ce}$  vs. B/Nb (**Fig. 7C,D**), then the role of SW and AOC1, along with 15%-NaCl and AOC2 is not yet justified, but the 50%-NaCl end-member appears to be potentially significant. The amount of SED to add to UAM

appears to be up to 50% in **Fig. 7C** and up to 100% in **Fig. 7D**, both being unrealistic. Finally, if we consider B/K<sub>2</sub>O vs.  $\delta^{11}\text{B}$  or B/Nb vs.  $\delta^{11}\text{B}$  systematics (**Fig. 8**), then the role of seawater, NaCl-brines and AOC1 is not confirmed at all, while the amount of AOC2 and SED to be assimilated approaches to 100% (**Fig. 8B**).

In conclusion we note that Cl and F often behave as incompatible elements during partial melting and magma fractionation (e.g., Fischer and Schmincke, 1984; Carroll and Webster, 1994; Jambon, 1994; Dalou et al., 2014). As shown by Dalou et al. (2014 and references therein), a Cl/F ratio of  $>1$  is expected in arc magmas resulting from partial melting of a depleted mantle source mixed with an aqueous dehydration fluid because the Cl/F ratio in the subduction slab fluids is usually high (e.g.,  $\sim 4$  in Mount Shasta rocks and up to 9.5 in Izu Bonin arc magmas). In contrast, the low Cl/F ratios ( $<1$ ; e.g., Central and South American arcs, Kermadec) suggest that the magmas were produced by melting of a mantle source mixed with a hydrous slab melt. In this context, it is essential to note that Cl/F ratios in the studied melt inclusions range from 1.5 to 2.0, pointing towards predominant role of the subduction slab fluid in the origin of Chickuracki magmas. Finally, the observed relationships of S/K<sub>2</sub>O with K<sub>2</sub>O-normalised concentrations of H<sub>2</sub>O, Cl and F (**Fig. 5C,D** and **Fig. 2.1A** – *Supporting online material*), as well as the H<sub>2</sub>O/K<sub>2</sub>O vs. Cl/K<sub>2</sub>O and F/Zr vs. Cl/Nb correlations (**Fig. 5E,F**) strongly suggest that the variations of H<sub>2</sub>O, S, Cl and F contents were not primarily affected by degassing or contamination, but more likely related to the transport of these components to magma source region by subduction slab fluids, in agreement with the previous studies of the Kurile-Kamchatka arc volcanism (e.g., Hochstaedter et al., 1996; Ishikawa and Tera, 1997; Kepezhinskas et al., 1997; Dorendorf et al., 2000; Churikova et al., 2001, 2007; Ishikawa et al., 2001; Portnyagin et al., 2007, 2015, Konrad-Schmolke et al., 2016).

## 5.2. Origin of sulphur concentration and S-isotope ranges

Sulphur solubility in basaltic magmas was shown to be mostly a function of iron content of the melt, if sulphur is predominately present in sulphide ( $S^{2-}$ ) form (e.g., Connolly and Haughton, 1972; Haughton et al., 1974; Mathez, 1976; Carroll and Rutherford, 1985; Wallace and Carmichael, 1992; ), but it also depends strongly on temperature, pressure, redox conditions,  $H_2O$  content, and the composition of coexisting melt and vapour phase (Nagashima and Katsura, 1973; Katsura and Nagashima, 1974; Carroll and Rutherford, 1985, 1987; Luhr, 1990; Wallace and Carmichael, 1992; Mavrogenes and O'Neill, 1999; Matjuschkin et al., 2016; Canil and Fellows, 2017; Mavrogenes and Blundy, 2017). The solubility of sulphur is generally higher (at a given  $f_{O_2}$ ) in mafic (tholeiite and hawaiite), as compared to more  $SiO_2$ -rich magmas (rhyodacite) (e.g., Katsura and Nagashima, 1974).

As a sulphide-saturated melt becomes progressively oxidised, higher amount of sulphur is required to be dissolved in the melt to cause saturation of the melt with sulphide (e.g., Carroll and Webster, 1994 and references therein). This is also evident from the exponential character of FeO-S relationships observed in the more S-rich inclusions (**Fig. 5A**). The sulphur valence state changes from  $S^{2-}$  to  $S^{6+}$  within a limited range of oxygen fugacity, from NNO-1 to NNO+2 (Carroll and Rutherford, 1988; Wallace and Carmichael, 1994; Metrich and Clocchiatti, 1996; Gurenko and Schmincke, 2000; Jugo et al., 2005a,b, 2010; Fiege et al., 2015). Further melt oxidation causes a sulphide phase to become unstable. At these and more oxidised conditions (i.e.,  $f_{O_2}$  is greater than NNO+2), anhydrite ( $CaSO_4$ ) may become stable and appear as phenocrysts in the magma, as it was observed, for example, in some El Chichón and Mount Pinatubo trachyandesite and dacite magmas (Luhr et al., 1984; Varekamp et al., 1984; Carroll and Rutherford, 1987; Luhr, 1990; Bernard et al., 1991; Kress, 1997). Moreover, the solubility of sulphate ( $SO_4^{2-}$ ) is much higher than that of sulphide ( $S^{2-}$ ) in mafic magmas and can reach up to 2.5 wt.%  $SO_3$  (Carroll and Rutherford, 1985, 1987; Luhr, 1990; Metrich and Clocchiatti, 1996; Gurenko and Schmincke, 2000; Jugo et al., 2005a,b; Fiege et al., 2015). Since S concentrations

in many of our inclusion glasses significantly exceed the level of  $S^{2-}$  solubility, and they show no correlation with FeO (**Fig. 5A**), we contend that a significant proportion of sulphur is dissolved as  $SO_4^{2-}$  species. This is consistent with generally higher oxidation state of subduction-zone-related magmas, and also with previously estimated oxygen redox conditions of the Chikurachki magmas (i.e.,  $\Delta NNO = +1$  to  $+2$ ; Gurenko et al., 2005a). According to the relationship given in Wallace and Carmichael (1994):

$$\log (X_{SO_4^{2-}}/X_{S^{2-}}) = a \log fO_2 + b/T + c \quad (1)$$

where  $X_{SO_4^{2-}}$  and  $X_{S^{2-}}$  are mole fractions of  $SO_4^{2-}$  and  $S^{2-}$  sulphur species in the melt (i.e.,  $X_{SO_4^{2-}} + X_{S^{2-}} = 1$ ),  $T$  is temperature in Kelvin, and coefficients  $a = 1.02$ ,  $b = 25410$ , and  $c = -10$ , between 80 and 98% sulphur dissolved as  $SO_4^{2-}$  is expected in the basaltic melts representing the studied Plinian 1986, 1853 and prehistoric eruptions of the Chikurachki volcano, thereby explaining the elevated concentrations of sulphur in the studied melt inclusions.

The fact that sulphur isotopic composition (i.e.,  $\delta^{34}S$ ) strongly correlates with S concentrations in the melt inclusions and groundmass glasses (**Fig. 6A,B**), as well as with  $K_2O$ -normalised concentrations of  $H_2O$  and Cl (**Fig. 6C,D**) suggests that sulphur was more likely introduced to the magma by the same process along with  $H_2O$  and Cl. One possible and relatively simple explanation would be a contamination of magma by sulphate-bearing seawater-derived material at shallow-crustal depth, as it was previously demonstrated on example of Miocene basaltic hyaloclastites drilled during the ODP Leg 157 southwest of Gran Canaria (Canary Islands, Spain) by Gurenko et al., 2001. However, as discussed above, shallow-depth contamination was not a factor controlling budget of volatile components in the studied Chikurachki magmas. Thus, the origin of sulphur and its S-isotope signature could more likely

be ascribed to slab-derived fluids. Whether a consequent degassing of the magmas could result in the observed, nearly 14‰-range of  $\delta^{34}\text{S}$  values is a subject of further discussion.

### 5.2.1. S-isotope fractionation during magma degassing

Magma degassing is an alternative process, which is capable to significantly affect the concentration and isotopic composition of sulphur in the melt by changing the proportion of the reduced ( $\text{S}^{2-}$ ) and oxidised ( $\text{SO}_4^{2-}$ ) forms (e.g., Taylor, 1986; Marini et al., 2011; Mandeville et al., 2009 and references therein). This is mostly because sulphide and sulphate species may possess strongly contrasting  $\delta^{34}\text{S}$  values, with an average fractionation factor of  $+7.3 \pm 1.4\%$  between sulphate and sulphide (e.g., Sakai et al., 1982, 1984). According to Holloway and Blank (1994), the variations of  $\delta^{34}\text{S}$  values resulted from continuous volatile exsolution can be modelled either as closed-system (Eq. 2) or as open-system (i.e., Rayleigh distillation) degassing (Eq. 3):

$$\delta^{34}\text{S}_f = \delta^{34}\text{S}_i + 1000 (1 - F) \ln \alpha_{\text{gas-melt}} \quad (2)$$

$$\delta^{34}\text{S}_f = (\delta^{34}\text{S}_i + 1000) (F^{\alpha_{\text{gas-melt}} - 1}) - 1000 \quad (3)$$

where subscripts  $f$  and  $i$  refer to the final and initial isotopic compositions of S in the melt, respectively,  $F$  is a fraction of sulphur remaining in the melt upon degassing, and  $\alpha_{\text{gas-melt}}$  is a gas-melt S-isotope fractionation factor defined as:

$$\alpha_{\text{gas-melt}} = ({}^{34}\text{S}/{}^{32}\text{S})_{\text{gas}}/({}^{34}\text{S}/{}^{32}\text{S})_{\text{melt}} \quad (4)$$

To calculate the parameter  $F$  in Eq. 2 and 3, we used S-concentrations in final ( $[S]_f$ ) and initial ( $[S]_i$ ) melts normalised to a concentration of an incompatible trace element (metal;  $[Me]_f$  and  $[Me]_i$ , respectively) by the following equation:

$$F = ([S]_f \times [Me]_i) / ([S]_i \times [Me]_f) \quad (5)$$

using, in particular, the concentrations of Ce and K in the respective melts, and then the average was used. Two melt inclusions with the highest S concentrations and  $\delta^{34}\text{S}$  values (3/4-4-1 and 13/2-1-2, PEC- and exsolution-corrected values; **Table S2.1** – *Supporting online material*) were taken to calculate the initial melt composition ( $[S]_i = 3287 \mu\text{g/g}$ ,  $[\text{Ce}]_i = 9.54 \mu\text{g/g}$ ,  $[\text{K}]_i = 3592 \mu\text{g/g}$ , and  $\delta^{34}\text{S}_i = +11.2\%$ ).

Modelling of S isotopic fractionation caused by degassing is rather complicated because  $\alpha_{\text{gas-melt}}$  can be either higher or lower than unity because it depends strongly on  $T$ ,  $f_{\text{O}_2}$  and the proportion of  $\text{SO}_4^{2+}$  and  $\text{S}^{2-}$  species dissolved in the melt. Generally, a fractionation factor of  $^{32}\text{S}$  and  $^{34}\text{S}$  isotopes between gas phase and silicate melt ( $\Delta^{34}\text{S}_{\text{gas-melt}}$ ) can be expressed as following (e.g., Sakai et al., 1982), assuming that isotopic equilibrium is maintained among all sulphur species in the gaseous and melt phases:

$$\begin{aligned} \Delta^{34}\text{S}_{\text{gas-melt}} &= \delta^{34}\text{S}_{\text{gas}} - \delta^{34}\text{S}_{\text{melt}} \approx 1000 \ln \alpha_{\text{gas-melt}} = \\ &= [X\delta^{34}\text{S}_{\text{SO}_2} + (1 - X)\delta^{34}\text{S}_{\text{H}_2\text{S}}] - [Y\delta^{34}\text{S}_{\text{sulfide}} + (1 - Y)\delta^{34}\text{S}_{\text{sulfate}}], \end{aligned} \quad (6)$$

where  $X$  is the mole fraction of  $\text{SO}_2$  in the  $\text{SO}_2$ - $\text{H}_2\text{S}$  gas phase (i.e.,  $X_{\text{SO}_2} + X_{\text{H}_2\text{S}} = 1$ ), and  $Y$  is the mole fraction of sulphur dissolved in the melt as sulphide, and  $1 - Y$  mole fraction of sulphur dissolved as sulphate. After multiplying and collecting terms, Eq. 6 can be expressed as:

$$1000 \ln \alpha_{\text{gas-melt}} = X 1000 \ln \alpha_{\text{SO}_2\text{-H}_2\text{S}} + Y 1000 \ln \alpha_{\text{sulphate-sulphide}} - 1000 \ln \alpha_{\text{sulphate-H}_2\text{S}} \quad (7)$$

The temperature dependences of the fractionation factors in Eq. 7 can be approximated by the relationships studied by Richet et al. (1977) and Miyoshi et al. (1984) (for more detail see a compilation by Taylor, 1986):

$$1000 \ln \alpha_{\text{SO}_2\text{-H}_2\text{S}} = -0.42 \times 10^9/T^3 + 4.367 \times 10^6/T^2 - 0.105 \times 10^3/T - 0.41 \quad (8)$$

$$1000 \ln \alpha_{\text{sulphate-sulphide}} = 7.4 \times 10^6/T^2 - 0.19 \quad (9)$$

$$1000 \ln \alpha_{\text{sulphate-H}_2\text{S}} = 6.5 \times 10^6/T^2 \quad (10)$$

where  $T$  is temperature in Kelvin. For the temperature range from 910 to 1180°C determined for the studied Chikurachki melt inclusions (Gurenko et al., 2005a), the  $\Delta^{34}\text{S}_{\text{gas-melt}}$  fractionation factor is within the range of  $-2.1\text{‰}$  to  $-0.9\text{‰}$  (i.e.,  $\alpha_{\text{gas-melt}} = 0.9979\text{--}0.9991$ ). The calculations were done assuming  $X = 0.98$  because usually the volcanic gases emitted at temperatures  $\sim 1000^\circ\text{C}$  contain mostly  $\text{SO}_2$ , whereas  $\text{H}_2\text{S}$  is present only in trace amount, as shown for Kilauea volcano, Hawaii, and Mnt. Etna, Sicily (Symonds et al., 1994) and according to the recent gas-melt equilibria model of Burgisser and Scaillet (2007) for oxidised (i.e.,  $\Delta\text{NNO} = +1.5$ ) conditions. The variations of  $X$  within  $\pm 10\%$  (from 0.8 to 1) will cause an uncertainty in the resulting  $\Delta^{34}\text{S}_{\text{gas-melt}} \approx \pm 0.2\text{--}0.3\text{‰}$ . The  $Y$  value varied from 0.02 to 0.2, in agreement with the previously reported  $\Delta\text{NNO}$ -range from +1 to +2, resulting in 80 to 98% sulphur dissolved as sulphate in the melt (see above).

The diagram of  $\delta^{34}\text{S}$  versus  $F$  (**Fig. 9**) demonstrates that neither open- nor close-system degassing can account for the observed  $\sim 14\text{‰}$  range in the Chikurachki melt inclusions. This is mostly because the outgassed  $\text{SO}_2$  will preferentially drive out the lighter  $^{32}\text{S}$  isotope under

conditions of high  $f_{O_2}$  so that the remaining melt will be progressively enriched in  $^{34}\text{S}$ , as it was first shown by Sakai et al. (1982) and also demonstrated by Mandeville et al. (2009) for the case of climatic and pre-climatic eruptions of Mt. Mazama, Crater Lake, Oregon, but being completely opposite to what is observed in **Fig. 6A** and **Fig. 9**. Negative correlations of S with incompatible trace elements mentioned above (**Fig. 5B**) may suggest that magma fractionation (resulting in enrichment of melt by incompatible trace elements) might have been accompanied by S-depletion due to degassing. However, a weak correlation of S with  $\text{CO}_2$  and no correlations with other volatile components (like  $\text{H}_2\text{O}$  and Cl) imply that degassing, though being probably important, was not a single process affecting the concentrations of the dissolved S. Furthermore, the fact that the hyperbola-shaped relationship of  $\delta^{34}\text{S}$  vs. S concentrations (**Fig. 6A**) is transformable into a straight line in the  $1/\text{S}-\delta^{34}\text{S}$  space (**Fig. 6B**) represents a strong argument in favour of two-component mixing hypothesis (Langmuir et al., 1978), whereas host lapilli groundmass glasses, which presumably suffered degassing, do not fit the line. We thus contend that sulphur abundances, the entire S-isotope range, as well as the other volatile components such as  $\text{H}_2\text{O}$ , Cl and F correlating with each other (**Fig. 5**) were largely controlled by two-component mixing process. The possible mixing end-members could be the magmas having similar major and trace element compositions but strongly contrasting volatile contents and S isotopes.

### 5.3. Trace element constraints on magma source components

Mafic arc magmatism results from melting of a mantle wedge, whose solidus is lowered by input of slab-derived hydrous melts and/or fluids (e.g., Gill, 1981; McCulloch and Gamble, 1991; Hawkesworth et al., 1993; Woodhead et al., 1993; Pearce and Peate, 1995). The composition of mantle wedge is relatively well constrained and thought to be represented by a MORB-type, or even more depleted upper mantle material (e.g., Woodhead et al., 1993;

Kepezhinskas et al., 1997). However, a relative contribution of slab-derived H<sub>2</sub>O-rich fluids and/or melts, as well as the proportion of the sediments relative to the mafic part of the oceanic crust in the slab remains enigmatic. Numerous previous studies attempted to assess the effect of fluid transport vs. slab melting in the origin of the Kurile-Kamchatka arc magmas (e.g., Hochstaedter et al., 1996; Ishikawa and Tera, 1997; Kepezhinskas et al., 1997; Khubunaya and Sobolev, 1998; Dorendorf et al., 2000; Churikova et al., 2001, 2007; Ishikawa et al., 2001; Portnyagin et al., 2007, 2015, Konrad-Schmolke et al., 2016, among others). One of the main purposes of the present study is to constrain a contribution of the above mentioned components to the origin of the Chikurachki magmas using the behaviour of fluid-immobile (i.e., Nb, Zr, Ti, Yb) vs. fluid-mobile (i.e., B, Ba, K, Sr, ±Th) elements.

### 5.3.1. Mantle wedge composition

In our model calculations, it is essential to assess chemical composition of the mantle wedge. Generally, the HFSE and HREE are immobile in the fluid and not enriched in the oceanic crust and in sedimentary components of the subduction slab, and the ratios of these elements can be used as reasonable proxy to characterise the nature of the mantle wedge of a particular volcano in the SZ setting (e.g., Pearce, 1983; Pearce and Peate, 1995).

Several trace element characteristics of the studied Chikurachki melt inclusions allow us to conclude that the composition of the mantle wedge is similar or even more depleted than the source of typical N-MORB magmas. They are following: (i) the average N-MORB-normalised concentrations of  $[Nb]_n = 0.55 \pm 0.21$ ,  $[Sm]_n = 1.16 \pm 0.37$ ,  $[Zr]_n = 0.74 \pm 0.30$ ,  $[Ti]_n = 0.74 \pm 0.27$ ,  $[Yb]_n = 0.83 \pm 0.29$  are close to, or lower, than unity; (ii) the element ratios  $Zr/Nb = 35\text{--}49$ ,  $Zr/Yb = 16\text{--}25$ , and  $Nb/Yb = 0.35\text{--}0.7$  are similar to those of N-MORB (i.e.,  $Zr/Nb = 32$ ,  $Zr/Yb = 24$ ,  $Nb/Yb = 0.76$ ; Hofmann, 1988); (iii) the studied melt inclusions plot in the field of MORB in the Pearce and Peate (1995)  $Zr/Yb$  vs.  $Nb/Yb$  discrimination diagram (not shown). Thus, to simplify our calculations, we assume the trace element concentrations of the Kurile arc mantle

wedge is the same as depleted N-MORB mantle of Salters and Stracke (2004) (**Table 1**). We note that using more depleted composition of the mantle wedge will not significantly change the calculation results, but slightly increase a role of the slab-derived components.

### 5.3.2. Subduction slab melt vs. fluid as a volatile transport media

The origin of arc magmas is tightly related to dehydration and/or melting of the slab, where both melt and fluid components are added in different proportions, namely higher proportion of fluid at shallower depths and *vice versa* when the slab penetrates deeper (Hickey and Frey, 1982; Cameron, 1985; Tatsumi et al., 1986; Crawford et al., 1989; Elliott et al., 1997). Experiments on element partitioning confirm that slab-derived H<sub>2</sub>O-rich fluids may cause remarkable enrichment by incompatible elements typical for volcanic arcs worldwide (e.g., Brenan et al., 1995; Ayers et al., 1997; Stalder et al., 1998; Johnson and Plank, 1999). Silica-rich partial melts released from the subducting crust and/or sediment are also potential contributors to the chemical balance (Defant and Drummond, 1990; Drummond and Defant, 1990).

Several studies proposed a concurrent addition of slab-derived melt and fluid components in the arc magmas (e.g., Kay, 1978; Rogers et al., 1985; Stern et al., 1984; Defant et al., 1991; Stern and Kilian, 1996; Class et al., 2000; Yogodzinski et al., 2001). In particular, it has been concluded that hydrous fluids released in the arc front due to amphibole and serpentine dehydration are enriched in boron, LILE, sulphur and chlorine and likely represent a main fluid transport and reaction agent affecting mantle wedge of the Kurile and Kamchatka arcs (e.g., Hochstaedter et al., 1996; Ishikawa and Tera, 1997; Churikova et al., 2001, 2007; Portnyagin et al., 2007). Moreover, Churikova et al. (2007) concluded that S, Cl and B could have been transported by a fluid released from AOC-like subduction slab at shallow depth, whereas Li appeared to be related to Li-F-rich fluids resulting from breakdown of volatile-rich minerals (e.g., amphibole) at much greater depths (~400 km). Furthermore, Churikova et al. (2007) have noted that some Kurile-Kamchatka arc magmas may also exhibit an adakitic signature, whereas

Portnyagin et al. (2015) argued that melting of the mantle wedge beneath Tolbachik was triggered by slab-derived hydrous melts.

We think that in the case of Chikurachki, the significant positive correlations of  $H_2O/K_2O$  vs.  $Cl/K_2O$  and  $F/Zr$  vs.  $Cl/Nb$  (**Fig. 5E,F**),  $Th/Nb$  vs.  $Ba/Nb$  ( $R = 0.649$ ; see **Fig. 9** and discussion below), as well as  $B/Nb$  vs.  $Li/Nb$  and  $Be/Nb$  ( $R = 0.833$  and  $0.796$ , respectively, except for one outlier; **Fig. S2.2** – *Supporting online material*) imply a complex nature of the Kurile arc subduction component composed of both melt and fluid. The observed strong correlations of  $Li/Nb$  and  $Be/Nb$  with  $B/Nb$  and  $F/Zr$  with  $Cl/Nb$  in the Chikurachki melt inclusions suggest that these elements have a common source (i.e., slab component), support a major role of a Cl-rich fluid and imply that B, Cl and F are likely to relate to a single transport process, in contrast to the mechanism of B and Li decoupling suggested by Churikova et al. (2007).

### 5.3.3. Trace element constraints

Thorium and LREE are enriched in the subducted slab, being however less mobile or nearly immobile in the fluid phase, as compared to fluid-mobile elements, such as Li, B, Ba, Pb and Sr. The difference in geochemical behaviour of these elements can be used to elucidate a contribution of slab-derived melts vs. dehydration fluids in the origin of arc magmas (e.g., Kamenetsky et al., 1997; Class et al., 2000; Hochstaedter et al., 2001). Namely, the transport by fluid would leave  $Th/Nb$  ratios virtually unchanged since both elements are immobile but affect the ratios of fluid-mobile to fluid-immobile elements (e.g.,  $Ba/Nb$ ,  $B/Nb$ ), and *vice versa*, if the element transport would have been caused solely by the slab-derived melt. Systematic variations of  $Th/Nb$  vs.  $Ba/Nb$  or  $B/Nb$  ratios would imply the contribution of both slab-derived melt and fluid components.

We modelled the contribution of the anticipated subduction sediment-derived melt and fluid components to the mantle wedge following the approach of Hochstaedter et al. (2001). The

AOC- and SED-derived fluids, as well as a partial melt from SED were considered as possible end-members (**Table 2**). The element partitioning data used in the modelling are from experimentally-determined bulk sediment-melt, sediment-fluid, rutile-melt and rutile-fluid partition coefficients (Brenan et al. 1994, 1995; Stalder et al., 1998; Johnson and Plank, 1999; Foley et al., 2000). The sediment subducted at the Kurile trench (its composition is taken from Plank and Langmuir, 1998) is thought to contain up to 5% Ti-rich residual phase (i.e. rutile), which has the potential to fractionate Nb from Th (Class et al., 2000), whereas 1% rutile was assigned for AOC (Hochstaedter et al., 2001). Equilibrium batch melting and Rayleigh fractionation equations (Shaw, 1970) were used to calculate the compositions of partial melt and fluid in equilibrium with the subduction sediment, respectively. We assumed 5% partial melting of the Kurile sediment to produce a sediment melt mixing end-member, and 2% fluid removal from both the Kurile sediment and AOC to account for a possible sediment fluid end-member (**Fig. 10**). We note also that because Ba, Nb and Th are chemical elements of very similar incompatibility to minerals composing magma source rocks, as follows from their solid-melt partition coefficients, they show little or no fractionation during partial melting and/or magma fractionation. The calculated Ba/Nb and Th/Nb ratios of the multi-component magma source region are thus relevant for the Th/Nb and Ba/Nb of the produced partial melts, and can be used to assess the origin of the Chikurachki parental magmas.

In the model presented in **Fig. 10**, the Th/Nb vs. Ba/Nb variations observed in the Chikurachki melt inclusions (presented alongside whole-rock chemical compositions of the Kurile arc mafic lavas) can be explained by partial melting of a magma source composed of the following three components (**Table 2**):

1. A mantle wedge (*MW*) component with trace element concentrations of depleted mantle after Salters and Stracke (2004).

2. A composite dehydration fluid, which is enriched in Ba relative to Th or Nb and presents a 9:1 mixture of the fluid released from the altered oceanic crust (*AOCFluid*, the composition is taken from Hochstaedter et al., 2001) and a fluid produced by dehydration of a subduction sediment (*SedFluid*), whose trace element composition was calculated from that of the Kurile sediment after Plank and Langmuir (1998) (for more detail see explanatory notes in **Table 2**). The 9:1 proportion of *AOCFluid* to *SedFluid* was selected arbitrary, given a subordinate amount of the sediment relative to a total mass of the slab, taking also into account that sediment is more volatile-rich, as compared to its basaltic and ultramafic counterparts. We note however that variations of *AOCFluid* to *SedFluid* ratio do not affect significantly the calculation results.
3. A sediment-derived melt (*SedMelt*) with slightly elevated Th/Nb ratios and significantly lower Ba/Nb ratios, as compared *AOCFluid* and *SedFluid* mixing end-members, whose trace element composition was also calculated from the composition of the Kurile sediment (see also explanatory notes in **Table 2**).

As it follows from our calculations, the composition of subduction component (*SUBD*) is variable. It ranges from the composition of the pure *SedMelt* component (representing a lower limit of the range) to that of *SedMelt* mixed with up to 60% composite slab-derived fluid presenting, in turn, a 9:1-mixture of *AOCFluid* and *SedFluid*, crudely corresponding to the AOC to SED proportion in the dehydrating model slab (Tonarini et al., 2011) and assuming also equal amount of fluid released from both lithologies. The ~1–8% admixture of *SUBD* to *MW* appeared to satisfactorily explain the entire compositional range of the Chikurachki melt inclusions, and from ~0.4% to ~10% of *SUBD* to be mixed with *MW* to account for the range of the Kurile arc magmas (**Fig. 10**). In conclusion, it is worth emphasising that the subduction component contributed to the origin of the Tolbachik magmas (Portnyagin et al., 2015) appeared to be very similar to *SUBD* inferred during the present study (**Fig. 9; Table 2**).

#### 5.3.4. Implications for chemical composition and amount of slab-derived fluid

Here we evaluate a possible contribution of slab-derived fluid to the origin of volatile components and boron in the studied magmas, assessing the observed B-isotope range. The light lithophile element boron is a powerful proxy to trace fluid transport of chemical elements from subducting slab to the mantle wedge because of strongly contrasting B concentrations and  $\delta^{11}\text{B}$  values in these geochemical reservoirs. A significant fraction of initial boron is contained in the uppermost few km of the subducting crust and sediments, being then progressively released as subduction proceeds (e.g., Leeman and Sisson, 1996). Systematically higher  $\delta^{11}\text{B}$  values in the volcanic arc magmas, as compared to the exhumed subduction-related metamorphic rocks, suggest that dehydration reactions cause significant decrease of  $\delta^{11}\text{B}$  values of subducted crustal material, implying strong fractionation of  $^{10}\text{B}$  and  $^{11}\text{B}$  isotopes as boron partitions in the fluid (e.g., Peacock and Hervig, 1999). This conclusion was supported by later experimental study of Wunder et al. (2005), who have demonstrated that a wide range of  $\delta^{11}\text{B}$  in the arc magmas may result from continuous breakdown of micas, if they are present in the down-dragged slab. Finally, a serpentinitised mantle wedge at the slab-mantle interface represents another major inventory of boron, being also a source of variable but generally positive  $\delta^{11}\text{B}$  values (e.g., Benton et al., 2001; Straub and Layne, 2002; Savov et al., 2005, 2007; Scambelluri and Tonarini, 2012).

Boron mobilised from the slab was shown to be systematically enriched in  $^{11}\text{B}$  than the parental AOC at 400–500°C, resulting in +5 to +10‰ enrichment of the fluid relative to the solid (Ishikawa and Nakamura, 1992; Leeman and Sisson, 1996; Rose et al. 2001; Rosner et al. 2003; Leeman et al., 2004; Le Voyer et al., 2008). Since the composition of slab-derived fluid may vary strongly, depending on the composition of subducting crust and the degree of its devolatilisation, a wide  $\delta^{11}\text{B}$  range from –7‰ up to +15‰ (assuming +10‰ enrichment of the

fluid phase relative to the upper end of the  $\delta^{11}\text{B}$  range given for AOC by Smith et al., 1997) is very probable (Rose et al. 2001; Leeman et al., 2004; Gurenko et al., 2005b; Le Voyer et al., 2008; Tonarini et al., 2011, Scambelluri and Tonarini, 2012).

The effects of possible interaction of slab-derived fluids with depleted MORB-like mantle source are illustrated in diagrams in coordinates of  $\delta^{11}\text{B}$  vs. B/Nb and Ba/Nb ratios (**Fig. 11**). Given the  $^{11}\text{B}/^{10}\text{B}$ , B/Nb and Ba/Nb do not change significantly during partial melting or magma fractionation, these ratios, being calculated for the magma source, can be directly transferred to the resulting partial melts. The following mixing end-members were defined (**Table 1**):

1. Mantle wedge (*MW*) was chosen to contain 1.2  $\mu\text{g/g}$  Ba, 0.21  $\mu\text{g/g}$  Nb, 0.06  $\mu\text{g/g}$  B and  $\delta^{11}\text{B} = -4\%$  (Chaussidon and Marty, 1995; Salters and Stracke, 2004).
2. Subducting slab fluid (*SSF*) varies significantly in chemical composition, depending on the particular chemical and lithological composition of the slab and the dehydration depth. In particular, Tonarini et al. (2011) have demonstrated that slab-derived fluids released at the depth range from 30 to 120 km may contain from  $\sim 10$  and  $\sim 200$   $\mu\text{g/g}$  B and vary between  $-10$  to higher than  $+20\%$   $\delta^{11}\text{B}$ . Three contrasting fluid compositions (*SSF1* through *SSF3*) were compiled based on the data of Hochstaedter et al. (2001), Rose et al. (2001), Straub and Layne (2002), Rosner et al. (2003), Leeman et al. (2004), Gurenko et al. (2005b), Le Voyer et al. (2008), Tonarini et al. (2011), Scambelluri and Tonarini (2012), using also the compositions of *SED* and *AOC*-derived fluids calculated during the present study (*SSF1*: trace elements concentrations are those as of *SEDFluid*, containing 300  $\mu\text{g/g}$  B and  $\delta^{11}\text{B} = +15\%$ ; *SSF2* is the 9:1 mixture of *AOCFluid* and *SedFluid* as calculated above, containing 100  $\mu\text{g/g}$  B and  $\delta^{11}\text{B} = +3\%$ ; and *SSF3* is equal to *AOCFluid* with 50  $\mu\text{g/g}$  B and  $\delta^{11}\text{B} = -8\%$ ). The decrease of B and Ba contents and

$\delta^{11}\text{B}$  values in the fluid components qualitatively reflects the increase of slab temperature and degree of its devolatilisation.

As it follows from the calculations, the slab-derived fluid may represent a major agent affecting the abundances of volatile and fluid-mobile elements, and B isotopes in the melt inclusions. About 1–10% of SSF2 and SSF3 have to be added to MW in order to explain the  $\sim 9\%$  range of  $\delta^{11}\text{B}$  values of the studied melt inclusions. We note that this estimation is very similar to that inferred from the trace element modelling (1 to 8% addition of SUBD; **Fig. 9**). It is also worth emphasising that the selected compositions of SSF2 and SSF3 are very close to the model fluid compositions released from the South Sandwich Island arc uppermost slab at 90–120 km depth (170–190  $\mu\text{g/g}$  B and  $\delta^{11}\text{B} = -0.7$  to  $-3.9\%$ ; Tonarini et al., 2011), but having less boron and more negative  $\delta^{11}\text{B}$  values. This discrepancy can be explained by somewhat stronger dehydration of the residual Kurile slab because of the greater depth of the Wadati-Benioff zone beneath Chikurachki ( $\sim 130$ – $140$  km; e.g., Ishikawa and Tera, 1997; Ishikawa et al., 2001).

#### *5.4. Magma ascent, degassing history and eruption style*

It is well known that conduit geometry, magma viscosity, degassing history, changing conditions of magma ascent and fragmentation, and even the dynamics of syn-eruptive microlite growth may strongly affect the eruption rate and style of a volcano (e.g., Fischer and Schmincke, 1984; Carey and Sparks, 1986; Houghton et al., 2004; Szramek et al., 2006, 2010; Houghton and Gonnermann, 2008). Previously, Gurenko et al. (2005a) suggested intensive pre-eruptive degassing of  $\text{H}_2\text{O}$  and S from the magma at the stage of phenocryst crystallisation, pointing on the strongly varying to very low volatile contents in the melt inclusions trapped by phenocrysts of the same or very close chemical composition. The authors suggested a scenario by which local volumes of magma could be partially degassed before being trapped by growing crystals and

employed a mechanism of vertical cycling of magma within one or between several connected reservoirs, being akin to the magma drainback phenomenon described in particular for Kilauea Volcano, Hawaii (e.g., Wallace and Anderson, 1998). Our present data (i.e., numerous correlations of volatile component concentrations and S isotope compositions preserved in the studied melt inclusions with incompatible trace elements and between each other, inability of magma degassing and crustal contamination to account for these variations coupled with the suggested here two-component mixing scenario to explain  $\delta^{34}\text{S}$  vs. S concentrations) could be interpreted in support of the mechanism previously suggested by Gurenko et al. (2005a) to control the variations of H<sub>2</sub>O, S, Cl and F in the erupting magmas.

The required insignificant correction of melt inclusion compositions for post-entrapment crystallisation of host mineral (i.e., less than 7.8 wt.% Ol, 0.8 wt.% Opx, and no correction for Pl) also suggests that the eruption and quenching of magma likely occurred shortly after the inclusions have been entrapped by growing phenocrysts. As previously discussed (e.g., Roggensack et al., 1997; Gurenko et al., 2005a; Shinohara, 2008; Ruth et al., 2016), eventual eruptions may be non-explosive, if magma degassing is effective at shallow crustal depth, whereas rapid ascent of a volatile-rich magma may result in strong volatile release and highly explosive character of eruptions. In this context, the range of volatile contents recorded by melt inclusions and their relationships with other chemical components may indicate whether the erupted magma underwent a prolonged stage of relatively passive degassing or degassing was essentially a single-stage process upon the eruption. It is worth emphasising that most of the historic eruptions from Chikurachki have been of the non-explosive character, possibly linked to the episodes where volatile components were able to dissipate smoothly. Explosive eruptions, in contrast, are thought to occur when access to shallow levels is blocked or when the flux of magma from depth is too rapid for shallow degassing to prevent a catastrophic decompression and explosive fragmentation of the magma. We thus think that rapid ascent of the volatile-rich

basaltic magmas from ~16-km initial depth (inferred from our CO<sub>2</sub>-H<sub>2</sub>O data; **Fig. 4B**) was a primary reason caused the Plinian character of the Chikurachki eruptions.

## 6. Summary and conclusions

The present study focuses on the unravelling the origin and behaviour of volatile components resulting in highly explosive (Plinian) basaltic eruptions of the Chikurachki volcano.

The following main conclusions were derived:

1. The obtained concentrations of volatile components in the studied melt inclusions agree well with our previously reported data. After correction for post-entrapment crystallisation of the host mineral, the concentrations of CO<sub>2</sub> and H<sub>2</sub>O reveal minimum pressure of 33 to 276 MPa (or ~9.5 to ~1 km depth), corresponding to the last equilibrium between melt and vapour phases upon magma eruption and quenching. About 5–69% of the original CO<sub>2</sub> and 0.1–3.2% of H<sub>2</sub>O were lost to the shrinkage bubble of melt inclusions after their entrapment during phenocryst crystallisation. After correction, the obtained 47–1580 μg/g CO<sub>2</sub>, 0.4–4.2 wt.% H<sub>2</sub>O, 399–633 μg/g F, 619–3402 μg/g S and 805–1240 μg/g Cl ranges imply strong pressure release from 456 to 35 MPa, corresponding to ~15.7–1.2-km-depth range of magma ascent.
2. Significant negative correlations of S with incompatible trace elements, no clear relationships of S with H<sub>2</sub>O, CO<sub>2</sub> and Cl but strong positive correlations of S/K<sub>2</sub>O with H<sub>2</sub>O/K<sub>2</sub>O, Cl/K<sub>2</sub>O and F/K<sub>2</sub>O suggest that later magma degassing and/or contamination by crustal rocks were not the governing processes affecting the amounts of S and probably of the other volatile components preserved in inclusions
3. Sulphur isotopes in the studied inclusion and groundmass glasses vary strongly, ranging from -1.6 to +12.3‰ of δ<sup>34</sup>S, decrease with decreasing S concentrations and show significant

positive correlations with H<sub>2</sub>O, CO<sub>2</sub> and S, as well as negative correlations with light lithophile (Li, Be and B), and with a number of incompatible trace elements. Our modelling has demonstrated that either open- or close-system magma degassing cannot account for the observed ~14‰ range of  $\delta^{34}\text{S}$  variations in the Chikurachki melt inclusions. The linear character of the 1/S vs.  $\delta^{34}\text{S}$  relationship suggests mixing of two components possibly represented by melts having similar major and trace element compositions but strongly contrasting volatile contents and S isotopes.

4. The  $\delta^{11}\text{B}$  values of the studied melt inclusions range from -7.0 to +2.4‰ within the interval of B concentrations of 13–23  $\mu\text{g/g}$ . Similarly, the relationships of  $\delta^{11}\text{B}$  with B/K<sub>2</sub>O and B/Nb, as well as of Cl/K<sub>2</sub>O with B/K<sub>2</sub>O (and equally, Cl/Nb with B/Nb) cannot be accounted by magma contamination at shallow crustal depths.
5. The observed relationships of S/K<sub>2</sub>O with K<sub>2</sub>O-normalised concentrations of H<sub>2</sub>O, Cl and F, as well as the H<sub>2</sub>O/K<sub>2</sub>O vs. Cl/K<sub>2</sub>O and F/Zr vs. Cl/Nb correlations strongly suggest that variations in H<sub>2</sub>O, S, Cl and F contents were not primarily affected by degassing or contamination, but likely originated due to the mixing of magmas, which in turn reflect the composition of source components. We contend that subduction slab-derived melt and/or fluid were major agents in controlling chemical and isotopic composition of the mantle wedge and, consequently, of the magmas originated from this hybrid source.
6. The behaviour of fluid-mobile vs. fluid-immobile incompatible trace elements points towards strongly varying chemical composition of the subduction component. It ranges from the composition of the pure subducted sediment-derived melt (a lower limit of the range) to that of the sediment melt mixed with up to 60% composite slab-derived fluid, representing in turn a 9:1-mixture of fluid components released from both altered oceanic crust and the sediment of the slab. The ~1–8% admixture of the inferred subduction component to the mantle wedge is required to account for the whole compositional range of the studied Chikurachki magmas,

and addition from ~0.4% to ~10% of the subduction component accounts for the broad compositional range of the Kurile arc magmas.

7. We conclude that rapid ascent of the volatile-rich basaltic magmas from ~16-km initial depth without their prolonged stagnation and crystallisation in shallower depth magma reservoir but accompanied by near-surface bubble nucleation and growth followed by magma fragmentation was a primary reason caused Plinian character of the Chikurachki eruptions.

### **Acknowledgements**

We thank Peter Landry for his help and ensuring stable working conditions of the NENIMF CAMECA IMS 1280 instrument, Charlie Mandeville and Alberto Saal for fruitful discussions, Anastassia Borisova for review and comments of the early manuscript version. The insightful reviews, useful comments and suggestions, as well as a thorough correction of English by John Caulfield and two anonymous referees are very much appreciated. This work was benefited from the NENIMF financial support of AAG during his training as a SIMS research specialist, the NSF grant EAR 0911093 to AAG, and partially from the Russian Science Foundation grant #16-17-10145 to VSK and MEZ. This is CRPG contribution #24??

**References**

- Alt, J.C., Shanks III, W.C., Jackson, M.C., 1993. Cycling of sulfur in subduction zones: The geochemistry of sulfur in the Mariana Island Arc and back-arc trough. *Earth and Planetary Science Letters* 119, 477–494.
- Anderson, A.T., 1974. Chlorine, sulfur and water in magmas and oceans. *Bulletin of Geological Society of America* 85, 1485–1492.
- Arculus, R.J., Powell, R., 1986. Source component mixing in the regions of arc magma generation. *Journal of Geophysical Research* 91, 5913–5926.
- Aster, E.M., Wallace, P.J., Moore, L.R., Watkins, J., Gazel, E., Bodnar, R.J., 2016. Reconstructing CO<sub>2</sub> concentrations in basaltic melt inclusions using Raman analysis of vapor bubbles. *Journal of Volcanology and Geothermal Research* 323, 148–162.
- Ayers J.C., Dittmer, S.K., Layne G., 1997. Partitioning of elements between peridotite and H<sub>2</sub>O at 2.0–3.0 GPa and 900–1100°C, and application to models of subduction zone processes. *Earth and Planetary Science Letters* 150, 381–398.
- Bailey, J.C., Frolova, T.I., Burikova, I.A., 1987a. Mineralogy, geochemistry and petrogenesis of Kurile Island Arc basalts. *Contribution to Mineralogy and Petrology* 102, 265–280.
- Bailey, J.C., Larsen, O., Frolova, T.I., 1987b. Strontium isotope variations in lower Tertiary-Quaternary volcanic rocks from the Kurile Island Arc. *Contribution to Mineralogy and Petrology* 95, 155–165.
- Belousov, A.B., Belousova, M.G., Grushin, S.Yu., Krestov, P.B., 2003. Historic eruptions of the Chikurachki volcano (Paramushir, Kurile Islands). *Volcanology and Seismology* 3, 15-34 (in Russian).
- Belousov, A., Belousova, M., Edwards, B., Volynets, A., Melnikov, D., 2015. Overview of the precursors and dynamics of the 2012–13 basaltic fissure eruption of Tolbachik Volcano, Kamchatka, Russia. *Journal of Volcanology and Geothermal Research* 307, 22-37.

- Belousov, A., Belousova, M., Kozlov, D. 2017. Distribution of tephra deposit and reconstruction of parameters of the 1973 explosive eruption of Tyatya volcano, Kunashir Island, Kuriles. *Volcanology and Seismology*, in print (in Russian).
- Benton, L.D., Ryan, J.G., Tera, F., 2001. Boron isotope systematics of slab fluids as inferred from a serpentine seamount, Mariana forearc, *Earth and Planetary Science Letters* 187, 273–282.
- Bernard, A., Demaiffe, D., Mattielli, N., Punanongbayan, R.S., 1991. Anhydrite-bearing pumices from Mount Pinatubo: further evidence for the existence of sulfur-rich silicic magmas. *Nature* 354, 139–140.
- Bindeman, I.N., Ponomareva, V.V., Bailey, J.C., Valley, J.W., 2004. Volcanic arc of Kamchatka: a province with high- $\delta^{18}\text{O}$  magma sources and large-scale  $^{18}\text{O}/^{16}\text{O}$  depletion of the upper crust. *Geochimica et Cosmochimica Acta* 68, 841–865.
- Brasseur, G., Granier, C., 1992. Mount Pinatubo aerosols, chlorofluorocarbons and ozone depletion. *Science* 257, 1239–1242.
- Brenan, J.M., Shaw, H.F., Phinney, D.L., Ryerson, F.J., 1994. Rutile-aqueous fluid partitioning of Nb, Ta, Hf, Zr, U and Th: implications for high field strength element depletions in island-arc basalts. *Earth and Planetary Science Letters* 128, 327–339.
- Brenan, J.M., Shaw, H.F., Ryerson, F.J., Phinney, D.L., 1995. Mineral-aqueous fluid partitioning of trace elements at 900°C and 2.0 GPa: constraints on the trace element chemistry of mantle and deep crustal fluids. *Geochimica et Cosmochimica Acta* 59, 3331–3350.
- Burgisser, A., Scaillet, B., 2007. Redox evolution of a degassing magma rising to the surface. *Nature* 445, 194–197.
- Cabral, R.A., Jackson, M.G., Koga, K.T., Rose-Koga, E.F., Hauri, E.H., Whitehouse, M.J., Price, A.A., Day, J.M.D., Shimizu, N., Kelley, K.A., 2014. Volatile cycling of  $\text{H}_2\text{O}$ ,  $\text{CO}_2$ , F, and Cl

- in the HIMU mantle: A new window provided by melt inclusions from oceanic hot spot lavas at Mangaia, Cook Islands. *Geochemistry Geophysics Geosystems* 15, 4445–4467.
- Cameron, W.E., 1985. Petrology and origin of primitive lavas from the Troodos ophiolite, Cyprus. *Contribution to Mineralogy and Petrology* 89, 239–255.
- Canil, D., Fellows, S.A., 2017. Sulphide–sulphate stability and melting in subducted sediment and its role in arc mantle redox and chalcophile cycling in space and time. *Earth and Planetary Science Letters* 470, 73–86.
- Carey, S., Sparks, R.S.J., 1986. Quantitative models of the fallout and dispersal of tephra from volcanic eruption columns. *Bulletin of Volcanology* 48, 109–125.
- Carroll, M.R., Rutherford, M.J., 1985. Sulfide and sulfate saturation in hydrous silicate melts. *Journal of Geophysical Research* 90, C601–C612.
- Carroll, M.R., Rutherford, M.J., 1987. The stability of igneous anhydrite: experimental results and implications for sulfur behavior in the 1982 El Chichon trachyandesite and other evolved magmas. *Journal of Petrology* 28, 781–801.
- Carroll, M.R., Rutherford, M.J., 1988. Sulfur speciation in hydrous experimental glasses of varying oxidation state: results from measured wavelength shifts of sulfur X-ray. *American Mineralogist* 73, 845–849.
- Carroll, M.R., Webster, J.D., 1994. Solubilities of sulfur, noble gases, nitrogen, chlorine, and fluorine in magmas, in: Carroll, M.R., Holloway, J.R. (Eds.), *Volatiles in Magmas, Reviews in Mineralogy* 30, Mineralogical Society of America, Washington DC, pp 231–279.
- Caulfield, J.T., Cronin, S.J., Turner, S.P., Cooper, L.B., 2011. Mafic Plinian volcanism and ignimbrite emplacement at Tofua volcano, Tonga. *Bulletin of Volcanology* 73, 1259–1277.
- Chaussidon, M., Jambon, A., 1994. Boron content and isotopic composition of oceanic basalts: geochemical and cosmochemical implications. *Earth and Planetary Science Letters* 121, 277–291.

- Chaussidon, M., Marty, B., 1995. Primitive boron isotope composition of the mantle. *Science*, 269, 383–386.
- Chaussidon, M., Albarède, F., Sheppard, S.M.F., 1987. Sulphur isotope heterogeneity in the mantle from ion microprobe measurements of sulphide inclusions in diamonds. *Nature* 330, 242–244.
- Churikova T., Dorendorf F., Wörner G., 2001. Sources and fluids in the mantle wedge below Kamchatka, evidence from across-arc geochemical variation. *Journal of Petrology* 42, 1567–1593.
- Churikova, T., Wörner, G., Mironov, N., Kronz, A., 2007. Volatile (S, Cl and F) and fluid mobile trace element compositions in melt inclusions: implications for variable fluid sources across the Kamchatka arc. *Contributions to Mineralogy and Petrology* 154, 217–239.
- Class, C., Miller, D.M., Goldstein, S.L., Langmuir, C.H., 2000. Distinguishing melt and fluid subduction components in Umnak Volcanics, Aleutian Arc. *Geochemistry Geophysics Geosystems* 1, 1999GC000010.
- Connolly, J.W.D., Haughton, D.R., 1972. The valence of sulfur in glass of basaltic composition formed under low oxidation potential. *American Mineralogist* 57, 1515–1517.
- Crawford, A.J., Falloon, T.J., Green, D.H., 1989. Classification, petrogenesis and tectonic setting of boninites, in: Crawford, A.J. (Ed.), *Boninites*. Unwin Hyman, London, pp. 1–49.
- Dalou, C., Koga, K.T., Le Voyer, M., Shimizu, N., 2014. Contrasting partition behaviour of F and Cl during hydrous mantle melting: implications for Cl/F signature in arc magmas. *Progress in Earth and Planetary Science* 1, 26.
- Davidson, J.P., 1987. Crustal contamination versus subduction zone enrichment: examples from the Lesser Antilles and implications for mantle source composition of island arc volcanic rocks. *Geochimica et Cosmochimica Acta* 51, 2185–2198.

- Defant, M.J., Drummond, M.S., 1990. Derivation of some modern arc magmas by melting of young subducted lithosphere. *Nature* 347, 662–665.
- Defant, M.J., Richerson, P.M., De Boer, J.Z., Stewart, R.H., Maury, R.C., Bellon, H., Drummond, M.S., Feigenson, M.D., Jackson, T.E., 1991. Dacite genesis via both slab melting and differentiation: Petrogenesis of La Yeguada volcanic complex, Panama. *Journal of Petrology* 32, 1101–1142.
- Ding, T., Valkiers, S., Kipphardt, H., De Bièvre, P., Taylor, P.D.P., Gonfiantini, R., Krouse, R., 2001. Calibrated sulfur isotope abundance ratios of three IAEA sulfur isotope reference materials and V-CDT with a reassessment of the atomic weight of sulfur. *Geochimica et Cosmochimica Acta* 65: 2433–2437.
- Dorendorf, F., Wiechert, U., Wörner, G., 2000. Hydrated sub-arc mantle: a source for the Kluchevskoy volcano, Kamchatka/Russia. *Earth and Planetary Science Letters* 175, 69–86.
- Drummond, M.S., Defant, M.J., 1990. A model for trondhjemite-tonalite-dacite genesis and crustal growth via slab melting: Archaean to modern comparisons. *Journal of Geophysical Research* 95, 503–521.
- Elliott, T., Plank, T., Zindler, A., White, W., Bourdon, B., 1997. Element transport from slab to volcanic front at the Mariana arc. *Journal of Geophysical Research* 102, 14991–15019.
- Esposito, R., Bodnar, R.J., Danyushevsky, L.V., De Vivo, B., Fedele, L., Hunter, J., Lima, A., Shimizu, N., 2011. Volatile evolution of magma associated with the Solchiaro eruption in the Phlegrean Volcanic District (Italy). *Journal of Petrology* 52, 2431–2460.
- Fedotov, S.A., Markhinin, Ye.K., 1983. *The Great Tolbachik Fissure Eruption*. Cambridge Univ. Press, New York.
- Fiege, A., Holtz, F., Behrens, H., Mandeville, C.W., Shimizu, N., Crede, L.S., Göttlicher, J., 2015. Experimental investigation of the S and S-isotope distribution between H<sub>2</sub>O-S ± Cl fluids and basaltic melts during decompression. *Chemical Geology* 393–394, 36–54.

- Fisher, R.V., Schmincke, H.-U., 1984. Pyroclastic rocks. Springer-Verlag, Berlin, Heidelberg, New York, Tokyo.
- Foley, S.F., Barth, M.G., Jenner, G.A., 2000. Rutile/melt partition coefficients for trace elements and an assessment of the influence of rutile on the trace element characteristics of subduction zone magmas. *Geochimica et Cosmochimica Acta* 64, 933–938.
- Francis, P., 1993. *Volcanoes: a planetary perspective*. Clarendon Press, Oxford.
- Gill, J.B., 1981. *Orogenic andesites and plate tectonics*. Springer-Verlag, Berlin-Heidelberg.
- Gill, J., Torssander, P., Lapierre, H., Taylor, R., Kaiho, K., Koyama, M., Kusakabe, M., Aitchison, J., Cisowski, S., Dadey, K., Fujioka, K., Klaus, A., Lovell, M., Marsaglia, K., Pezard, P., Taylor, B., Tazaki, K., 1990. Explosive deep-water basalt in the Sumisu Backarc Rift. *Science* 248, 1214–1217.
- Gorshkov, G.S., 1967. *Volcanism of the Kurile Island Arc*. Nauka, Moscow (in Russian).
- Gurenko, A.A., Chaussidon, M., 1997. Boron concentrations and isotopic composition in the Icelandic mantle: evidence from glass inclusions in olivine. *Chemical Geology* 135, 21–34.
- Gurenko, A.A., Kamenetsky, V.S., 2011. Boron isotopic composition of olivine-hosted melt inclusions from Gorgona komatiites, Colombia: New evidence supporting wet komatiite origin. *Earth and Planetary Science Letters* 312, 201–212.
- Gurenko, A.A., Schmincke, H.-U., 2000. S concentrations and its speciation in Miocene basaltic magmas north and south of Gran Canaria (Canary Islands): Constraints from glass inclusions in olivine and clinopyroxene. *Geochimica et Cosmochimica Acta* 64, 2321–2337.
- Gurenko, A.A., Chaussidon, M., Schmincke, H.-U., 2001. Magma ascent and contamination beneath one intraplate volcano: Evidence from S and O isotopes in glass inclusions and their host clinopyroxenes from Miocene basaltic hyaloclastites southwest of Gran Canaria (Canary Islands). *Geochimica et Cosmochimica Acta* 65, 4359–4374.

- Gurenko, A.A., Belousov, A.B., Trumbull, R.B., Sobolev, A.V., 2005a. Explosive basaltic volcanism of the Chikurachki Volcano (Kurile arc, Russia): Insights on pre-eruptive magmatic conditions and volatile budget revealed from phenocryst-hosted melt inclusions and matrix glasses. *Journal of Volcanology and Geothermal Research* 147, 203–232.
- Gurenko, A.A., Trumbull, R.B., Thomas, R., Lindsay J.M., 2005b. A melt inclusion record of volatiles, trace elements and Li-B isotope variations in a single magma system from the Plat Pays Volcanic Complex, Dominica, Lesser Antilles. *Journal of Petrology* 46, 2495–2526.
- Hartley, M.E., Maclennan, J., Edmonds, M., Thordarson, T., 2014. Reconstructing the deep CO<sub>2</sub> degassing behaviour of large basaltic fissure eruptions. *Earth and Planetary Science Letters* 393, 120–131.
- Hasegawa, T., Nakagawa, M., Yoshimoto, M., Ishizuka, Y., Hirose, W., Seki, S. I., Ponomareva, V., Alexander, R., 2011. Tephrostratigraphy and petrological study of Chikurachki and Fuss volcanoes, western Paramushir Island, northern Kurile Islands: Evaluation of Holocene eruptive activity and temporal change of magma system. *Quaternary International* 246, 278–297.
- Haughton, D.R., Roeder, P.L., Skinner, B.J., 1974. Solubility of sulfur in mafic magmas. *Economic Geology* 69, 541–567.
- Hawkesworth, C.J., Gallagher K., Hergt J. M., and McDermott F. (1993) Mantle and slab contributions in arc magmas. *Annual Review of Earth and Planetary Sciences* 21, 175–204.
- Hickey, R.L., Frey, F.A., 1982. Geochemical characteristics of boninite series volcanics: Implications for their source. *Geochimica et Cosmochimica Acta* 46, 2099–2115.
- Hickey, R., Frey, F.A., Gerlach, D.C., Lopez-Escobar, L., 1986. Multiple sources for basaltic arc rocks from the southern volcanic zone of the Andes (34°–41°S): Trace element and isotopic evidence for contributions from subducted oceanic crust, mantle and continental crust. *Journal of Geophysical Research* 91, 5963–5983.

- Hochstaedter, A.G., Kepezhinskas, P., Defant, M., 1996. Insights into volcanic arc mantle wedge from magnesian lavas from the Kamchatka arc. *Journal of Geophysical Research* 101, 697–712.
- Hochstaedter, A., Gill, J., Peters, R., Broughton, P., Holden, P., Taylor, B., 2001. Across-arc geochemical trends in the Izu-Bonin arc: Contributions from the subducting slab. *Geochemistry Geophysics Geosystems* 2, 2000GC000105.
- Hofmann, A.W., 1988. Chemical differentiation of the Earth: the relationship between mantle, continental crust and oceanic crust. *Earth and Planetary Science Letters* 90, 297–314.
- Holloway, J.R., Blank, J.G., 1994. Application of experimental results to C-O-H species in natural melts, in: Carroll, M.R., Holloway, J.R. (Eds.), *Volatiles in Magmas. Reviews in Mineralogy* 30, Mineralogical Society of America, Washington, D.C., pp. 187–230.
- Houghton, B.F., Gonnermann, H.M., 2008. Basaltic explosive volcanism: Constraints from deposits and models. *Chemie der Erde* 68, 117–140.
- Houghton, B.F., Wilson, C.J.N., Del Carlo, P., Coltelli, M., Sable, J.E., Carey, R., 2004. The influence of conduit processes on changes in style of basaltic Plinian eruptions: Tarawera 1886 and Etna 122 BC. *Journal of Volcanology and Geothermal Research* 137, 1–14.
- Ikeda, Y., 1998. Geochemistry of Miocene back-arc basin basalts from northeast Hokkaido, Japan. *The Journal of the Geological Society of Japan* 104, 99–106.
- Ikeda, Y., Stern, R.J., Kagami, H., Sun, C.-H., 2000. Pb, Nd, and Sr isotopic constraints on the origin of Miocene basaltic rocks from northeast Hokkaido, Japan: implications for opening of the Kurile back-arc basin. *Island Arc* 9, 161–172.
- Ishikawa, T., Nakamura, E., 1992. Boron isotope geochemistry of the oceanic crust from DSDP/ODP hole 504B. *Geochimica et Cosmochimica Acta* 56, 1633–1639.

- Ishikawa, T., Tera, F., 1997. Source, composition and distribution of the fluid in the Kurile mantle wedge: constraints from across-arc variations of B/Nb and B isotopes. *Earth and Planetary Science Letters* 152, 123–138.
- Ishikawa, T., Tera, F., Nakazawa, T., 2001. Boron isotope and trace element systematics of the three volcanic zones in the Kamchatka arc. *Geochimica et Cosmochimica Acta* 65, 4523–4537.
- Jambon, A. 1994. Earth degassing and large-scale geochemical cycling of volatile elements, in: Carroll, M.R., Holloway, J.R. (Eds.), *Volatiles in Magmas. Reviews in Mineralogy* 30, Mineralogical Society of America, Washington, D.C., pp 479–517.
- Jambon, A., Déruelle, B., Dreibus, G., Pineau, F., 1995. Chlorine and bromine abundance in MORB: the contrasting behaviour of the Mid-Atlantic Ridge and East Pacific Rise and implications for chlorine geodynamic cycle. *Chemical Geology* 126, 101–117.
- Johnson, M.C., Plank, T., 1999. Dehydration and melting experiments constrain the fate of subducted sediments. *Geochemistry Geophysics Geosystems* 1, 1999GC000014.
- Jugo, P.J., Luth, R.W., Richards, J.P., 2005a. An experimental study of the sulfur content in basaltic melts saturated with immiscible sulfide or sulfate Liquids at 1300°C and 1.0 GPa. *Journal of Petrology* 46, 783–798.
- Jugo, P.J., Luth, R.W., Richards, J.P., 2005. Experimental data on the speciation of sulfur as a function of oxygen fugacity in basaltic melts. *Geochimica et Cosmochimica Acta* 69, 497–503.
- Jugo, P.J., Wilke, M., Botcharnikov, R.E., 2010. Sulfur K-edge XANES analysis of natural and synthetic basaltic glasses: Implications for S speciation and S content as function of oxygen fugacity. *Geochimica et Cosmochimica Acta* 74, 5926–5938.
- Kamenetsky, V.S., Crawford, A.J., Eggins, S., Mühe, R., 1997. Phenocryst and melt inclusion chemistry of near-axis seamounts, Valu Fa Ridge, Lau basin: insight into mantle wedge

- melting and the addition of subduction components. *Earth and Planetary Science Letters* 151, 205–223.
- Katsura, T., Nagashima, S., 1974. Solubility of sulfur in some magmas at 1 atmosphere. *Geochimica et Cosmochimica Acta* 38, 517–531.
- Kay, R.W., 1978. Aleutian magnesian andesites: melts from subducted Pacific ocean crust. *Journal of Volcanology and Geothermal Research* 4, 117–132.
- Kendrick, M.A., Arculus, R., Burnard, P., Honda, M., 2013. Quantifying brine assimilation by submarine magmas: Examples from the Galápagos Spreading Centre and Lau Basin. *Geochimica et Cosmochimica Acta* 123, 150–165.
- Kent, A.J.R., Norman, M.D., Hutcheon, I.D., Stolper, E.M., 1999. Assimilation of seawater-derived components in an oceanic volcano: evidence from matrix glasses and glass inclusions from Loihi seamount, Hawaii. *Chemical Geology* 156, 299–319.
- Kepezhinskas, P., McDermott, F., Defant, M.J., Hochstaedter, A., Drummond, M.S., Hawkesworth, C., Koloskov, A., Maury, R.C., Bellon, H., 1997. Trace element and Sr-Nd-Pb isotopic constraints on a three-component model of Kamchatka Arc petrogenesis. *Geochimica et Cosmochimica Acta* 61, 577–600.
- Khubunaya, S.A., Sobolev, A.V., 1998. Primary melts of calc-alkaline magnesian basalts from the Klyuchevskoi volcano, Kamchatka. *Doklady Akademii Nauk* 360, 100–102.
- Konrad-Schmolke, M., Halama, R., Manea, V.C., 2016. Slab mantle dehydrates beneath Kamchatka – Yet recycles water into the deep mantle. *Geochemistry Geophysics Geosystems* 17, 2987–3007.
- Kress, V., 1997. Magma mixing as a source for Pinatubo sulphur. *Nature* 389, 591–593.
- Langmuir, C.H., Vocke, R.D., Hanson, G.N., Hart, S.R. (1978). A general mixing equation with applications to Icelandic basalts. *Earth and Planetary Science Letters* 37, 380–392.

- Lassiter, J.C., Hauri, E.H., Nikogosian, I.K., Barszczus, H.G., 2002. Chlorine-potassium variations in melt inclusions from Raivavae and Rapa, Austral Islands: Constraints on chlorine recycling in the mantle and evidence for brine-induced melting of oceanic crust. *Earth and Planetary Science Letters* 202, 525–540.
- Leeman, W.P., Sisson, V.B., 1996. Geochemistry of boron and its implications for crustal and mantle processes, in: Grew, E.S., Anovitz, L.M. (Eds.), *Boron: Mineralogy, Petrology and Geochemistry*. Mineralogical Society of America 33, Washington, D.C., pp. 645–707.
- Leeman, W.P., Tonarini, S., Chan, L.H., Borg, L.E., 2004. Boron and lithium isotopic variations in a hot subduction zone—the southern Washington Cascades. *Chemical Geology* 212, 101–124.
- Le Voyer, M., Rose-Koga, E.F., Laubier, M., Schiano, P., 2008. Petrogenesis of arc lavas from the Rucu Pichincha and Pan de Azucar volcanoes (Ecuadorian arc): Major, trace element, and boron isotope evidences from olivine-hosted melt inclusions. *Geochemistry Geophysics Geosystems* 9, doi: 12010.11029/12008GC002173.
- Le Voyer, M., Cottrell, E., Kelley, K.A., Brounce, M., Hauri, E.H., 2015. The effect of primary versus secondary processes on the volatile content of MORB glasses: an example from the equatorial Mid-Atlantic Ridge (5°N–3°S). *Journal of Geophysical Research* 120, 2014JB011160.
- le Roux, P.J., Shirey, S.B., Hauri, E.H., Perfit, M.R., Bender, J.F., 2006. The effects of variable sources, processes and contaminants on the composition of northern EPR MORB (8–10°N and 12–14°N): Evidence from volatiles (H<sub>2</sub>O, CO<sub>2</sub>, S) and halogens (F, Cl). *Earth and Planetary Science Letters* 251, 209–231.
- Luhr, J.F., 1990. Experimental phase relations of water and sulfur saturated arc magmas and the 1982 eruptions of El Chichón volcano. *Journal of Petrology* 31, 1071–1114.

- Luhr, J.F., Carmichael, I.S.E., Varekamp, J.C., 1984. The 1982 eruptions of El Chichón Volcano, Chiapas, Mexico: mineralogy and petrology of the anhydrite-bearing pumices. *Journal of Volcanology and Geothermal Research* 23, 69–108.
- Macpherson, C.G., Matthey, D.P., 1998. Oxygen isotope variations in Lau Basin lavas. *Chemical Geology* 144, 177–194.
- Mandeville, C.W., Webster, J.D., Tappen, C., Taylor, B.E., Timbal, A., Sasaki, A., Hauri, E., Bacon, C.R., 2009. Stable isotope and petrologic evidence for open-system degassing during the climactic and pre-climactic eruptions of Mt. Mazama, Crater Lake, Oregon. *Geochimica et Cosmochimica Acta* 73, 2978–3012.
- Marini, L., Moretti, R., Accornero, M., 2011. Sulfur isotopes in magmatic-hydrothermal systems, melts, and magmas. *Reviews in Mineralogy and Geochemistry* 73, 423–492.
- Mathez, E.A., 1976. Sulfur solubility and magmatic sulfides in submarine basalt glasses. *Journal of Geophysical Research* 81, 4269–4276.
- Mavrogenes, J.A., O'Neill, H.S.C., 1999. The relative effects of pressure, temperature and oxygen fugacity on the solubility of sulfide in mafic magmas. *Geochimica et Cosmochimica Acta* 63, 1173–1180.
- Mavrogenes, J., Blundy, J., 2017. Crustal sequestration of magmatic sulfur dioxide. *Geology* 45, 211–214.
- Matjuschkin, V., Blundy, J.D., Brooker, R.A., 2016. The effect of pressure on sulphur speciation in mid- to deep-crustal arc magmas and implications for the formation of porphyry copper deposits." *Contributions to Mineralogy and Petrology* 171, 66.
- McCulloch, M.T., Gamble, J.A., 1991. Geochemical and geodynamical constraints on subduction zone magmatism. *Earth and Planetary Science Letters* 102, 358–374.
- Metrich, N., Clocchiatti R., 1996. Sulfur abundance and its speciation in oxidized alkaline melts. *Geochimica et Cosmochimica Acta* 60, 4151–4160.

- Métrich, N., Schiano, P., Clocchiatti, R., Maury, R.C., 1999. Transfer of sulfur in subduction settings: an example from Batan Island (Luzon volcanic arc, Philippines). *Earth and Planetary Science Letters* 167, 1–14.
- Michael, P.J., Cornell, W.C., 1998. Influence of spreading rate and magma supply on crystallisation and assimilation beneath mid-ocean ridges: Evidence from chlorine and major element chemistry of mid-ocean ridge basalts. *Journal of Geophysical Research* 103, 18325–18356.
- Michael, P., Schilling, J.-G., 1989. Chlorine in mid-ocean ridge magmas: evidence for assimilation of seawater-influenced components. *Geochimica et Cosmochimica Acta* 53, 3131–3143.
- Mironov, N., Portnyagin, M., Botcharnikov, R., Gurenko, A., Hoernle, K., Holtz, F., 2015. Quantification of the CO<sub>2</sub> budget and H<sub>2</sub>O–CO<sub>2</sub> systematics in subduction-zone magmas through the experimental hydration of melt inclusions in olivine at high H<sub>2</sub>O pressure. *Earth and Planetary Science Letters* 425, 1–11.
- Moore, L.R., Gazel, E., Tuohy, R., Lloyd, A.S., Esposito, R., Steele-MacInnis, M., Hauri, E.H., Wallace, P.J., Plank, T., Bodnar, R.J., 2015. Bubbles matter: An assessment of the contribution of vapor bubbles to melt inclusion volatile budgets. *American Mineralogist* 100, 806–823.
- Miyoshi, T., Sakai, H., Chiba, H., 1984. Experimental study of sulfur isotope fractionation factors between sulfate and sulfide in high temperature melts. *Geochemical Journal* 18, 75–84.
- Newman, S., Lowenstern, J.B., 2002. VolatileCalc: a silicate melt-H<sub>2</sub>O-CO<sub>2</sub> solution model written in Visual Basic for Excel. *Computers and Geosciences* 28, 597–604.
- Ovsyannikov, A.A., Muravyev, Y.D., 1992. Eruptions of the Chikurachki volcano. *Volcanology and Seismology* 5-6, 3–21 (in Russian).

- Peacock, S.M., Hervig, R.L., 1999. Boron isotopic composition of subduction-zone metamorphic rocks. *Chemical Geology* 160, 281–290.
- Pearce, J.A., 1983. Role of the sub-continental lithosphere in magma genesis at active continental margins, in: Hawkesworth, C.J., Norry, M.J. (Eds.), *Continental Basalts and Mantle Xenoliths*. Shiva Publishing, Nantwich, UK, pp. 230–249.
- Pearce, J.A., Peate, D.W., 1995. Tectonic implications of the composition of volcanic arc magmas. *Annual Review of Earth and Planetary Sciences* 24, 251–285.
- Plank, T., Langmuir, C.H., 1998. The chemical composition of subducting sediment and its consequences for the crust and mantle. *Chemical Geology* 145, 325–394.
- Portnyagin, M., Hoernle, K., Plechov, P., Mironov, N., Khubunaya, S., 2007. Constraints on mantle melting and composition and nature of slab components in volcanic arcs from volatiles (H<sub>2</sub>O, S, Cl, F) and trace elements in melt inclusions from the Kamchatka Arc. *Earth and Planetary Science Letters* 255, 53–69.
- Portnyagin, M., Duggen, S., Hauff, F., Mironov, N., Bindeman, I., Thirlwall, M., Hoernle, K., 2015. Geochemistry of the late Holocene rocks from the Tolbachik volcanic field, Kamchatka: quantitative modelling of subduction-related open magmatic systems. *Journal of Volcanology and Geothermal Research* 307, 133–155.
- Richet, P., Bottinga, Y., Javoy, M., 1977. A review of hydrogen, carbon, nitrogen, oxygen, sulfur and chlorine stable isotope fractionation among gaseous molecules. *Annual Review of Earth and Planetary Sciences* 5, 65–110.
- Robin, C., Eissen, J.P., Monzier, M., 1993. Giant tuff cone and 12-km-wide associated caldera at Ambrym Volcano (Vanuatu, New Hebrides Arc). *Journal of Volcanology and Geothermal Research* 55, 225–238.

- Roggensack, K., Hervig, R.L., McKnight, S.B., Williams, S.N., 1997. Explosive basaltic volcanism from Cerro Negro volcano: Influence of volatiles on eruptive style. *Science* 277, 1639–1642.
- Rogers, G., Saunders, A.D., Terrell, D.J., Verma, S.P., Marriner, G.F., 1985. Geochemistry of Holocene volcanic rocks associated with ridge subduction in Baja, California, Mexico. *Nature* 315, 389–392.
- Rose, E.F., Shimizu, N., Layne, G.D., Grove, T.L., 2001. Melt production beneath Mt. Shasta from boron data in primitive melt inclusions. *Science* 293, 281–283.
- Rosner, M., Erzinger, J., Franz, G., Trumbull, R.B., 2003. Slab-derived boron isotope signatures in arc volcanic rocks from the Central Andes and evidence for boron isotope fractionation during progressive slab dehydration. *Geochemistry Geophysics Geosystems* 4, doi: 0.1029/2002GC000438.
- Ruth, D.C.S., Cottrell, E., Cortés, J.A., Kelley, K.A., Calder, E.S., 2016. From passive degassing to violent strombolian eruption: the case of the 2008 eruption of Llaima volcano, Chile. *Journal of Petrology* 57, 1833–1864.
- Ryan, J.G., Langmuir, C.H., 1993. The systematics of boron abundances in young volcanic rocks. *Geochimica et Cosmochimica Acta* 57, 1489–1498.
- Ryan, J.G., Morris, J., Tera, F., Leeman, W.P., Tsvetkov, A., 1995. Cross-arc geochemical variations in the Kurile Arc as a function of slab depth. *Science* 270, 625–627.
- Saal, A.E., Hauri, E.H., Langmuir, C.H., Perfit, M.R., 2002. Vapour undersaturation in primitive mid-ocean-ridge basalt and the volatile content of Earth's upper mantle. *Nature* 419, 451–455.
- Sable, J.E., Houghton, B.F., Del Carlo, P., Coltelli, M., 2006. Changing conditions of magma ascent and fragmentation during the Etna 122 BC basaltic Plinian eruption: Evidence from clast microtextures. *Journal of Volcanology and Geothermal Research* 158, 333–354.

- Sakai, H., Casadevall, T.J., Moore, J.G., 1982. Chemistry and isotopic ratios of sulfur in basalts and volcanic gases at Kilauea volcano, Hawaii. *Geochimica et Cosmochimica Acta* 46, 729–738.
- Sakai, H., Des Marais, D.J., Ueda, A., Moore, J.G., 1984. Concentrations and isotope ratios of carbon, nitrogen and sulfur in ocean floor basalt. *Geochimica et Cosmochimica Acta* 48, 2433–2441.
- Salters, V.J.M., Stracke, A., 2004. Composition of the depleted mantle. *Geochemistry Geophysics Geosystems* 5, doi: 10.1029/2003GC000597.
- Savov, I.P., Ryan, J.G., D'Antonio, M., Kelley, K., Mattie, P., 2005. Geochemistry of serpentinized peridotites from the Mariana Forearc-Conical Seamount, ODP Leg 125: Implications for the elemental recycling at subduction zones. *Geochemistry Geophysics Geosystems* 6, Q04J15, doi: 10.1029/2004GC000777.
- Savov, I.P., Ryan, J.G., D'Antonio, M., Fryer, P., 2007. Shallow slab fluid release across and along the Mariana arc-basin system: Insights from geochemistry of serpentinized peridotites from the Mariana Forearc. *Journal of Geophysical Research* 112, doi: 10.1029/2006JB004749.
- Scambelluri, M., Tonarini, S., 2012. Boron isotope evidence for shallow fluid transfer across subduction zones by serpentinized mantle. *Geology* 40, 907–910.
- Selvaraja, V., Fiorentini, M.L., LaFlamme, C.K., Wing, B.A., Bui, T.-H., 2017. Anomalous sulfur isotopes trace volatile pathways in magmatic arcs. *Geology* 45, 419–422.
- Shaw, D.M., 1970. Trace element fractionation during anatexis. *Geochimica et Cosmochimica Acta* 34, 237–243.
- Shimizu, K., Shimizu, N., Komiya, T., Suzuki, K., Maruyama, S., Tatsumi, Y., 2009. CO<sub>2</sub>-rich komatiitic melt inclusions in Cr-spinels within beach sand from Gorgona Island, Colombia. *Earth and Planetary Science Letters* 288, 33–43.

- Simons, K., Dixon, J., Schilling, J.-G., Kingsley, R., Poreda, R., 2002. Volatiles in basaltic glasses from the Easter-Salas y Gomez Seamount Chain and Easter Microplate: Implications for geochemical cycling of volatile elements. *Geochemistry Geophysics Geosystems* 3, doi: 10.1029/2001GC000173.
- Shinohara, H., 2008. Excess degassing from volcanoes and its role on eruptive and intrusive activity. *Reviews of Geophysics* 46, RG4005, doi: 10.1029/2007RG000244.
- Smith, H.J., Leeman, W.P., Davidson, J., Spivack, A.J., 1997. The B isotopic composition of arc lavas from Martinique, Lesser Antilles. *Earth and Planetary Science Letters* 146, 303–314.
- Smythe, D.J., Wood, B.J., Kiseeva, E.S., 2017. The S content of silicate melts at sulfide saturation: New experiments and a model incorporating the effects of sulfide composition. *American Mineralogist* 102, 795–803.
- Spivack, A.J., Edmond, J.M., 1986. Determination of boron isotope ratios by thermal ionization mass spectrometry of the cesium metaborate cation. *Analytical Chemistry* 58, 31–35.
- Soto, G.J., Alvarado G.E., 2006. Eruptive history of Arenal Volcano, Costa Rica, 7 ka to present. *Journal of Volcanology and Geothermal Research* 157, 254–269.
- Stalder, R., Foley, S.F., Brey, G.P., Horn, I., 1998. Mineral-aqueous fluid partitioning of trace elements at 900–1200°C and 3.0–5.7 GPa: new experimental data for garnet, clinopyroxene, and rutile, and implications for mantle metasomatism. *Geochimica et Cosmochimica Acta* 62, 1781–1801.
- Staudigel, H., Plank, T., White, B., Schmincke, H.-U., 1996. Geochemical fluxes during seafloor alteration of the basaltic upper oceanic crust: DSDP Sites 417 and 418, in: Bebout, G.E., Scholl, D.W., Kirby, S.H., Platt, J.P. (Eds.), *Subduction: Top to Bottom*. Geophysical Monography 96, American Geophysical Union, pp. 19–38.

- Steele-Macinnis, M., Esposito, R., Bodnar, R.J., 2011. Thermodynamic model for the effect of post-entrapment crystallization on the H<sub>2</sub>O–CO<sub>2</sub> systematics of vapor-saturated, silicate melt Inclusions. *Journal of Petrology* 52, 2461–2482.
- Stern, C.R., Kilian, R., 1996. Role of the subducted slab, mantle wedge and continental crust in the generation of adakites from the Andean Austral Volcanic Zone. *Contribution to Mineralogy and Petrology* 123, 263–281.
- Stern, C.R., Futa, K., Muehlenbachs, K., 1984. Isotope and trace element data for orogenic andesites from the Austral Andes, in: Harmon, R.S., Barreiro, B.A. (Eds.), *Andean Magmatism: Chemical and Isotopic Constraints*. Shiva Publishing, Cheshire, UK, pp. 31–46.
- Straub, S.M., Layne G.D., 2002. The systematics of boron isotopes in Izu arc front volcanic rocks. *Earth and Planetary Science Letters* 198, 25–39.
- Straub, S.M., Layne, G.D., 2003. The systematics of chlorine, fluorine, and water in Izu arc front volcanic rocks: Implications for volatile recycling in subduction zones. *Geochimica et Cosmochimica Acta* 67, 4179–4203.
- Symonds, R.B., Rose, W.I., Bluth, G.J.S., Gerlach, T.M., 1994. Volcanic-gas studies: methods, results, and applications, in: Carroll, M.R., Holloway, J.R. (Eds.), *Volatiles in Magmas*. *Reviews in Mineralogy* 30, Mineralogical Society of America, Washington, D.C., pp 1–66.
- Szramek, L., Gardner, J.E., Larsen, J., 2006. Degassing and microlite crystallization of basaltic andesite magma erupting at Arenal Volcano, Costa Rica. *Journal of Volcanology and Geothermal Research* 157, 182–201.
- Takagi, T., Orihashi, Y., Naito, K., Watanabe, Y., 1999. Petrology of a mantle-derived rhyolite, Hokkaido, Japan. *Chemical Geology* 160, 425–445.
- Tatsumi, Y., Hamilton, D.L., Nesbitt, R.W., 1986. Chemical characteristics of fluid phase released from a subducted lithosphere and origin of arc magmas: evidence from high-

- pressure experiments and natural rocks. *Journal of Volcanology and Geothermal Research* 29, 293–309.
- Tatsumi, Y., Furukawa, Y., Kogiso, T., Yamada, Y., Yokoyama, T., Fedotov, S.A., 1994. A third volcanic chain in Kamchatka: Thermal anomaly at transform/convergence plate boundary. *Geophysical Research Letters* 21, 537–540.
- Taylor, B.E., 1986. Magmatic volatiles: isotopic variation of C, H, and S, in: Valley, J.W. Taylor, Jr., H.P., O'Neil, J.R. (Eds.), *Stable Isotopes in High Temperature Geological Processes*, *Reviews in Mineralogy* 16, Mineralogical Society of America, Washington, D.C., pp. 185–225.
- Thirlwall, M.F., Graham, A.M., Arculus, R.J., Harmon, R.S., Macpherson, C.G., 1996. Resolution of the effects of crustal assimilation, sediment subduction, and fluid transport in island arc magmas: Pb-Sr-Nd-O isotope geochemistry of Grenada, Lesser Antilles. *Geochimica et Cosmochimica Acta* 60, 4785–4810.
- Tonarini, S., Leeman, W.P., Leat, P.T., 2011. Subduction erosion of forearc mantle wedge implicated in the genesis of the South Sandwich Island (SSI) arc: Evidence from boron isotope systematics. *Earth and Planetary Science Letters* 301, 275–284.
- Turner, S., Hawkesworth, C., Rogers, N., Bartlett, J., Worthington, T., Hergt, J., Pearce, J.A., Smith, I., 1997.  $^{238}\text{U}$ - $^{230}\text{Th}$  disequilibria, magma petrogenesis, and flux rates beneath the depleted Tonga-Kermadec island arc. *Geochimica et Cosmochimica Acta* 61, 4855–4884.
- Varekamp, J.C., Luhr, J.F., Prestegard, K.L., 1984. The 1982 eruptions of El Chichón Volcano (Chiapas, Mexico): character of the eruptions, ash-fall deposits, and gas phase. *Journal of Volcanology and Geothermal Research* 23, 39–68.
- Vroon, P.Z., Lowry, D., van Bergen, M.J., Boyce, A.J., Matthey, D.P., 2001. Oxygen isotope systematics of the Banda Arc: Low  $\delta^{18}\text{O}$  despite involvement of subducted continental material in magma genesis. *Geochimica et Cosmochimica Acta* 65, 589–609.

- Walker, G.P.L., 1973. Explosive volcanic eruptions – a new classification scheme. *Geologische Rundschau* 62, 431–446.
- Walker, G.P.L., 1980. The Taupo pumice: product of the most powerful known (ultraplinian) eruption? *Journal of Volcanology and Geothermal Research* 8, 69–94.
- Walker, G.P.L., Self, S., Wilson, L., 1984. Tarawera 1886, New Zealand — A basaltic plinian fissure eruption. *Journal of Volcanology and Geothermal Research* 21, 61–78.
- Wallace, P.J., Anderson, A.T.J., 1998. Effects of eruption and lava drainback on the H<sub>2</sub>O contents of basaltic magmas at Kilauea Volcano. *Bulletin of Volcanology* 59, 327–344.
- Wallace, P., Carmichael, I.S.E., 1992. Sulfur in basaltic magmas. *Geochimica et Cosmochimica Acta* 56, 1863–1874.
- Wallace, P.J., Carmichael, I.S.E., 1994. S speciation in submarine basaltic glasses as determined by measurements of SK<sub>α</sub> X-ray wavelength shifts. *American Mineralogist* 79, 161–167.
- Wallace, P.J., Kamenetsky, V.S., Cervantes, P., 2015. Melt inclusion CO<sub>2</sub> contents, pressures of olivine crystallization, and the problem of shrinkage bubbles. *American Mineralogist* 100, 787–794.
- Walowski, K.J., Wallace, P.J., Clyne, M.A., Rasmussen, D.J., Weis, D., 2016. Slab melting and magma formation beneath the southern Cascade arc. *Earth and Planetary Science Letters* 446, 100–112.
- Wilson, L., Walker, G.P.L., 1987. Explosive volcanic eruptions – VI. Ejecta dispersal in plinian eruptions: the control of eruption conditions and atmospheric properties. *Geophysical Journal of the Royal Astronomical Society* 89, 657–679.
- Williams, S.N., 1983. Plinian airfall deposits of basaltic composition. *Geology* 11, 211–214.
- Woodhead, J., Eggins, S., Gamble, J., 1993. High field strength and transition element systematics in island arc and back-arc basin basalts: evidence for multi-phase melt extraction and a depleted mantle wedge. *Earth and Planetary Science Letters* 114, 491–504.

Wunder, B., Meixner, A., Romer, R.L., Wirth, R., Heinrich, W., 2005. The geochemical cycle of boron: Constraints from boron isotope partitioning experiments between mica and fluid. *Lithos* 84, 206–216.

Yogodzinski, G.M., Lees, J.M., Churikova, T.G., Dorendorf, F., Wörner, G., Volynets, O.N., 2001. Geochemical evidence for the melting of subducting oceanic lithosphere at plate edges. *Nature* 409, 500–504.

ACCEPTED MANUSCRIPT

**Figure captions**

**Fig. 1.** (A) Schematic map of the northern Kurile Islands (Russia) with the locality of the Chikurachki volcano on Paramushir Island shown in relation to the Kamchatka Peninsula. Contours show isobars for the top of the Wadati-Benioff zone from Tatsumi et al. (1994). (B) The Chikurachki volcano, a view from the eastern side ~6 km away.

**Fig. 2.** Isopachs (lines of equal thickness) given in cm of the tephra representing Plinian (A) 1853 and (B) 1986 eruptions of the Chikurachki volcano after Belousov et al. (2003).

**Fig. 3.** Multi-element diagram presenting trace and rare-earth element distribution in olivine-hosted glass inclusions and groundmass glasses and demonstrating positive H<sub>2</sub>O, B, Cl and F enrichment. The concentrations of trace elements in N-MORB used for normalisation are from Hofmann (1988), the normalisation values for volatile elements and water are 1350 µg/g H<sub>2</sub>O, 0.6 µg/g B, 54.5 µg/g Cl, and 146 µg/g F, which were calculated from the average B/K ≈ 0.001, H<sub>2</sub>O/Ce ≈ 180, K/Cl ≈ 11, and F/Nd ≈ 20, or F/Zr ≈ 2 ratios inferred for MORB using the data of Ryan and Langmuir (1993), Jambon et al. (1995), Saal et al. (2002), Kendrick et al. (2013), Le Voyer et al. (2015) and references therein. The shaded field presents whole-rock data for the Kurile island arc basalts and andesites (MgO > 2.5 wt.%) from Bailey et al. (1987a,b), Ishikawa and Tera (1997), Ikeda (1998), Ikeda et al. (2000), Takagi et al. (1999), Ishikawa et al. (2001).

**Fig. 4.** Concentrations of CO<sub>2</sub> and H<sub>2</sub>O dissolved in glass inclusions measured by SIMS. (A) CO<sub>2</sub> and H<sub>2</sub>O corrected for the amount of host mineral crystallised inside melt inclusions between entrapment and magma eruption and final quenching. (B) CO<sub>2</sub> and H<sub>2</sub>O corrected for the loss of volatiles to the inclusion shrinkage bubble (see text). Isobars of melt compositions in equilibrium with CO<sub>2</sub>-H<sub>2</sub>O fluid (*solid lines*) and isopleths of fluid composition (*dashed lines*)

with labels returning molecular fraction of H<sub>2</sub>O in the fluid) were calculated using *VolatileCalc* solution model (Newman and Lowenstern, 2002). The isopleths and isobars were calculated for basaltic magma composition at 1000°C i.e., the average temperature of the Chikurachki magmas (Gurenko et al., 2005a). *Error bars* represent  $\pm 2\sigma$  average integral analytical uncertainty, i.e.,  $\pm 18\%$  relative for CO<sub>2</sub> and  $\pm 12\%$  relative for H<sub>2</sub>O, they are less than the size of a symbol, if not shown.

**Fig. 5.** Concentrations of S, Cl and F in glass inclusions and lapilli matrix glasses. (A) FeO vs. S variations. The larger shaded grey symbols denote the compositions of the respective host matrix glasses. The *MORB and OIB* shaded field presents data for presumably undegassed MORB and Hawaiian submarine tholeiitic and alkali basaltic glasses from Wallace and Carmichael (1992). The *S saturation* shaded field represents compositional dependence of sulphur saturation of MORB-type magmas with immiscible sulfide liquid of pure FeS based on experimental data of Smythe et al. (2017) and references therein. Legend: *MI* = melt inclusions, *GL* = matrix glasses. (B) Ce vs. S, (C) H<sub>2</sub>O/K<sub>2</sub>O vs. S/K<sub>2</sub>O, (D) F/K<sub>2</sub>O vs. S/K<sub>2</sub>O, (E) H<sub>2</sub>O/K<sub>2</sub>O vs. Cl/K<sub>2</sub>O, and (F) F/Zr vs. Cl/Nb. The observed relationships do not support shallow-depth contamination of the magma, and rather favour a deep origin of volatiles resulted from slab-derived fluid released during subduction (see text).

**Fig. 6.** Sulphur isotope composition of glass inclusion and one host matrix glass of the Chikurachki prehistoric eruption related to (A) S concentrations, (B) H<sub>2</sub>O/K<sub>2</sub>O and (C) Cl/K<sub>2</sub>O ratios.

**Fig. 7.** Diagrams of (A) H<sub>2</sub>O/K<sub>2</sub>O vs. Cl/K<sub>2</sub>O, (B) H<sub>2</sub>O/Ce vs. Cl/Nb, (C) H<sub>2</sub>O/K<sub>2</sub>O vs. B/K<sub>2</sub>O, and (D) H<sub>2</sub>O/Ce vs. B/Nb (given as weight ratios) illustrate the effects of contamination (mixing lines) of presumably “uncontaminated” arc magma (*UAM*) with different types of contaminants,

i.e., seawater (*SW*), 15%- and 50%-NaCl saline brines (labelled as *15%-NaCl* and *50%-NaCl*, respectively), altered oceanic crust (*AOC1* and *AOC2*) and siliceous marine sediment (*SED*) (see text and **Table 1** for the end-member chemical compositions and further details). Numbers on the mixing lines refer to wt.% of the contaminant added to the magma. The calculated mixing trends allow us to conclude that direct contamination of UAM by altered basaltic rocks and marine sediments at shallow depth and/or by seawater or seawater-derived components (saline brine) is unable to account for the H<sub>2</sub>O, Cl and B enrichment observed in the studied melt inclusions. Legend: *MI* = melt inclusions, *GL* = matrix glasses.

**Fig. 8.** Boron isotope composition of the studied glass inclusions given in relation to B/K<sub>2</sub>O (A) and B/Nb (B) in context of possible magma contamination. The mixing end-members used (and their labels) are the same as in **Fig. 7**. It is demonstrated that magma contamination by seawater or seawater-derived components, altered oceanic crust and marine sediments is unable to account for B isotopic composition of the studied melt inclusions.

**Fig. 9.** Sulphur isotope fractionation during magma degassing. The  $\delta^{34}\text{S}$  values in the melt inclusions are shown versus fraction of S remaining (*F*). Shifts of  $\delta^{34}\text{S}$  were calculated assuming closed-system (equilibrium, solid and dashed *straight lines*) and open-system (Rayleigh distillation; solid and dashed *curves*) degassing (see text), the  $\alpha_{\text{gas-melt}}$  values of 0.996 and 0.998, and maximum, undegassed S concentration found in the most S-rich melt inclusions having the highest  $\delta^{34}\text{S}$  values. At the temperature range from 910 to 1180°C and the redox conditions of NNO+1 to NNO+2 obtained for the studied Chikurachki magmas, the outgassed SO<sub>2</sub> will drive out the lighter <sup>32</sup>S isotope so that the remaining melt will be progressively enriched in <sup>34</sup>S. Thus, degassing alone cannot explain the observed, ~10‰-S-isotope of the studied melt inclusions.

**Fig. 10.** Diagram Th/Nb vs. Ba/Nb illustrating a possible origin of the Chikurachki magmas as a result of melting of a three-component magma source. Legend: *MI* = melt inclusions, *GL* = matrix glasses. The source components are mantle wedge (*MW*) with Ba, Nb and Th concentrations of the depleted mantle after Salters and Stracke (2004) and hybrid subduction component (*SUBD*) representing a binary 4:6-mixture of a sediment melt (*SedMelt*) with a composite slab-derived fluid, which in turn is a 9:1-mixture of a dehydration fluids released from the altered oceanic crust (*AOCFluid*) and Kurile sediment (*SedFluid*) (see text and **Table 2** for definition). The compositions of *Kurile lavas* are from Bailey et al. (1987a,b), Ishikawa and Tera (1997), Ikeda (1998), Ikeda et al. (2000), Takagi et al. (1999), Ishikawa et al. (2001). *KurileSed* is a bulk composition of the sediment column subducted at the Kurile trench taken from Plank and Langmuir (1998). *AOC* is altered oceanic crust taken from Staudigel et al. (1996), whose composition was used by Hochstaedter et al. (2001) to calculate *AOCFluid* composition. *SlabMelt* is a composition of composite hydrous melt released from the slab and contributed to the origin of magmas erupted in the Tolbachik volcanic field, Kamchatka, which was taken from Portnyagin et al. (2015) for comparison.

**Fig. 11.** Diagrams of (A)  $\delta^{11}\text{B}$  vs. B/Nb and (B)  $\delta^{11}\text{B}$  vs. Ba/Nb illustrate the proposed interpretation to account for B-isotope variations in the Chikurachki mineral-hosted melt inclusions by addition of slab-derived fluids to the mantle wedge. Mixing lines between the depleted MORB-type mantle wedge (*MW*) and subduction slab fluids (*SSF1*, *SSF2* and *SSF3*) are shown (see **Table 1** for chemical composition). Numbers on mixing lines refer to wt% of component added to MW end-member.




Fig. 1. Gurenko et al.

ACCEPTED



**(A) 1853 eruption****(B) 1986 eruption**

Fig. 2. Gurenko et al.




Fig. 3. Gurenko et al.

ACCEPTED MANUSCRIPT




Fig. 4. Gurenko et al.




Fig. 5. Gurenko et al.




Fig. 6. Gurenko et al.




Fig. 7. Gurenko et al.




Fig. 8. Gurenko et al.




Fig. 9. Gurenko et al.




Fig. 10. Gurenko et al.

ACCEPTED




Fig. 11. Gurenko et al.